

60ml when reconstituted

MoxipenTM

Amoxycillin Trihydrate

FOR ORAL SUSPENSION

Amoxycillin 125mg / 5ml

To reconstitute: Shake to loosen powder. Add water and shake vigorously to disperse powder. Add water to the 60ml mark and shake well.

SHAKE WELL BEFORE EACH DOSE

After reconstitution, store in a cool, dry place below 25°C

Use within 5 days after reconstitution.

DOSE: 8 hourly

Infants/Children:

up to 6kg

6-8kg

8-20kg

Above 20kg

Adults:

25mg-50mg

50mg-100mg

6.7mg-13.3mg per kg body weight

250mg-500mg

250mg-500mg

Preservative: Sodium Benzoate 0.1% w/v.

Store below 25°C.

Not to be used in patients with known hypersensitivity to Penicillin.

Refer to enclosed leaflet for further information.

CONTROLLED MEDICINE

FOR ORAL USE ONLY

KEEP OUT OF REACH OF CHILDREN / JAUHI DARI KANAK-KANAK

BATCH MFG EXP



Xepa^{SP}

Manufactured and Marketed by :

Xepa-Soul Pattinson (Malaysia) Sdn Bhd

1-5 Cheng Industrial Estate, 75250 Melaka, Malaysia

CODE M00200

MAL 19900374AZ

SIN 7725P

HK-36163

MM Reg: No. R2002A3783