

90ml

Each 5ml contains:

Dextromethorphan HBr 15mg
Triprolidine HCl 1.25mg
Pseudoephedrine HCl 30mg
Sodium Benzoate 0.1% w/v as preservative.

Store in a cool place below 30°C.

Refer to accompanying package insert for further information.

MAL 19940070AZ CODE C00505
SIN 8780P BRU15111967P
HK-40687

CONTROLLED MEDICINE

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

儲存于兒童的活動範圍以外。



Manufactured and Marketed by

Xepa-Soul Pattinson (Malaysia) Sdn Bhd

1-5, Cheng Industrial Estate, 75250 Melaka, Malaysia

Cough-en[®] Linctus

**Antitussive, Antihistamine,
Nasal Decongestant**

Relieves

- Dry Irritating Cough
- Runny Nose
- Sneezing
- Stuffy Nose
- Itchy & Watery Eyes

Blackcurrant flavour

Dosage:

- Adults & children over 12 years
10ml 3 times a day
- Children 6 to 12 years
5ml 3 times a day
- Children 2 to 5 years
2.5ml 3 times a day

*Sip and swallow slowly,
preferably taken
without dilution.*

Caution: This medicine may cause drowsiness. If affected, do not drive or operate machinery. Avoid alcoholic drinks.

Warnings: Not to be used in children below 2 years old. To be used with cautions, and as advised by the physician/pharmacist for children age 2 to 6 years.

C00505

BATCH

MFG EXP

004

120ml

Each 5ml contains:

Dextromethorphan HBr 15mg
Triprolidine HCl 1.25mg
Pseudoephedrine HCl 30mg
Sodium Benzoate 0.1% w/v as preservative.

Store in a cool place below 30 °C.

Refer to accompanying package insert for further information.

MAL 19940070AZ

CODE C00504

HK-40687

BRU15111967P

SIN 8780P

CONTROLLED MEDICINE

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

儲存于兒童的活動範圍以外。

FOR ORAL USE ONLY



Xepa^{sp}

Manufactured and Marketed by

Xepa-Soul Pattinson (Malaysia) Sdn Bhd

1-5, Cheng Industrial Estate, 75250 Melaka, Malaysia

Cough-en[®] Linctus

Antitussive, Antihistamine,
Nasal Decongestant

Relieves

- Dry Irritating Cough
- Runny Nose
- Sneezing
- Stuffy Nose
- Itchy & Watery Eyes

Blackcurrant flavour

Dosage:

- Adults & children over 12 years
10ml 3 times a day
- Children 6 to 12 years
5ml 3 times a day
- Children 2 to 5 years
2.5ml 3 times a day

*Sip and swallow slowly,
preferably taken
without dilution.*

Caution: This medicine may
cause drowsiness. If affected,
do not drive or operate machinery.
Avoid alcoholic drinks.

Warnings: Not to be used in children
below 2 years old. To be used with
cautions, and as advised by the
physician/pharmacist for children
age 2 to 6 years.

C00504

BATCH
MFG EXP

005