

PBX0128



12JFMAMJJAASOND171819



0318



Febricol®

Each tablet contains:

Paracetamol 500mg

Pseudoephedrine HCl 30mg


Chlorpheniramine Maleate 2mg

Dosage: Adults and children over 12 years : 1-2 tablets every 4-6 hours. Maximum 8 tablets per day.

See enclosed leaflet for additional information.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.
Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/Toxic Epidermal Necrolysis have been reported.

Febricol®



Febricol®


Relieves symptoms of headache and nasal congestion in influenza and common cold

100 TABLETS



004

Febricol®

BN
MPC
EXP

Caution : This may cause drowsiness. If affected, do not drive or operate machinery.
注意：此藥可使人昏昏欲睡。服後如有此情形。不得駕駛或動用機械。

Caution : It is dangerous to exceed the stated dose.
注意：服食過量有危險。

This preparation contains PARACETAMOL. Do not take any other paracetamol-containing medicines at the same time
There is potential for liver toxicity and severe liver damage. Do not take more than the recommended dose.
含有 Paracetamol，不能同時服用其他含有 Paracetamol 藥方。
可能引起肝臟致毒和損傷。不能超過指定服量。

Storage: Store in cool dry place below 30°C
儲存於攝氏30度以下。**CONTROLLED MEDICINE**

KEEP OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

FOR ORAL USE ONLY



Manufactured and Marketed by
Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5 Cheng Industrial Estate
75250 Melaka, Malaysia

MAL 20013375AZ
SIN 11488P
HK-50347
CODE FE0105

9 556492 000857

Febricol®

Melegakan sakit kepala & kesesakan hidung
semasa demam selsema & influenza

100 TABLETS

