

Glenix®
Glimepiride 2mg

Xepa^{sp}

HOLOGRAM

Glenix®
Glimepiride 2mg

30 Tablets
Each tablet contains Glimepiride 2mg

Glenix®
Glimepiride 2mg

Glenix®
Glimepiride 2mg

Dosage: Take immediately before breakfast.

Initially 1mg once daily, adjust the dose according to response in 1mg steps at 1-2 weeks intervals; or as prescribed by the physician.

Maximum dosage : 6mg daily.

Dosis: Ambil sebelum sarapan pagi.

Ambil 1mg sekali sehari pada mulanya, dan sesuaikan dosis pada kadar 1mg setiap 1-2 minggu berdasarkan respon kepada ubat; atau mengikut preskripsi doktor.

Dosis maksimum : 6mg sehari.

服量：早餐前即時服用。

每日服用1次，每次1mg，可依據對藥物的反應調節服量（每1-2星期調節1mg）；或遵照醫生指示服用。

每日總服量不可超過6mg。

Please refer to enclosed leaflet for details of dosage & administration, indications and other information.

Caution: It is dangerous to exceed the stated dose.

注意：服量過量有危險。

Store below 25°C.

Simpan di bawah 25°C.

置放在溫度 25°C 以下。

CONTROLLED MEDICINE

KEEP OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

儲存于兒童的活動範圍以外。

FOR ORAL USE ONLY

CODE GL0200 HK-58053

MAL 07021423A



9 556492 002455

Xepa^{sp}

Manufactured by

Xepa-Soul Pattinson (Malaysia) Sdn. Bhd.

1-5, Cheng Industrial Estate,

75250 Melaka, Malaysia.