

# PROGESIC SUSPENSION

Paracetamol (250mg/5ml)

## What is in this leaflet

1. What Progesic Suspension is used for
2. How it works
3. Before you use Progesic Suspension
4. How to use Progesic Suspension
5. While you are using Progesic Suspension
6. Side Effects
7. Storage and Disposal of Progesic Suspension
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

## What Progesic Suspension is used for

It is used to relieve:

- Fever.
- Pain, such as toothache and headache.

## How Progesic Suspension works

Paracetamol is a pain killer and fever reducer.

It works by reducing the production of chemicals that cause inflammation, pain, and fever.

## Before you use Progesic Suspension

### - When you must not use it

Do not use Progesic Suspension if you are allergic to Paracetamol or any ingredient in this medicine.

### - Before you start to use it

Tell your doctor or pharmacist before using Progesic Suspension if you:

- are pregnant or plan to pregnant.
- are breast-feeding or plan to start breast-feeding.
- have kidney or liver disease.

## WARNING

This preparation contains  
PARACETAMOL.

Do not take any other paracetamol containing medicines at the same time.

- **Allergy alert:** Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

### - Taking other medicines

Tell your doctor or pharmacist if you are taking or having recently taken any other

medicines. This includes any medicines which you have bought without a prescription especially coumarin medicine (blood thinning agent).

## How to use Progesic Suspension

### - How much to use

- **Adults:** 10ml- 20ml every 4-6 hours. Maximum 80ml in a day.
- **Children:**  
1-5 years: 2.5ml -5ml every 4-6 hours.  
6-12 years: 5ml -10ml every 4-6 hours.

### - When to use it

To be taken with or without food.

Follows the dose prescribed by doctor or pharmacist.

### - How long to use it

Follow the prescription from your doctor or pharmacist. Ask your doctor or pharmacist before prolonged use of this medicine.

### - If you forget to use it

Use it as soon as you remember, unless it is nearly time for your next dose, skip the missed dose. Then go on as before. Never double up on the next dose to make up for the one missed.

### - If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Over dosage may cause nausea, vomiting, abdominal pain, severe liver damage and sometimes kidney damage, coma and may leads to death.

## While you are using it

### - Things you must do

- Always wash your hand before handle the medicine.
- Take the medicine at the same time each day.

### - Things you must not do

Do not exceed the stated dose and do not give this product to anyone else, even if they have same symptoms or condition as you

### - Things to be careful of

Tell your doctor or pharmacist if fever does not subside in 36 hours, consult your doctor or pharmacist.

## Side effects

Progesic Suspension is usually well tolerated. However, if rashes occur, treatment should be discontinued.

Seek immediate medical attention if you experience any side effects or allergic reactions.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my) [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

## Storage and Disposal of Progesic Suspension

### - Storage

Store below 30°C. Keep container well closed.

Keep out of the reach of the children.

Do not use this medicine after expiry date.

### - Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required.

## Product Description

### - What it looks like

Orange colour suspension with raspberry and cherry flavour.

Available in pack of 100ml.

### - Ingredients

- Active Ingredient(s)  
Paracetamol

- Inactive ingredients  
Cherry Flavour, Sunset Yellow, Aluminium Magnesium Silicate, Syrup, Raspberry Flavour, Methyl Hydroxybenzoate, Propyl Hydroxybenzoate, Xanthan Gum, Purified Water, Propylene Glycol, Saccharin Sodium.

### - MAL number:

MAL19912891XZ

# PROGESIC SUSPENSION

Paracetamol (250mg/5ml)

---

## **Manufacturer and Product Registration**

### **Holder**

Xepa-Soul Pattinson (Malaysia) Sdn Bhd  
1-5 Cheng Industrial Estate,  
75250 Melaka, Malaysia.

### **Date of revision**

09/06/2020

### **Serial Number**

NPRA(R4/1)130520/00077