

Important information. Please read carefully.

# Bacidin<sup>TM</sup>

## Antiseptic Cream 1%

**Bacidin<sup>TM</sup>** contains an antibacterial agent that is useful for:

1. Healing of cuts
2. Burns
3. Abrasions
4. Minor skin infections

### COMPOSITION

Contains Chlorhexidine Gluconate 1% w/w in white, smooth, homogeneous cream. Methyl Hydroxybenzoate 0.15% w/w and Propyl Hydroxybenzoate 0.08% w/w as preservatives.

### USES

**Bacidin<sup>TM</sup>** Antiseptic Cream is used as a disinfectant for wounds, burns, abrasions and minor skin infections.

### DIRECTION

Apply to the affected area as required.

### CONTRAINDICATIONS

**Bacidin<sup>TM</sup>** Antiseptic Cream should not be used on the brain, meninges or perforated ear-drum.

### SIDE EFFECTS

Occasionally, skin sensitivity may occur

#### Immune system disorders

Frequency not known: Hypersensitivity including anaphylactic shock

### WARNINGS AND PRECAUTIONS

Do not use on sensitive tissues. Do not use with soap.

#### Allergy Alert

Bacidin contains chlorhexidine. Chlorhexidine is known to induce hypersensitivity, including generalised allergic reactions and anaphylactic shock. The prevalence of chlorhexidine hypersensitivity is unknown, but available literature suggests this is likely to be very rare. Bacidin should not be administered to anyone with a possible history of an allergic reaction to chlorhexidine.

### 消毒藥膏

**Bacidin<sup>TM</sup>** 含有一種抗菌藥物，適于：

1. 愈合割傷
2. 燒傷
3. 擦傷
4. 輕微被細菌感染的肌膚

### 成份

含 Chlorhexidine Gluconate 1%w/w 白色，柔質均勻軟膏。Methyl Hydroxybenzoate 0.15% w/w 和 Propyl Hydroxybenzoate 0.08% w/w 為防腐劑。

### 適應症

對一般傷口，燒傷，擦傷與輕微皮膚細菌感染有消毒功效。

### 用法

在需要處搽用。

### 禁忌症

勿用在頭腦，腦膜或受傷之耳膜。

### 副作用

偶爾可能造成皮膚敏感。

### 免疫系統紊亂

頻率不詳：過敏症，包括過敏性休克。

### 警語

勿塗在敏感地方，勿與肥皂一起使用。

**Bacidin** 含 chlorhexidine. Chlorhexidine 能導致過敏症，包括過敏性休克。對 chlorhexidine 過敏的頻率不詳，但極可能很罕見。**Bacidin** 不應給與任何可能對 chlorhexidine 的患者使用。



• Insect Bites • Gigitan Serangga  
• 蚊蟲咬



• Cuts • Luka • 割傷



• Abrasions • Terparut • 擦傷



• Burns • Terkena Api • 燒傷

If any signs or symptoms of suspected hypersensitivity reaction such as itching, skin rash, hives, redness, swelling (including facial swelling), breathing difficulties/ wheezing, shock, light headedness, and rapid heart rate develop, immediately stop using the product and seek medical help right away. Appropriate therapeutic countermeasures must be instituted as clinically indicated.

#### **SYMPTOMS & TREATMENT OF OVERDOSAGE**

Overdosage is rare. Chlorhexidine is poorly absorbed and systemic effects are unlikely even if large amounts are swallowed. However, gastric lavage followed by supportive measures may be used as appropriate.

#### **PRESENTATION**

White, smooth, homogeneous cream in collapsible aluminium tube of 15g and 10g.

#### **SHELF LIFE**

The expiry date is indicated on the packaging.

#### **STORAGE**

Store below 30°C.

#### **KEEP OUT OF REACH OF CHILDREN**

#### **JAUHI DARI KANAK-KANAK**

*For further information, please consult your pharmacist or physician.*

Revision Date: 08-Oct-2018

Manufactured and Marketed by  
Xepa-Soul Pattinson (Malaysia) Sdn Bhd  
1-5 Cheng Industrial Estate, 75250 Melaka, Malaysia.

若發現過敏症的症狀如搔癢，皮膚紅腫(包括面部腫)，麻疹，呼吸困難/喘息，休克，頭暈，及心率增加則應立即停藥,尋求醫療幫助。根據臨床指征採取適當的治療措施。

#### **用藥過量症狀與治療**

服用藥過量是罕見的。Clorhexidine 不易被吸收，並且在大量吞服後也不大可能有全身效應。但是，可以適當地使用洗胃接著支持治療。

#### **包裝**

白色，柔質，均勻軟膏，包裝在鋁質管。

#### **貯存期限**

有效日期备注于包装外。

#### **貯存**

存放在攝氏 30 度以下，陰涼及避免日晒的地方。把容器蓋好。

將此藥物放在兒童不能觸及的地方。

欲知其它資料，請向藥劑師或醫生詢問。