

90ml

Each 5ml contains:

Carbocisteine 100mg
Promethazine Hydrochloride 2.5mg

Methyl Hydroxybenzoate 0.1%w/v and
Propyl Hydroxybenzoate 0.024%w/v as preservative.
Store below 25°C.

Refer package insert for details.

Cautions: It is dangerous to exceed the stated dose. This medicine may cause drowsiness. If affected, do not drive or operate machinery.

CODE MU0500 • FOR ORAL USE ONLY
MAL06071081AZ HK-58930

CONTROLLED MEDICINE
KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK
储存于儿童的活动范围以外。



Manufacturer and Product Registration Holder:

Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5, Cheng Industrial Estate
75250 Melaka, Malaysia

Mucoprom[®] Syrup

Mucolytic, Antihistamine

Alcohol-Free & Sugar-Free
Chocolate vanilla flavour

Dosage: To be taken in divided doses, preferably in the evening.

- Adults
15 to 20ml every 6 to 8 hours
not to exceed 60ml in 24 hours
- Children 12-15 years
10 to 15ml every 6 to 8 hours
not to exceed 45ml in 24 hours
- Children 6-12 years
5 to 10ml every 6 to 8 hours
not to exceed 30ml in 24 hours
- Children 2-5 years
5ml every 6 to 8 hours
not to exceed 20ml in 24 hours

WARNING

When used for treatment of cough and cold
(a) It should not be used in pediatric patients less than 2 years of age because of the potential for fatal respiratory depression.
(b) To be used with caution and doctor's/pharmacist's advice in children 2 to 6 years of age.

Mucoprom may cause severe allergy and serious skin reactions. Stop using it and seek medical assistance immediately if you experience any of the following symptoms:

- Severe allergy: breathing difficulties, light headedness, skin swellings or rash.
- Severe skin reaction: skin reddening, blisters, rash, fever, sore throat or eye irritation.

MU0500

BATCH
MFG EXP

005

120ml

Each 5ml contains :

Carbocisteine 100mg

Promethazine Hydrochloride 2.5mg

Methyl Hydroxybenzoate 0.1%w/v and

Propyl Hydroxybenzoate 0.024%w/v as preservative.

Store below 25°C.

Refer package insert for details.

Cautions: It is dangerous to exceed the stated dose. This medicine may cause drowsiness. If affected, do not drive or operate machinery.

CODE MU0501 FOR ORAL USE ONLY

MAL06071081AZ HK-58930

CONTROLLED MEDICINE

KEEP OUT OF REACH OF CHILDREN

JAUH! DARI KANAK-KANAK

储存于儿童的活动范围以外。



Kepa^{SP}

Manufacturer and Product Registration Holder:

Xepa-Soul Pattinson (Malaysia) Sdn Bhd

1-5, Cheng Industrial Estate

75250 Melaka, Malaysia

Mucoprom[®] Syrup

Mucolytic, Antihistamine

Alcohol-Free & Sugar-Free

Chocolate vanilla flavour

Dosage : To be taken in divided doses, preferably in the evening.

- Adults
15 to 20ml every 6 to 8 hours
not to exceed 60ml in 24 hours
- Children 12-15 years
10 to 15ml every 6 to 8 hours
not to exceed 45ml in 24 hours
- Children 6-12 years
5 to 10ml every 6 to 8 hours
not to exceed 30ml in 24 hours
- Children 2-5 years
5ml every 6 to 8 hours
not to exceed 20ml in 24 hours

WARNING

When used for treatment of cough and cold

(a) It should not be used in pediatric patients less than 2 years of age because of the potential for fatal respiratory depression.

(b) To be used with caution and doctor's / pharmacist's advice in children 2 to 6 years of age.

Mucoprom may cause severe allergy and serious skin reactions. Stop using it and seek medical assistance immediately if you experience any of the following symptoms:

- Severe allergy: breathing difficulties, light headedness, skin swellings or rash.
- Severe skin reaction: skin reddening, blisters, rash, fever, sore throat or eye irritation.

MU0501

BATCH
MFG EXP

005