



Alcohol-Free & Sugar-Free

Mucolytic, Antihistamine

Mucoprom[®]
Syrup

90ml



Mucoprom[®]
Syrup

Mucolytic,
Antihistamine

Alcohol-Free & Sugar-Free

Chocolate vanilla flavour

4

005

Mucoprom[®]
Syrup



Free dosage cup & dosage
indication enclosed

Each 5ml contains:

Carbocisteine100mg

Promethazine Hydrochloride..... 2.5mg

Methyl Hydroxybenzoate 0.1% w/v and

Propyl Hydroxybenzoate 0.024% w/v as preservatives.

WARNING

When used for treatment of cough and cold

(a) It should not be used in pediatric patients less than 2 years

of age because of the potential for fatal respiratory depression.

(b) To be used with caution and doctor's/ pharmacist's advice

in children 2 to 6 years of age.

Mucoprom may cause severe allergy and serious skin reactions.

Stop using it and seek medical assistance immediately if you

experience any of the following symptoms:

- Severe allergy: breathing difficulties, light headedness, skin

swellings or rash.

- Severe skin reaction: skin reddening, blisters, rash, fever,

sore throat or eye irritation.

Store below 25°C.

Refer package insert for details.



9 556492 002424

Cautions : It is dangerous to exceed
the stated dose. This medicine may
cause drowsiness. If affected, do not
drive or operate machinery.

90ml



Mucoprom[®]
Syrup

Name/ Nama:
姓名:

Date/ Tarikh:
日期:

ml/ 毫升

Times Daily/ Kali Sehari/ 次

B
MFG
EXP



Manufacturer and Product Registration Holder
Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5, Cheng Industrial Estate
75250 Melaka, Malaysia



Mucoprom[®]
Syrup



Cawan penyukat dos
percuma
免费度量器

Simpan di bawah suhu 25°C.

Amaran : Adalah berbahaya jika melebihi
dos yang dinyatakan.

Ubat ini mungkin menyebabkan
mengantuk. Jika ada kesannya,
jangan memandu kenderaan
atau mengendalikan mesin.

置放在25°C以下。

注意： 服食过量有危险。此药可使人昏昏欲睡，
服食后如有此情况，不得驾驶或动用机械。

FOR ORAL USE ONLY

CONTROLLED MEDICINE

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

储存于儿童的活动范围以外。

CODE MU0500

MAL 06071081AZ

HK-58930





Alcohol-free & Sugar-free

Mucolytic, Antihistamine
Mucoprom[®]
Syrup

120ml



4

004

Mucoprom[®]
Syrup

Mucolytic,
Antihistamine

Alcohol-Free & Sugar-Free
Chocolate vanilla flavour

Mucoprom[®]
Syrup

Dosage: To be taken in divided doses, preferably in the evening.

- Adults
15 to 20ml every 6 to 8 hours
not to exceed 60ml in 24 hours
- Children 12-15 years
10 to 15ml every 6 to 8 hours
not to exceed 45ml in 24 hours
- Children 6-12 years
5 to 10ml every 6 to 8 hours
not to exceed 30ml in 24 hours
- Children 2-5 years
5ml every 6 to 8 hours
not to exceed 20ml in 24 hours

Each 5ml contains:
Carbocisteine 100mg
Promethazine Hydrochloride 2.5mg
Methyl Hydroxybenzoate 0.1% w/v and
Propyl Hydroxybenzoate 0.024% w/v as preservatives.

WARNING

When used for treatment of cough and cold
(a) It should not be used in pediatric patients less than 2 years of age because of the potential for fatal respiratory depression.
(b) To be used with caution and doctor's/ pharmacist's advice in children 2 to 6 years of age.

Mucoprom may cause severe allergy and serious skin reactions. Stop using it and seek medical assistance immediately if you experience any of the following symptoms:

- Severe allergy: breathing difficulties, light headedness, skin swellings or rash.
- Severe skin reaction: skin reddening, blisters, rash, fever, sore throat or eye irritation.

Store below 25°C.

Refer package insert for details.




9 556492 002431

Cautions: It is dangerous to exceed the stated dose. This medicine may cause drowsiness. If affected, do not drive or operate machinery.



Free dosage cup & dosage indication enclosed

120ml



Mucoprom[®]
Syrup

Mucolytic,
Antihistamine

Alcohol-Free & Sugar-Free
Chocolate vanilla flavour

Mucoprom[®]
Syrup

Dos: Untuk diambil dalam dos berasingan, disyorkan pada waktu lewat petang.

- Dewasa
15 hingga 20ml setiap 6 hingga 8 jam
tidak melebihi 60ml dalam tempoh 24 jam
- Kanak-kanak 12 hingga 15 tahun
10 hingga 15ml setiap 6 hingga 8 jam
tidak melebihi 45ml dalam tempoh 24 jam
- Kanak-kanak 6 hingga 12 tahun
5 hingga 10ml setiap 6 hingga 8 jam
tidak melebihi 30ml dalam tempoh 24 jam
- Kanak-kanak 2 hingga 5 tahun
5ml setiap 6 hingga 8 jam
tidak melebihi 20ml dalam tempoh 24 jam

Setiap 5ml mengandungi:
Carbocisteine 100mg
Promethazine Hydrochloride 2.5mg
Methyl Hydroxybenzoate 0.1% w/v dan
Propyl Hydroxybenzoate 0.024% w/v sebagai bahan pengawet.
Simpan di bawah suhu 25° C.

服量：应分开服食，傍晚时分为佳。

- 成人：每6至8小时服15至20毫升
24小时内不可超过60毫升
- 12至15岁儿童：每6至8小时服10至15毫升
24小时内不可超过45毫升
- 6至12岁儿童：每6至8小时服5至10毫升
24小时内不可超过30毫升
- 2至5岁儿童：每6至8小时服5毫升
24小时内不可超过20毫升

每5毫升含：

Carbocisteine 100毫克
Promethazine Hydrochloride 2.5毫克

Methyl Hydroxybenzoate 0.1% w/v 和
Propyl Hydroxybenzoate 0.024% w/v 为防腐剂
置放在25° C以下。

注意：服食过量有危险。此药可使人昏昏欲睡，
服食后如有此情况，不得驾驶或动用机械。

CONTROLLED MEDICINE

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

储存于儿童的活动范围以外。

FOR ORAL USE ONLY

CODE MU0501

MAL 06071081AZ HK-58930



Cawan penyukat dos
percuma
免费度量器

B
MFG
EXP



Manufacturer and Product Registration Holder
Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5, Cheng Industrial Estate
75250 Melaka, Malaysia

