

Palavo

Suspension

Paracetamol (120mg/5ml)

What is in this leaflet:

1. What is Palavo Suspension used for
2. How Palavo Suspension works
3. Before you use Palavo Suspension
4. How to use Palavo Suspension
5. While you are using Palavo Suspension
6. Side effects
7. Storage and disposal of Palavo Suspension
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What is Palavo Suspension used for

Palavo Suspension is used for mild to moderate pain and for fever.

How Palavo Suspension works

Palavo Suspension contains paracetamol that has analgesic (relieve pain) and antipyretic (reduce fever) properties and weak anti-inflammatory activity (reduce inflammation).

Before you use Palavo Suspension

When you must not to use it

Do not use Palavo Suspension to your children if they:

- are known hypersensitivity to any ingredients in this medicine
- are taking other medicines containing paracetamol at the same time

Before you start to use it

Tell your doctor or pharmacist if:

- Your children are less than 1 month and/or 3 months old.

Taking other medicines

Tell your doctor if your children are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

How to use Palavo Suspension

How much to use

If you know the body weight of your child, it is always more accurate to use the weight to get the correct dose. Please refer below.

$$\text{Dose} = \frac{\text{BW (kg)} \times 15 \times 5\text{ml}}{120}$$

Body Weight	Dose	How Often
10 kg	6 ml	Do not take more

15 kg	9 ml	frequently than every 4 hours and do not exceed more than four doses in 24 hours.
20 kg	13 ml	
30 kg	19 ml	

For oral use only.

May be given every 4 to 6 hours, up to a maximum of 4 doses in a day.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

When to use it

Use as directed by your doctor or pharmacist.

How long to use it

- Maximum duration of continued use without advice from doctor or pharmacist: 3 days.
- Continue taking for as long as your doctor recommends.
- Ask your doctor or pharmacist before prolonged using.

If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to give the medicine to your children.

Give your children the missed dose as soon as you remember. If it is almost time for the next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Over dosage may cause damage to the liver and kidney. Prompt treatment with acetylcysteine or methionine is essential.

While you are using Palavo Suspension

Things you must do

- This product is for oral use only.

- Shake well before use.

Give your children the medicine exactly as your doctor has told you.

Tell all the doctors and pharmacists treating your children that they are taking Palavo Suspension.

Tell your doctor immediately if your children (under 3 months) persist to have fever for more than 24 hours (with maximum 4 doses in day). This is to ensure that fever that may be due to a serious infection is quickly diagnosed.

Things you must not do

- Do not exceed the stated dose.
- Do not take any new medicines without consulting your doctor.
- Do not give this product to anyone else, even if they have the same symptoms or condition as your children.

Things to be careful of

- Please always consult your doctor or pharmacist before giving your children this product especially when there is a known to hypersensitivity.

WARNING

This preparation contains PARACETAMOL. Do not take any other paracetamol containing medicines at the same.

There is potential for liver toxicity and severe liver damage. Do not take more than the recommended dose

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash.

These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Side effects

- Side effects are rare and usually mild, although haematological reactions including thrombocytopenia (reduce platelet in blood) and leucopenia (less white blood cell).

- Skin rashes and other hypersensitivity reactions occur occasionally.
- Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/ Toxic Epidermal Necrolysis have been reported.

Like all medicines, this product can cause side effects, although not everybody gets them. Visit your doctor or pharmacist immediately if your children experience any side effects after taking this medicine.

If your children experience any symptoms persistently for more than 7 days, please consult your doctor, dentist or pharmacist.

You may report any side effects or adverse drug reactions directly to The National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.moh.gov.my [Public → Reporting Side Effects to Medicinal (ConSERF) or Vaccine (AEFI)].

Storage and disposal of Palavo Suspension

Storage

Store below 30°C.

Keep out of reach of children.

The expiry date of this product is indicated on the packaging.

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

How it looks like

Each bottle contains orange colour suspension with orange flavour.

Available in pack size of 60ml.

Ingredients

Active ingredients

Paracetamol

Inactive ingredients

Syrup, Hydroxypropyl Methylcellulose, Xanthan Gum, Glycerin, Saccharin Sodium, Citric Acid Anhydrous, Sodium Citrate, Colourant, Orange Flavour, Methyl Hydroxybenzoate, Propyl Hydroxybenzoate, Polyethylene Glycol and Purified Water.

MAL Number

MAL

Manufacturer and Product Registration Holder

Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5, Cheng Industrial Estate,
75250 Melaka,
Malaysia.

Date of revision

25/11/2021

Serial Number

NPRA (R2) 21/113