

Palavo Tablet 650mg

Paracetamol (650mg)

What is in this leaflet:

1. What is Palavo Tablet used for
2. How Palavo Tablet works
3. Before you use Palavo Tablet
4. How to use Palavo Tablet
5. While you are using Palavo Tablet
6. Side effects
7. Storage and disposal of Palavo Tablet
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial number

What is Palavo Tablet used for

Palavo Tablet is used for mild to moderate pain and for fever.

How Palavo Tablet works

Palavo Tablet contains paracetamol that has analgesic (relieve pain) and antipyretic (reduce fever) properties and weak anti-inflammatory activity (reduce inflammation).

Before you use Palavo Tablet

When you must not to use it

Do not use Palavo Tablet if you:

- are known hypersensitivity to any ingredients in this medicine
- are taking other medicines containing paracetamol at the same time

Before you start to use it

Tell your doctor or pharmacist if:

- You are pregnant or plan to get pregnant.
- You are breast-feeding.

Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

How to use Palavo Tablet

How much to use

For oral use only.

Adults: 1 - 1½ tablets every 4-6 hours (not more than 6 tablets in one day unless prescribed by doctor).

To be taken before or after food with a glass of water.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the

instructions on the label, ask your doctor or pharmacist for help.

When to use it

Use as directed by your doctor or pharmacist.

How long to use it

- Continue taking for as long as your doctor recommends.
- Ask your doctor or pharmacist before prolonged using.

If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Over dosage may cause nausea, vomiting, sweating, abdominal pain, severe liver damage and sometimes acute renal tubular necrosis, coma and the worst is death. Prompt treatment with acetylcysteine or methionine is essential.

While you are using Palavo Suspension

Things you must do

This product is for oral use only.

Take your medicine exactly as your doctor has told you.

Tell all the doctors and pharmacists treating you that you are taking Palavo Tablet.

Tell your doctor immediately if you become pregnant while taking this medication.

Things you must not do

- Do not exceed the stated dose.
- Do not take any new medicines without consulting your doctor.
- Do not give this product to anyone else, even if they have the same symptoms or condition as you.

Things to be careful of

- Please always consult your doctor or pharmacist before using this product especially when there is a known to hypersensitivity.

WARNING

This preparation contains
PARACETAMOL.

Do not take any other paracetamol containing medicines at the same.

There is potential for liver toxicity and severe liver damage. Do not take more than the recommended dose.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Side effects

- Side effects are rare and usually mild, although haematological reactions including thrombocytopenia (reduce platelet in blood) and leucopenia (less white blood cell).
- Skin rashes and other hypersensitivity reactions occur occasionally.
- Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/ Toxic Epidermal Necrolysis have been reported.

Like all medicines, this product can cause side effects, although not everybody gets them. Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

If you experience any symptoms persistently for more than 7 days, please consult your doctor, dentist or pharmacist.

You may report any side effects or adverse drug reactions directly to The National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.moh.gov.my [Public → Reporting Side Effects to Medicinal (ConSERF) or Vaccine (AEFI)].

Storage and disposal of Palavo Tablet

Storage

Store below 30°C.

Keep out of reach of children.

The expiry date of this product is indicated on the packaging.

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

How it looks like

Palavo Tablet is white, oval, with “XS” and breakline on one side and plain on the reverse.

Each tablet contains Paracetamol 650mg. Available in pack size of 70 tablets per box (10 tablets per blister).

Ingredients

Active ingredients

Paracetamol

Inactive ingredients

Corn Starch, Pregelatinized Starch, PVP K-30, Sodium Starch Glycolate and Stearic Acid.

MAL Number

MAL

Manufacturer and Product Registration Holder

Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5, Cheng Industrial Estate,
75250 Melaka,
Malaysia.

Date of revision

12/11/2021

Serial Number

NPRA (R2) 21/135