

MIOLIV FILM-COATED TABLET

Domperidone (10mg)

What is in this leaflet

1. What is Mioliv used for
2. How Mioliv works
3. Before you use Mioliv
4. How to use Mioliv
5. While you are using it
6. Side effects
7. Storage and Disposal of Mioliv
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision
11. Serial Number

What is Mioliv used for

Mioliv is indicated for the relief of the symptoms of nausea and vomiting.

How Mioliv works

MIOLIV is an antiemetic. It works by blocking the action of a chemical messenger in the brain which causes the feeling of nausea and vomiting, as well as increasing the movement or contractions of the stomach and intestines, allowing food to move more easily through the stomach.

Before you take Mioliv

When you must not use it

Do not take Mioliv if:

- you are taking any of the following medicines: clarithromycin (antibiotic).
- you have an allergy to Mioliv or any of the ingredients at the end of this leaflet.
- you have a tumour of the pituitary gland called prolactinoma, bleeding, obstruction or perforation in the stomach and intestines.
- for prevention of post-operative vomiting or for long term usage.

Before start to take it

Tell your doctor or pharmacist:

- If you are pregnant or plan to become pregnant.
- If you have ever been told that you have problems with your liver.

Taking other medicines

Do not take Mioliv together with:

- Anticholinergic medicines used to treat stomach cramp

- Antacid medicines for heartburn
- Antisecretory medicines
- Clarithromycin (antibiotic)

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without prescription.

How to use Mioliv

How much to use

You should always take the lowest amount of Mioliv that works for you and you should not take it for longer than is necessary. Although the amount of Mioliv you should usually take is described below, your doctor may adjust your dose to your personal needs.

Mioliv is most effective if taken 15-30 minutes before meals.

Adults and adolescents (12 years of age and over) weighing 35kg or more; children weighing 35kg or more:

Take 1 tablet 3 to 4 times a day. Do not take more than 4 tablets per day (40mg/day).

Infants and children less than 12 years of age and weighing less than 35kg:

The effectiveness of Mioliv has not been established in infants and children under 12 years of age with a body weight of <35kg.

When to use it

Mioliv is most effective if taken 15 to 30 minutes before meals.

How long to use it

Continue taking your medicine for as long as your doctor tells you.

If you forget to use it

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Otherwise, take it as soon as you remember, and then go back to taking your capsules as you would normally.

Do not take a double dose to make up for the dose that you missed. This may

increase the chance of you getting an unwanted side effect.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Some of the symptoms of an overdose may include:

- nervous
- uncontrolled movements

While you are using it

Things you must do

- Tell all the doctors, dentists and pharmacists who are treating you that you are taking Mioliv.
- If you are about to start taking a new medicine, tell your doctor and pharmacist that you are taking Mioliv.

Things you must not do

- Do not give Mioliv to anyone else, even if they have the same symptoms or condition as you.
- Do not use Mioliv to treat any other complaints unless your doctor tells you to.
- Do not use Mioliv if you are breast feeding or wish to breastfeed unless the expected benefits outweigh any potential risk.

Things to be careful of

Be careful driving or operating machinery until you know how Mioliv affects you.

Side effects

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

Advise to consult doctor/pharmacist for further information.

MIOLIV FILM-COATED TABLET

Domperidone (10mg)

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Mioliv

Storage

Store below 30 °C.

Keep out of the reach of the children.

Do not use this medicine after expiry date.

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

What it looks like

Pink, normal convex, round shape film-coated tablet, with “XS” marking on one side and plain on the reverse.

Ingredients

- Active Ingredient

Domperidone

- Inactive Ingredients

Lactose, Microcrystalline Cellulose, Starch, Sodium Lauryl Sulphate, Magnesium Stearate, Colloidal Silicon Dioxide, HPMC Compound Mix, Colour Pigment Red.

MAL Number

MALxxxxxxxAZ

Manufacturer and Product

Registration Holder

Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5 Cheng Industrial Estate,
75250 Melaka, Malaysia.

Date of Revision

11/07/2024

Serial Number

NPRA (R2) 23/185