

90ml

Each 5ml contains:

Diphenhydramine HCl 14mg
Ammonium Chloride 135mg
Sodium Benzoate 0.15% w/v as preservative.

Store in a cool place below 30°C.

Refer to accompanying package
insert for further information.

MAL 16105026AZ BRU16022010P



Xepa^{SP}
Manufacturer and Product Registration Holder
Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5, Cheng Industrial Estate
75250 Melaka, Malaysia

CONTROLLED MEDICINE

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

儲存于兒童的活動範圍以外。

FOR ORAL USE ONLY

CODE BE0109

Bena[®] Expectorant

Expectorant, Antihistamine

Clears chesty cough

Mixed fruity flavour

Dosage:

- Adults & children over 12 years:
5 to 10ml every 4 to 6 hours
not to exceed 60ml in 24 hours
- Children 6 to 12 years:
2.5 to 5ml every 4 to 6 hours
not to exceed 30ml in 24 hours
- Children 2 to 5 years:
2.5ml every 4 to 6 hours
not to exceed 20ml in 24 hours

*Take medication with
a full glass of water.*

Caution:

This medicine may cause
drowsiness. If affected, do not
drive or operate machinery.
Avoid alcoholic drinks.

Warnings:

When used for treatment cough and cold:
Not to be used in children less than 2
years of age. To be used with caution
and doctor's/pharmacist's advice in
children 2 to 6 years of age.

BE0109

BATCH
MFG EXP
005

120ml

Each 5ml contains:

Diphenhydramine HCl14mg

Ammonium Chloride135mg

Sodium Benzoate 0.15% w/v as preservative.

Store in a cool place below 30°C.

Refer to accompanying package insert for further information.

CODE BE0107

MAL 16105026AZ BRU16022010P

CONTROLLED MEDICINE

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

儲存于兒童的活動範圍以外。

FOR ORAL USE ONLY



Manufacturer and Product Registration Holder

Xepa-Soul Pattinson (Malaysia) Sdn Bhd

1-5 Cheng Industrial Estate, 75250 Melaka, Malaysia

Bena[®] Expectorant

Expectorant, Antihistamine

Clears chesty cough

Mixed fruity flavour

Dosage:

- Adults & children over 12 years:

5 to 10ml every 4 to 6 hours

not to exceed 60ml in 24 hours

- Children 6 to 12 years:

2.5 to 5ml every 4 to 6 hours

not to exceed 30ml in 24 hours

- Children 2 to 5 years:

2.5ml every 4 to 6 hours

not to exceed 20ml in 24 hours

*Take medication with
a full glass of water.*

Caution:

This medicine may cause drowsiness. If affected, do not drive or operate machinery. Avoid alcoholic drinks.

Warnings :

When used for treatment of cough and cold: Not to be used in children less than 2 years of age. To be used with caution and doctor's / pharmacist's advice in children 2 to 6 years of age.

BE0107

BATCH
MFG EXP

006