

# FEPRIL TABLET 500MG®

Consumer Medication Information Leaflet (RiMUP)

Paracetamol (500mg)

## What is in this leaflet

1. What Fepril Tablet is used for
2. How Fepril Tablet works
3. Before you use Fepril Tablet
4. How to use Fepril Tablet
5. While you are using it
6. Side effects
7. Storage and Disposal of Fepril Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial number

## What Fepril Tablet is used for

Fepril Tablet is used for symptomatic relief of mild to moderate pain and to reduce fever.

## How Fepril Tablet works

Fepril Tablet has analgesic (pain relief) and antipyretic (anti-fever) properties. It works by blocking the pain impulse generation and increasing the pain threshold. It also increases the blood flow across the skin, heat loss and sweating to reduce fever.

## Before you use Fepril Tablet

### - When you must not use it

Do not take this medicine if:

- you are allergic to paracetamol or any other ingredients of this medicine (listed in Product Description)
- you have liver problems

### - Before you start to use it

Tell your doctor if you have any of the following:

- an enzyme problem known as G6PD deficiency
- problems with nutrition (chronic malnutrition or dehydration)
- body weight below 50kg

### Pregnancy and lactation

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine.

### - Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a

pharmacy, supermarket or health food shop.

This preparation contains PARACETAMOL. Do not take any other paracetamol containing medicines at the same time. Inform your doctor or pharmacist if you are using this medicine in combination with the following:

- Anticoagulants such as warfarin (blood thinning medicine)
- Metoclopramide or domperidone (medicines for stomach problems)
- Cholestyramine (medicine to reduce cholesterol)

## How to use Fepril Tablet

### - How much to use

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

**Adults and children aged 12 years and over:** 1 to 2 tablets, 3 or 4 times a day up to a maximum of 8 tablets (4000mg) daily.

**Children aged 7 – 12 years:** ½ to 1 tablet, 3 or 4 times daily.

### - When to use it

Use as directed by your doctor or pharmacist.

### - How long to use it

Use Fepril Tablet for the duration that the doctor recommends.

### - If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### - If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have used too much of this medicine. Do this even if there are no

signs of discomfort or poisoning. You may need urgent medical attention.

Symptoms of paracetamol overdose in the first 24 hours include pallor, nausea, vomiting, loss of appetite and abdominal pain. Signs of liver damage (pain, tenderness, abdominal swelling) may be seen 12 to 48 hours after paracetamol overdose. Paracetamol overdose may also affect the kidneys or heart.

## While you are using it

### - Things you must do

Use your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are using Fepril Tablet.

Tell your doctor immediately if you become pregnant while using this medication.

### - Things you must not do

Children should not take paracetamol for more than 3 days with consulting a doctor.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Fepril Tablet to anyone else, even if they have the same symptoms or condition as you.

### - Things to be careful of

#### Driving and using machines

Some medicines may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

## Side effects

Like all medicines, Fepril Tablet can cause side effects, although not everybody gets them.

**ALLERGY ALERT:** Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters and rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

The side effects of Fepril tablet are usually mild, although reactions related to the blood and bone marrow cells have been reported. Other possible side effects include low blood pressure, increased heart rate, nausea, vomiting, constipation, difficulty sleeping, liver or kidney problems.

Tell your doctor or pharmacist if you notice anything that is making you feel unwell.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my) [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

### **Storage and Disposal of Fepril Tablet**

- Storage

Keep out of the reach and sight of children.

Keep in a dry place, store below 30°C and protect from light.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

### **Product Description**

- What it looks like

Round and biconvex tablets with one side scored. White to off white in colour.

- Ingredients

- Active ingredient

Each tablet contains paracetamol 500 mg

- Inactive ingredients

Microcrystalline cellulose 101, Pregelatinized starch, Croscarmellose Sodium, Povidone, Potassium Sorbate, Talc Powder, Stearic Acid.

- MAL number:

MAL19912157XZ

### **Manufacturer & Product Registration Holder**

Idaman Pharma Manufacturing Sdn Bhd, (661901-P)  
Lot 24 & 25, Jalan Perusahaan Lapan, Bakar Arang Industrial Estate, 08000 Sungai Petani, Kedah Darul Aman, Malaysia.

### **Date of revision**

12/11/2024

### **Serial Number**

NPRA(R4/1)111124/00122