

CORBETA TABLET 40MG

Propranolol Hydrochloride (40mg)

What is in this leaflet

1. What Corbета Tablet is used for
2. How Corbета Tablet works
3. Before you use Corbета Tablet
4. How to use Corbета Tablet
5. While you are using Corbета Tablet
6. Side Effects
7. Storage and Disposal of Corbета Tablet
8. Product Description
9. Product Registration Holder and Manufacturer
10. Date of Revision
11. Serial Number

What Corbета Tablet is used for

Corbета Tablet is used to:

- Treat high blood pressure
- Treat severe chest pain
- Control irregular or fast heart beats and other symptoms caused by an overactive thyroid gland
- To reduce migraine attacks
- To calm people who are anxious or worried
- Prevent stomach bleeding in patients with high blood pressure in their liver or swollen blood vessels in their gullet

How Corbета Tablet works

Corbета Tablet works by slowing down the heart and decreasing the amount of blood it pumps out. This helps to decrease blood pressure, the heart pumps more efficiently, and reduces the workload on the heart.

Before you use Corbета Tablet

- When you must not use it

Do not use Corbета Tablet in the following conditions:

- Allergic to any ingredient of the tablet
- Severe heart block
- Sick sinus syndrome
- Cardiogenic shock
- Bronchial asthma
- Child with diabetes and heart failure

If you are not sure whether you can use this medicine, please consult your doctor or pharmacist first.

- Before you start to use it

Tell your doctor or pharmacist if you have:

- Other heart problem
- A history of allergic reactions
- Kidney or liver disease

- Wolff-Parkinson-White syndrome
- Smoke or drink alcohol

Pregnancy and Lactation

Safety in pregnancy has not been established. It is not recommended in nursing mothers unless it outweighs any potential risk.

You must tell your doctor if you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby.

Your doctor or pharmacist will advise you if you can use Corbета Tablet during pregnancy or while you are breast-feeding.

- Taking other medicines

Corbета Tablet should not be used with verapamil, clonidine and mibefradil.

Tell your doctor or pharmacist if you are using any other medicines, including those that you buy without a prescription from pharmacy, medical hall or any other source.

How to use Corbета Tablet

Always use the medicine exactly as your doctor or pharmacist has told you. Read the directions on the label carefully. If you have any concerns about how to use this medicine, talk to your doctor or pharmacist.

- How much to use

Your doctor or pharmacist will advise you how much to use each day and for how long you will need to use it.

Always take the dose exactly as your doctor has told you.

Adults' dosage:

Hypertension: The usual effective dose range is 40-320mg per day.

Angina pectoris: The average optimum dosage is 160mg per day.

Migraine: Initial dose is 80mg/day, may increase up to 160-240mg/day.

Pediatric dosage:

Doses are decided using body weight.

- When to use it

Follow the instruction on the label carefully

and as instructed by your doctor or pharmacist.

Take Corbета Tablet before meals. If you are taking an antacid with aluminum in it, take it at least 2 hours after taking Corbета Tablet.

- How long to use it

Continue taking Corbета Tablet for as long as your doctor recommends.

- If you forget to use it

If you miss a dose of Corbета Tablet, take it as soon as you remember. If it is almost time for your next dose, wait until then to take the dose and skip the missed dose. Do not take extra dose to make up for the missed dose.

- If you take too much (overdose)

Immediately call your doctor or go to the emergency department at your nearest hospital if you or anyone else has taken a large amount of this medication.

Examples of overdose symptoms are dizziness, shortness of breath, swelling of the hands/ankles/feet, and very slow heart rate.

While you are using Corbета Tablet

- Things you must do

Use the medication exactly as your doctor prescribed.

- Things you must not do

While you are using this medicine, do not do the following action:

- Do not suddenly stop taking the medicine
- Do not take more than the recommended dose
- You are advised to avoid alcohol to consume this product
- Do not drive or operate machinery if there is visual disturbances, hallucinations, fatigue, confusion, dizziness or tiredness after taking the medicine
- If you have been told that you have intolerance to some sugars, contact your doctor before taking this medicine.

- Things to be careful of

- Driving and using machines

Propranolol may cause visual disturbances, hallucinations, fatigue, mental confusion,

CORBETA TABLET 40MG

Propranolol Hydrochloride (40mg)

dizziness or tiredness. Make sure you are not affected before you drive or operate machinery.

longer required. These measures will help to protect the environment.

Side Effects

Like all medicines, Corbeta Tablet also can cause side effects, although not everybody gets them. Inform your doctor or pharmacist immediately if you experience any side effects after using this medicine.

Tell your doctors if you have experienced the following side effects:

- Slow heart rate
- Low blood pressure causing dizziness, lightheadedness, fainting or blurred vision
- Allergic reaction such as itching, difficulty in breathing and swelling of the face, lips, throat or tongue.

Other side effects which may occur more commonly are congestive cardiac failure, Raynaud's Disease, hallucinations, nightmares, weakness, fatigue, visual disturbance and bronchospasm.

Other side effects not listed above may occur in some patients. Ask your doctor or pharmacist to answer any questions you may have.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website nptra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Corbeta Tablet

- Storage

Keep all medicines out of the sight and reach of children.

Do not use this medicine after the expiry date stamped on the pack.

Store below 30°C in a cool dry place. Protect from light and moisture.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no

Product Description

- What it looks like

Clean, uniform, round, flat, bevel edged tablets, scored on one side and pink in color.

- Ingredients

- Active Ingredient

Propranolol Hydrochloride

- Inactive Ingredient

- Ponceau 4R
- Lactose
- Maize Starch
- Magnesium Stearate

- MAL Number:

MAL19860175AZ

Product Registration Holder & Manufacturer

Idaman Pharma Manufacturing Sdn Bhd
(200401023395)

Lot 24 & 25, Jalan Perusahaan Lapan, Bakar Arang Industrial Estate,
08000 Sungai Petani
Kedah Darul Aman, Malaysia

Date of Revision

26/10/2023

Serial Number

NPRA(R2/1)191226/313