

IDAMAN PHARMA CHLORPROMAZINE TABLET 100MG

Chlorpromazine Hydrochloride (100mg)

What is in this leaflet

1. What Idaman Pharma Chlorpromazine Tablet 100mg is used for
2. How Idaman Pharma Chlorpromazine Tablet 100mg works
3. Before you use Idaman Pharma Chlorpromazine Tablet 100mg
4. How to use Idaman Pharma Chlorpromazine Tablet 100mg
5. While you are using Idaman Pharma Chlorpromazine Tablet 100mg
6. Side Effects
7. Storage and Disposal of Idaman Pharma Chlorpromazine Tablet 100mg
8. Product Description
9. Product Registration Holder and Manufacturer
10. Date of Revision

What Idaman Pharma Chlorpromazine Tablet 100mg is used for

Idaman Pharma Chlorpromazine Tablet 100mg contains Chlorpromazine Hydrochloride. It is used in the treatment of behavioral disturbances. It is also used for the short-term treatment of anxiety, agitation and violent or impulsive behavior. It is used to treat persistent hiccup and nausea and vomiting associated with terminal illness.

How Idaman Pharma Chlorpromazine Tablet 100mg works

Idaman Pharma Chlorpromazine Tablet 100mg is thought to work by blocking the action of a chemical called dopamine which acts on the nervous system in the brain.

Before you use Idaman Pharma Chlorpromazine Tablet 100mg

-When you must not use it

You must not take Idaman Pharma Chlorpromazine Tablet 100mg if you have:

- Allergic to chlorpromazine.
- Pregnant or breast-feeding
- History of blood problem.
- Severe heart disease.
- Dulled senses such as feeling sleeping or uncoordinated, having blurred vision, slurred speech or being less aware of your surroundings.

If you are not sure whether you should start using this medicine, talk to your doctor or pharmacist first.

Pregnancy and lactation

If you are pregnant or breast-feeding or think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking Idaman Pharma Chlorpromazine Tablet 100mg.

How to use Idaman Pharma Chlorpromazine Tablet 100mg

- How much to use

Always take Idaman Pharma Chlorpromazine Tablet 100mg exactly as your doctor has told you. If you do not understand the directions on the label, ask your doctor or pharmacist to explain the instructions to you.

Doses:

In most patient oral treatment may be used from the start, commencing with a dosage of 25-50 mg thrice daily and increasing as necessary. The usual daily maintenance dose is range from 25-100 mg thrice daily.

Psychotic patient may require daily doses of up to 1 g or more.

In the treatment of mild alcohol withdrawal, agitated patients may be given 25 to 50 mg twice daily by mouth. Patients with less severe withdrawal symptoms may be given 30 to 75 mg daily in divided doses by mouth.

Chlorpromazine should not be given to stupors patients.

Pediatric use of the tablets is not recommended.

- When to use it

Swallow the tablets with water. If you are taking antacids, you should take at least 2 hours after taking chlorpromazine.

Follow the instructions on the label carefully and as instructed by your doctor or pharmacist.

- How long to use it

Do not take more than 8 tablets in 24 hour period or 8 tablets in a day for more than 2 weeks except under the advice and supervision of your doctor or pharmacist.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

If you take too much (overdose)

If you have taken more than the prescribed dose, contact your doctor immediately. Signs of an overdose include:

- Quick and shallow breaths
- Low body temperature
- Low blood pressure
- Restlessness
- Fits
- Unusual heart beats
- Coma

Contact your doctor immediately or go to Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no sign of discomfort or poisoning. You may need urgent medical attention.

While you are using Idaman Pharma Chlorpromazine Tablet 100mg

- Things you must do

Take your medicine exactly as your doctor or pharmacist has told you.

Tell your doctor and pharmacist treating you that you are taking Idaman Pharma Chlorpromazine Tablet 100mg.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor. Do not take any new medicines without consulting your doctor. Do not give Idaman Pharma Chlorpromazine Tablet 100mg to anyone else even they have the same symptoms or condition as you.

- Things to be careful of

The high intake of aluminum containing antacids in elderly patients is not generally recommended because it may leads to Alzheimer's disease.

Risk benefits should be considered when the following medical problems exist:

- Alzheimer's disease
- Appendix or symptoms of appendicitis

IDAMAN PHARMA CHLORPROMAZINE TABLET 100MG

Chlorpromazine Hydrochloride (100mg)

- Undiagnosed gastrointestinal or rectal bleeding, ulcerative colitis (inflammation in colon).
- Constipation
- Fecal impaction (chronic constipation)
- Chronic diarrhea
- Intestinal obstruction (blockage of the bowel that prevents the contents of the intestine from passing through).
- Kidney problem.

Side Effects

Like all medicines, Idaman Pharma Chlorpromazine Tablet 100mg can cause side effects, although not everybody gets them.

Common side effects are:

- Drowsiness
- Dryness of the mouth
- Nasal congestion
- Postural hypotension
- Lowering of body temperature
- Tachycardia
- Arrhythmias
- Agitation
- Insomnia
- Depression
- Miosis and mydriasis
- Convulsions
- Photosensitivity
- Skin rashes
- Inhibition of ejaculation

This is not a complete list of side effects. You shall contact your doctor or pharmacist if you experience any side effects.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.moh.gov.my [Consumer → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Idaman Pharma Chlorpromazine Tablet 100mg

- Storage

Keep all medicines out of the sight and reach of children.

Store below 30°C. Protect from light.

Do not store in the bathroom or near a sink, or on a window sill.

Do not leave it in the car on hot days. Heat and dampness can destroy some medicines.

Do not take this medicine after the expiry date stamped on the pack. The expiry date refers to the last day of that month.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Clean, dry, round, uniform, biconvex, even edged film coated tablets and scored on one side, white to off white in colour.

- Ingredients

Active ingredient:

Chlorpromazine Hydrochloride

Inactive ingredients:

- Lactose Monohydrate
- Corn Starch (Maize Starch)
- Dioctyl sodium Sulphosuccinate
- Plasdone K30
- Ethyl Alcohol 96%
- Talc Powder
- Magnesium Stearate
- Sodium Starch Glycolate
- Opadry White 85F18422
- Purified Water

- MAL Number

MAL19880027AZ

Product Registration Holder & Manufacturer

InspiraPharma Sdn Bhd
(200401023395)
Lot 24 & 25, Jalan 8,
Kawasan Perusahaan Bakar Arang,
08000 Sungai Petani,
Kedah Darul Aman, Malaysia.

Date of Revision

08/07/2026

Serial Number:

NPRA(R2/1)200902/512