

# PHARMANIAGA CO-AMOXICLAV TABLET

Amoxycillin trihydrate/Clavulanic acid (500mg/125mg)

## What is in this leaflet

1. What Pharmaniaga Co-Amoxiclav Tablet is used for
2. How Pharmaniaga Co-Amoxiclav Tablet works
3. Before you use Pharmaniaga Co-Amoxiclav Tablet
4. How to use Pharmaniaga Co-Amoxiclav Tablet
5. While you are using it
6. Side effects
7. Storage and Disposal of Pharmaniaga Co-Amoxiclav Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What Pharmaniaga Co-Amoxiclav Tablet is used for

Pharmaniaga Co-Amoxiclav Tablet is used to treat the following bacterial infections:

- respiratory tract infections
- skin and soft tissue infections
- urinary tract infections
- bone and joint infections

## How Pharmaniaga Co-Amoxiclav Tablet works

Pharmaniaga Co-Amoxiclav Tablet contains two active ingredients which are amoxicillin trihydrate and clavulanic acid. Both are antibiotics that work by killing bacteria causing the infections.

## Before you use Pharmaniaga Co-Amoxiclav Tablet

### - When you must not take it

Do not take this medicine if:

- you are allergic or ever had severe allergic reactions to amoxicillin, clavulanic acid, other penicillin based antibiotics, or similar antibiotics called cephalosporins or any other ingredients of this medicine (listed in Product Description).
  - you have ever had liver problems or jaundice (yellowing of the skin and/or whites of the eyes) when taking antibiotic.
- Before you start to take it  
Tell your doctor or pharmacist if:
- you have liver or kidney problems

- has history of diarrhea during or after antibiotic therapy
- you are not urinating regularly

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking Pharmaniaga Co-Amoxiclav Tablet.

### *Pregnancy and lactation*

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine.

### - Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

You should tell your doctor if you are taking the following medicine as it may interact with Pharmaniaga Co-Amoxiclav Tablet:

- probenecid and allopurinol (medicine for gout)
- oral contraceptives
- blood thinning medications (such as warfarin)
- mycophenolate mofetil (medicine used to prevent the rejection after organ transplant)
- methotrexate (medicine used to treat cancer or rheumatic diseases)

## How to use Pharmaniaga Co-Amoxiclav Tablet

### - How much to take

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

For oral administration only.

### *Adult and children over 12 years old*

Mild to moderate infections: Take one tablet two times a day.

### *Kidney and liver problems*

If you have kidney problems, the frequency might be changed. If you have liver problems, you may have

more frequent blood tests to see how your liver is working.

### - When to take it

Take 2 times per day or as directed by your doctor or pharmacist.

Swallow the tablet whole with a glass of water at the start of a meal. Do not chew or cut the tablet.

### - How long to take it

Take Pharmaniaga Co-Amoxiclav Tablet for the duration that the doctor recommends.

The treatment duration is usually up to 14 days. If your condition worsens or no improvement seen, please see the doctor again.

Keep taking Pharmaniaga Co-Amoxiclav Tablet until the treatment is finished, even if you feel better.

### - If you forget to take it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### - If you take too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have use too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may cause upset stomach, imbalance of electrolytes, formation of crystals in the urine, kidney problems or convulsions.

## While you are using it

### - Things you must do

Tell all the doctors, dentists and pharmacists treating you that you are taking Pharmaniaga Co-Amoxiclav Tablet.

# PHARMANIAGA CO-AMOXICLAV TABLET

Amoxycillin trihydrate/Clavulanic acid (500mg/125mg)

Tell your doctor immediately if you become pregnant while taking this medication.

Take Pharmaniaga Co-Amoxiclav Tablet only with a doctor's prescription.

- **Things you must not do**

Do not stop using the medicine unless advised by your doctor.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Pharmaniaga Co-Amoxiclav Tablet to anyone else, even if they have the same symptoms or condition as you.

- **Things to be careful of**

**Driving and using machines**

Be careful driving or operating machinery until you know how Pharmaniaga Co-Amoxiclav Tablet affects you.

If any skin reactions occur while taking this medication, stop taking it and see the doctor immediately.

If you develop diarrhoea while taking this medication, or even after finishing the course of the antibiotic, stop taking it and see the doctor immediately.

**Blood and urine tests**

If you are having Coomb's test (red blood cell status tests), let the doctor or nurse know that you are taking Pharmaniaga Co-Amoxiclav Tablet, as this medicine can affect the test results.

**Side effects**

Like all medicines, Pharmaniaga Co-Amoxiclav Tablet can cause side effects, although not everybody gets them.

Stop taking Pharmaniaga Co-Amoxiclav Tablet and contact your doctor immediately if you experience any of the following:

- serious allergic reaction such as Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS). DRESS appears initially as flu-like

symptoms with a rash on the face and then with an extended rash, high temperature and enlarged lymph nodes.

Pharmaniaga Co-Amoxiclav Tablet may cause severe allergy and serious skin reactions. Stop using Pharmaniaga Co-Amoxiclav Tablet and seek medical assistance immediately if you experience any of the following symptoms:

- severe allergy: breathing difficulties, lightheadedness, skin swellings or rash
- severe skin reaction: skin reddening, blisters, rash, fever, sore throat or eye irritation

Very common and common side effects:

- diarrhea
- nausea
- vomiting
- fungal infection

Uncommon side effects:

- dizziness
- headache
- indigestion
- increase liver enzymes
- skin rash
- itchiness
- hives

Rare and very rare side effects:

- low white blood cells, red blood cells or platelets
- swelling of the skin (angioedema)
- inflammation of blood vessels which may be visible as red or purple raised spots on the skin (hypersensitivity vasculitis)
- unusually active
- convulsions
- black hairy tongue
- inflammation of the liver (hepatitis) & jaundice (yellowing of skin and whites of the eyes)
- inflammation of tubes in the kidney
- crystals in urine

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring visiting the website [npra.gov.my](http://npra.gov.my) [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

**Storage and Disposal of Pharmaniaga Co-Amoxiclav Tablet**

- **Storage**

Keep out of the reach and sight of children.

Store below 30°C and keep the remaining tablets in original packaging after open.

Use within 30 days after opening the pouch.

- **Disposal**

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

**Product Description**

- **What it looks like**

White to off white oblong shape film coated tablet with plain one side and word "RMB" on the other side.

- **Ingredients**

- Active ingredient(s)

Each film coated tablet contains:

- amoxicillin trihydrate equivalent to amoxicillin 500mg
- potassium clavulanate equivalent to clavulanic acid 125mg

- Inactive ingredients

Microcrystalline cellulose, magnesium stearate, colloidal silicon dioxide, pladone, croscarmellose sodium and opadry white.

- **MAL number:**

MAL20013767AZ

# PHARMANIAGA CO-AMOXICLAV TABLET

Amoxicillin trihydrate/Clavulanic acid (500mg/125mg)

---

**Manufacturer and Product  
Registration Holder**

Idaman Pharma Manufacturing Sdn Bhd  
Lot 120, Taman Farmaseutikal, 32610  
Bandar Sri Iskandar, Perak Darul  
Ridzuan, Malaysia.

**Date of revision**

8/8/2024

**Serial Number**

NPRA(R4/1)070824/00105