

IDAMOL TABLET 500MG

Paracetamol (500mg)

What is in this leaflet

1. What IDAMOL is used for
2. How IDAMOL works
3. Before you use IDAMOL
4. How to use IDAMOL
5. While you are using it
6. Side effects
7. Storage and disposal of IDAMOL
8. Product description
9. Product Registration Holder and Manufacturer
10. Date of revision

What IDAMOL is used for

IDAMOL is used for relief of mild to moderate pain including headache, toothache, sore throat and period pain. They are also used to reduce fever

How IDAMOL works

Paracetamol works as a painkiller by affecting chemicals in the body called prostaglandins. Prostaglandins are substances released in response to illness or injury. Paracetamol blocks the production of prostaglandins, making the body less aware of the pain or injury.

Paracetamol reduces temperature by acting on the area of the brain that is responsible for controlling temperature.

Before you use IDAMOL

- When you must not use it

Should not be used by person with allergic to paracetamol or any other ingredients listed at the end of this leaflet.

Should not be used by person with severe and active liver impairment.

Pregnancy and lactation

Consult with your doctor before taking IDAMOL if you are pregnant or think you may be pregnant. It can be considered as choice for analgesic in pregnancy and it considered to be safe even though it crosses placenta.

Consult with your pharmacist if you are breast feed your child since it excreted in breast milk.

Ask your doctor or pharmacist for advice before taking any medicine.

- Before you start to use it

Care should be taken if you having alcohol dependence, severe liver or urinary impairment when taking IDAMOL.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Some medicine might have interaction when taking together with IDAMOL. This include:

- Warfarin (medicine for blood thinning)
- Other anticoagulants medicine including Coumarin.

How to use IDAMOL

- How much to use

Following all direction given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

Doses:

Adults: 1 to 2 tablets taken every 4 to 6 hours as required up to a maximum of 8 tablets daily.

Children (6-11 years) : ½ to 1 tablet every 4 to 6 hours as required.

Maximum daily dose: 60 mg/kg presented in divided doses of 15 mg/kg throughout 24-hour period.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking IDAMOL for as long as your doctor recommends.

- If you forget to use it

Consult you doctor or pharmacist on what you should do if you forget to use it. Take the missed dose as soon as you remember. If it almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may cause serious liver damage.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not take new medicines without consulting your doctor.

Do not give IDAMOL to anyone else, even if they have the same symptoms or condition as you.

This preparation contains PARACETAMOL. Do not take any other paracetamol containing medicines at the same time.

- Things to be careful of

If you are pregnant or are breast-feeding, you should not take IDAMOL without discussing it with your doctor or pharmacist first.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blister or rash.

IDAMOL TABLET 500MG

Paracetamol (500mg)

These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Side Effects

Adverse effects of paracetamol are rare and usually mild, although hematological reactions have been reported.

Cutaneous hypersensitivity reactions including skin rashes, angioedema, Steven Johnson Syndrome/Toxic Epidermal Necrolysis have been reported.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npa.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of IDAMOL

- Storage

Keep in a dry place, below 30°C
Protect from light.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Clean, dry, round, biconvex tablets and scored on one side and white in colour

- Ingredients

Active ingredients :

Each tablet contains : Paracetamol
500mg

Inactive Ingredients

- Corn starch
- Pregelatinized starch
- Povidone
- Sodium carboxymethyl starch
- Stearic acid

MAL Number

MAL20046150XZ

Product Registration Holder & Manufacturer

InspiraPharma Sdn Bhd (200401023395)
Lot 24 & 25, Jalan 8,
Kawasan Perusahaan Bakar Arang,
08000 Sungai Petani,
Kedah Darul Aman, Malaysia.

Date of Revision

06/07/2026

Serial Number

NPRA (R2) 19/211