

Each vial contains 3.5 mg of Bortezomib

Reconstitution Information:

For IV Use: Add 3.5 ml of 0.9% Sodium Chloride to make 1 mg/ml final concentration.

For SC Use: Add 1.4 ml of 0.9% Sodium Chloride to make 2.5 mg/ml final concentration.

The total storage time for the reconstituted medicinal product should be not more than 8 hours at 25°C prior to administration.

For single use only.

Store below 20°C.

Keep the container in the outer carton box in order to protect from light.

Keep out of reach of children.

(Jauhkan daripada capaian kanak-kanak.)

For more information, please refer package insert.

pharmaniaga

Retrox

3.5 mg

Powder for Solution for Injection

(Bortezomib)

1 Vial

FOR IV or SC USE ONLY

CAUTION - CYTOTOXIC AGENT

Batch No.:

Mfg. Date:

Exp. Date:

MAL21066026AZ

Product Registration Holder:

InspiraPharma Sdn Bhd

Lot 24 & 25, Jalan 8, Kawasan Perusahaan Bakar Arang, 08000 Sungai Petani, Kedah Darul Aman, Malaysia

Manufacturer:

Shilpa Medicare Limited,

Noida/Mumbai - Distinct, 500091 Telangana, India


Code: TS/DUGS/1/2013

4500890

CONTROLLED MEDICINE (UBAT TERBAWAL)

(Unvarnished Area 18x10mm)

Dots not to be printed



CMYK

Superseeded Code: 4500890

Size: 70x22 mm

Market: Pharmaniaga (Malaysia)

R00 (13.02.2026)