

SAFFID CAPSULE 300MG

Gemfibrozil 300mg

What is in this leaflet

1. What Saffid Capsule 300mg is used for
2. How Saffid Capsule 300mg works
3. Before you use Saffid Capsule 300mg
4. How to use Saffid Capsule 300mg
5. While you are using Saffid Capsule 300mg
6. Side effects
7. Storage and Disposal of Saffid Capsule 300mg
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of Revision
11. Serial Number

What Saffid Capsule 300mg is used for

Saffid Capsule 300mg is used to treat hyperlipidaemia (having too much fat in the blood) and to prevent heart disease in people with hyperlipidaemia.

Saffid Capsule 300mg helps reduce cholesterol and triglycerides (fatty acids) in the blood. High levels of these types of fat in the blood are associated with an increased risk of atherosclerosis (hardening of blood vessels/arteries).

Saffid Capsule 300mg is also used to lower the risk of stroke, heart attack, or other heart complications in people with high cholesterol and triglycerides who have not responded to dietary and other measures.

How Saffid Capsule 300mg works

Gemfibrozil belongs to the class of medications known as *fibrates*. It works by lowering the level of certain types of fat and cholesterol, especially the type known as *triglycerides*.

Before you use Saffid Capsule 300mg

When you must not to use it

Do not take Saffid Capsule 300mg if you:

- are allergic to any medicine containing gemfibrozil.
- are allergic to fenofibrate or any other of the class of “fibrates” (e.g., clofibrate, bezafibrate)
- are breast-feeding
- are pregnant
- have severe liver disease, kidney disease, or gallbladder disease

Before you start to use it

Tell your pharmacist if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor or pharmacist if you are pregnant or plan to become pregnant or are breast-feeding. Your doctor or pharmacist can discuss with you the risks and benefits involved.

Taking other medicines

Tell your pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

If you are taking more than one medicine, they may interact with each other. At times your prescriber may decide to use medicines that interact, in other cases this may not be appropriate.

The decision to use medicines that interact depends on your specific circumstances. Your prescriber may decide to use medicines that interact, if it is believed that the benefits of taking the medicines together outweigh the risks. In such cases, it may be necessary to alter your dose or monitor you more closely.

Tell your prescriber the names of all the medicines that you are taking so that they can consider all possible interactions. This includes all the medicines which have been prescribed by your GP,

hospital doctor, dentist, nurse, health visitor, midwife or pharmacist. You must also tell your prescriber about medicines which you have bought over the counter without prescriptions.

The following medicines may interact with Saffid Capsule 300mg:

- Anticoagulants, Coumarin, Indandione derivative
Concurrent use may significantly increase the blood thinning effect of these medications, your doctor may need to do some blood test to adjust the anticoagulants dose.
- Chenodiol or ursodiol (used to treat gallstone)
Effects may be decreased when used concurrently with gemfibrozil.
- Lovastatin (cholesterol lowering agent)
Concurrent use may be associated with an increase in creatine kinase (CK) concentrations and myoglobinuria (presence of myoglobin in the urine). This leads to acute kidney failure that may be seen as early as 3 weeks or as late as several months after initiation of combined therapy.

If you are taking Saffid Capsule 300mg and one of the above medicines or types of medicines, make sure your prescriber knows about it.

Complementary preparations and vitamins

Medicines can interact with complementary preparations and vitamins. In general, there is not much information available about interactions between medicines and complementary preparations or vitamins.

If you are planning to take or are already taking any complementary preparations and vitamins you should ask your prescriber whether there are any known interactions with Gemfibrozil.

SAFFID CAPSULE 300MG

Gemfibrozil 300mg

Your prescriber can advise whether it is appropriate for you to take combinations that are known to interact. They can also discuss with you the possible effect that the complementary preparations and vitamins may have on your condition.

If you experience any unusual effects while taking this medicine in combination with complementary preparations and vitamins, you should tell your prescriber.

How to use Saffid Capsule 300mg

Follow all directions given to you by your pharmacist or doctor carefully. They may differ from the information contained in this leaflet.

If you do not understand the directions on your prescription label, ask your pharmacist, nurse or doctor to explain the instructions to you.

How much to use

Usual adult dose:

Orally 4 capsules (1.2g) a day in two divided dose thirty minutes before the morning and evening meals.

When to use it

Thirty minutes before the morning and evening meals.

How long to use it

Continue taking your medicine for as long as your doctor tells you.

It is important to keep taking your medicine even if you feel well. Suddenly stopping your medicine may cause your original condition to return. This is why you must speak to your prescriber if you are having any problems taking your medicine.

Your doctor may withdraw the treatment of Saffid Capsule 300mg if the response is inadequate after 3 months or if the results of liver function test show significant abnormalities.

If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

If you use too much (overdose)

Contact your doctor or go to your nearest hospital immediately. Do not drive if you have taken too many.

While you are using it

Things you must do

Talk to your pharmacist or doctor if your symptoms do not improve.

Your pharmacist or doctor will assess your condition and decide if you should continue to take the medicine.

Things you must not do

Do not take Saffid Capsule 300mg to treat any other complaints unless your pharmacist or doctor tells you to.

Do not give your medicine to anyone else, even if they have the same condition as you.

Do not take more than the recommended does unless your pharmacist or doctor tells you to.

Things to be careful of

Be careful driving or operating machinery until you know how Saffid Capsule 300mg affects you.

Do not drink alcohol while taking this medication.

Side effects

Tell your pharmacist or doctor if you do not feel well while you are taking Saffid Capsule 300mg. this medicine helps most people, but it may have unwanted side effects in a few people. All medicines can have side effects.

Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects. Do not be alarmed by the

following list of side effects. You may not experience any of them.

Ask your pharmacist doctor to answer any questions you may have.

Everyone's reaction to a medicine is different. It is difficult to predict which side effects will you have from taking a particular medicine, or whether you will not have any side effects at all. The important thing is to tell your prescriber or pharmacist with your medicine.

The reported side effects are:

- stomach pain, gas or heartburn
- diarrhoea
- nausea or vomiting
- skin rash
- tiredness
- anaemia or leucopenia (cough or hoarseness: fever or chills, lower back or side pain, painful or difficult urination)
- gallstones (severe stomach pain with nausea and vomiting)
- myositis (muscle pain, unusual tiredness or weakness)

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.moh.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Saffid Capsule 300mg

Storage

Keep all medicines out of the sight and reach of children.

Do not use Saffid Capsule 300mg after the expiry date stamped on the pack.

Store below 30°C. Protect from light

SAFFID CAPSULE 300MG

Gemfibrozil 300mg

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

What it looks like

A maroon and white '0' size elongated capsule containing white powder.

Dosage Form: Oral Solid

Pack Size:

In blister of 3 x 10's

Ingredients:

Active ingredients:

Saffid Capsule 300mg contains
Gemfibrozil 300mg

Inactive ingredients:

Microcrystalline Cellulose (101),
Pregelatinized Starch, Croscarmellose
Sodium, Colloidal Silicon Dioxide
(Aerosil 200) and Magnesium Stearate.

MAL Number:

MAL22106021AZ

Manufacturer

InspiraPharma Sdn Bhd
(200401023395),
Lot 24 & 25, Jalan 8,
Kawasan Perusahaan Bakar Arang,
08000 Sungai Petani,
Kedah Darul Aman, Malaysia.

Product Registration Holder

InspiraPharma Sdn Bhd
(200401023395),
Lot 24 & 25, Jalan 8,
Kawasan Perusahaan Bakar Arang,
08000 Sungai Petani,
Kedah Darul Aman, Malaysia.

Date of revision:

06/07/2026

Serial Number:

NPRA (R2) 22/026