

IDAMAN SILVER SULFADIAZINE CREAM 1% W/W

Silver Sulfadiazine (1% w/w)

What is in this leaflet

1. What Idaman Silver Sulfadiazine Cream 1% w/w is used for
2. How Idaman Silver Sulfadiazine Cream 1% w/w works
3. Before you use Idaman Silver Sulfadiazine Cream 1% w/w
4. How to use Idaman Silver Sulfadiazine Cream 1% w/w
5. While you are using Idaman Silver Sulfadiazine Cream 1% w/w
6. Side effects
7. Storage and Disposal of Idaman Silver Sulfadiazine Cream 1% w/w
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial number

What Idaman Silver Sulfadiazine Cream 1% w/w is used for

Silver Sulfadiazine is used:

- To prevent and treat infection in burn wounds
- As an aid to the short-term treatment in other skin conditions, such as leg ulcers
- As an aid to prevent infection in skin grafting.

How Idaman Silver Sulfadiazine Cream 1% w/w works

Silver Sulfadiazine has broad antimicrobial activity against Gram-positive and Gram-negative bacteria. It works by killing bacteria or stopping their growth.

Before you use Idaman Silver Sulfadiazine Cream 1% w/w

- When you must not use it

- Hypersensitive to silver sulphadiazine, other sulphonamides or any of the other ingredients in the preparation such as cetyl alcohol and propylene glycol.
- Pregnant women approaching or at term
- Premature infants
- Newborn infants during the first 2 months of life.

- Before you start to use it

Please tell your doctor:

- If you have kidney or liver problem
- If you have extensive burns
- If you have glucose-6-phosphate dehydrogenase (G6PD) deficiency

Absorption of propylene glycol which is contained in Silver Sulfadiazine may interfere with some laboratory test results.

If you are not sure whether you should start using this medicine, talk to your doctor or pharmacist first.

- Taking other medicines

As the silver content of silver sulphadiazine may inactivate topical proteolytic enzymes and debriding agents (a type of medicine used externally to clean and treat wounds), their concomitant use may be inappropriate.

The effects of oral hypoglycaemic agents (anti-diabetic medication) and phenytoin (anticonvulsant medication) may be potentiated when large areas of burns are treated. Tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription such as supplements or traditional medicines.

Pregnancy and lactation

Pregnancy and lactation

There are no adequate and controlled study being done in pregnant women, therefore the cream should be used during pregnancy only if clearly needed. Silver sulphadiazine should also be used with caution in nursing women. Ask your doctor or pharmacist before taking or using any medicines.

How to use Idaman Silver Sulfadiazine Cream 1% w/w

- How much to use

Silver Sulfadiazine is applied topically to cleansed burned/affected areas once or twice daily approximately 1.5mm thick, using a sterile-gloved hand or spatula. Dressing is usually not required but it can be used.

- When to use it

It should be applied to the burn wound/affected area as long as there is a possibility of infection or prescribed by the doctor. Whenever necessary, the cream should be re-applied to any areas from which it has been removed.

- How long to use it

Treatment should be continued until satisfactory healing has occurred, or until the burn site is ready for grafting.

- If you forget to use it

If you miss an application, apply it as soon as possible. Do not use a double dose to make up for a missed dose.

- If you use too much (overdose)

There is little experience of overdosage with Silver Sulfadiazine but if accumulation of sulphadiazine occurs, discontinuance of this cream should be considered.

For people who are on long-term/high dose therapy or people with impaired kidney function, it may be necessary to continuously monitor the possibility of crystal found in the urine.

While you are using Idaman Silver Sulfadiazine Cream 1% w/w

- Things you must do

- Always use Silver Sulfadiazine exactly according to the instructions given to you by your doctor or pharmacists. Check with your doctor or pharmacist if you are unsure.
- Tell your doctor or pharmacist immediately if you notice any side effects, or if the side effects gets serious or anything else that make you feel unwell.
- If you become pregnant, or plan on becoming pregnant while using Silver Sulfadiazine, tell your doctor immediately.
- If you are about to be started on a new medicine, remind your doctor and pharmacist that you are using Silver Sulfadiazine.

IDAMAN SILVER SULFADIAZINE CREAM 1% W/W

Silver Sulfadiazine (1% w/w)

- Things you must not do

Do not use Silver Sulfadiazine after the expiry date which is stated on the carton and tube.

- Things to be careful of

Be careful while driving or operating machinery until you know how Silver Sulfadiazine affects you.

Side effects

Like all medicines, Silver Sulfadiazine can cause side effects, although not everybody gets them.

Tell your doctor if you have any of these symptoms:

- Common: leukopenia (reduction in the number of white cells), application site burning, pruritus, application site rash (including eczema and contact dermatitis)
- Rare: argyria (exposure to or ingestion of silver)
- Very rare: renal failure

If you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Idaman Silver Sulfadiazine Cream 1% w/w

- Storage

Store below 30°C. Protect from light.

Once opened, product needs to be used within 6 months.

For external use only.

Keep out of reach of children.

- Disposal

If your doctors tell you to stop taking Silver Sulfadiazine or the expiry date has passed, ask your pharmacist the proper way to dispose the remaining products.

Product Description

- What it looks like

A smooth, soft, white homogenous cream.

- Ingredients

Active ingredient:

Silver Sulfadiazine

Inactive ingredients:

- White Soft Paraffin
- Light Liquid Paraffin
- Cetostearyl Alcohol
- Cetomacrogol 1000
- Chlorocresol
- Disodium Edetate
- Butylated Hydroxytoluene
- Povidone (Plasdone K 30)
- Purified Water
- Phosphoric Acid
- Disodium Hydrogen Phosphate (Anhydrous)

MAL Number

MAL25036010AZ

Product Registration Holder & Manufacturer

INSPIRAPHARMA SDN BHD
(200401023395)

Lot 24 & 25, Jalan 8, Kawasan
Perusahaan Bakar Arang,
08000 Sungai Petani,
Kedah Darul Aman, Malaysia.

Date of Revision

25 June 2026

Serial Number

NPRA(R2)23/133