

# UNI-PAIN LINIMENT

Methyl Salicylate (25% w/v)

## What is in this leaflet

1. What UNI-PAIN is used for
2. How UNI-PAIN works
3. Before you use UNI-PAIN
4. How to use UNI-PAIN
5. While you are using UNI-PAIN
6. Side effects
7. Storage and disposal of UNI-PAIN
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What UNI-PAIN is used for

It is indicated in relieving minor muscular and joints pains.

## How UNI-PAIN works

UNI-PAIN contains methyl salicylate which has the properties to reduce pain and inflammatory conditions.

## Before you use UNI-PAIN

### -When you must not use it

It should not be used if you had known to be hypersensitive to any of the components of the formulation.

### -Before you start to use it

#### Pregnancy and lactation

Talk to your doctor before using this medicine if you are pregnant, might become pregnant or are breastfeeding.

### -Using other medicines

Inform your doctor or pharmacist before using this medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

## How to use UNI-PAIN

### -How much to use

Apply a little to the affected part and gently massage. To be applied 3 or 4 times a day.

### -When to use it

Use as directed by your doctor or pharmacist.

### -How long to use it

Continue using UNI-PAIN as your doctor recommended. If symptoms persist, consult your doctor.

### -If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Use the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to use the medicine and skip the missed dose. Do not use a double dose to make up for the missed dose.

### -If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

## While you are using UNI-PAIN

### -Things you must do

Inform your doctor of all the medications you are using.

### - Things you must not do

- Do not use more than the recommended dose unless your pharmacist or doctor tells you.
- Do not use any new medicines without consulting your doctor or pharmacist.
- Do not give UNI-PAIN to anyone else, even if they have the same symptoms or condition as you.

### -Things to be careful of

- Discontinue use and consult a doctor if symptoms persist for more than 7 days.
- Do not apply to wounds or damaged skin
- Do not apply over large areas of the body or bandage.
- This product contains methyl salicylate and when applied or rub on to the skin, can be absorbed through the skin into the blood. For patients using warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding.

## Side effects

Like all medicines, UNI-PAIN can cause side effects such as skin irritation, redness or swelling, burning or stinging sensation.

If side effects persist, do not use the medicine and refer to the doctor.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npa.gov.my](http://npa.gov.my) [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

## Storage and disposal of UNI-PAIN

### -Storage

Store below 30°C, in a dry place, protected from direct sunlight. Keep out of reach of children.

### -Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## Product description

### -What it looks like

Yellow colour liniment with a characteristic odour.

### -Ingredients

#### -Active ingredient

Methyl salicylate.

#### -Inactive ingredients

Menthol, Methyl paraben, Alcohol denatured and palm oil.

### -MAL number

MAL06021263XC

## Product Registration Holder

UNIMED SDN. BHD.

53, Jalan Tembaga SD 5/2B,  
Bandar Sri Damansara,  
52200 Kuala Lumpur, Malaysia.

## Manufactured by

KCK Pharmaceutical Industries Sdn. Bhd.

No. 5 & 7 Hala Beringin, Diamond Valley Industrial Park, 11960 Bayan Lepas, Pulau Pinang, Malaysia.

## Date of revision

24/06/2024

## Serial Number

NPRA(R2/1)240621/1292