

ONEGABA CAPSULES

Gabapentin (300mg, 400mg)

What is in this leaflet

1. What ONEGABA is used for
2. How ONEGABA works
3. Before you use ONEGABA
4. How to use ONEGABA
5. While you are using it
6. Side effects
7. Storage and Disposal of ONEGABA
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What ONEGABA is used for

Onegaba Capsule is used:

- together with other medicines to help treat partial seizures (fits that start in one part of the brain) in adults and children who are 3 years old or older.
- to treat nerve pain in adults (18 years and older), such as pain from diabetes, shingles, or trigeminal neuralgia (a nerve pain in the face).

How ONEGABA works

Onegaba Capsule contains gabapentin as active ingredient, which is an anticonvulsant. Gabapentin works by calming overactive nerves. It reduces the extra electrical signals in the brain that trigger seizures. In nerve pain, it stops damaged nerves from sending the wrong pain messages to the brain

Before you use ONEGABA

- When you must not use it

Do not take Onegaba:

If you are allergic (hypersensitive) gabapentin or any other ingredients of this medicine (listed in Product Description)

- Before you start to use it

Tell your doctor if you have:

- breathing problem
- kidney disease
- other nerve/brain conditions

Pregnancy and lactation

- Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine
- Gabapentin must only be used during pregnancy if the benefits is more than the risks to the fetus.
- Gabapentin is secreted into breast milk and may affect your baby. Your doctor will discuss the risks and benefits of breast feeding while you're taking gabapentin.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Inform your doctor or pharmacist if you are taking:

- Antacids containing aluminium or magnesium (for heartburn/indigestion). Take Gabapentin at least 2 hours after antacids.
- Cimetidine (for stomach ulcers/acid reflux)
- Other epilepsy medications
- Morphine (opioid type painkiller)

How to use ONEGABA

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- How much to use

For epilepsy

- Adults and children aged 12 and above: The usual dose ranges from 900 mg to 3,600 mg per day, divided into three separate doses.
- Children age 3–12 years: The dose will be determined by the doctor based on the child's body weight.

For nerve pain in adults:

- The starting dose is usually 300 mg three times a day (total 900 mg/day).
- The doctor may increase the dose step-by-step if needed, up to 3,600 mg per day (in 3 divided doses).

For people with kidney issues:

Your doctor will adjust your dose according to your condition if you are undergoing hemodialysis or have kidney impairment.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

- Onegaba can be taken with or without food.
- Take three times per day or as directed by your doctor or pharmacist.
- Take each dose on time and do not leave more than 12 hours between doses.

- How long to use it

Take this medicine for the duration that your doctor recommends

- If you forget to use it

If you forget to take a dose, take it as soon as you remember unless it is time for your next dose. Do not take a double dose to make up for a forgotten dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many capsules may cause dizziness, double vision, slurred speech, drowsiness, lethargy, and mild diarrhea.

While you are using it

- Things you must do

- Take your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are taking Onegaba.
- Tell your doctor immediately if you become pregnant while using this medication.
- Take Onegaba only with a doctor's prescription.

- Things you must not do

- Do not take any new medicines without consulting your doctor.
- Do not give Onegaba to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Driving and using machines

- This medicine may affect your ability to drive or use machines. If the tablets make you feel drowsy, dizzy or affect your vision, do not drive or use machines and contact your doctor immediately.
- Do not stop taking Onegaba suddenly, as stopping too quickly may cause seizures to return.
- Watch out for any breathing problems that occurred while taking this medicine. If this happens, see your doctor as soon as possible.
- Gabapentin can sometimes cause false positive urine protein test results with certain dipstick tests. A more accurate test should be used if needed.

ONEGABA CAPSULES

Gabapentin (300mg, 400mg)

Side effects

Like all medicines, ONEGABA can cause side effects, although not everybody gets them.

Very common:

- viral infection
- drowsiness
- dizziness
- loss of muscle control and balance.

Common:

- respiratory infection
- urinary tract infection
- middle ear infection
- low white blood cell count
- eating disorder
- increased appetite
- hostility
- confusion and emotional liability
- depression
- anxiety
- nervousness
- thinking abnormal
- seizure
- increase in muscular activity
- motor speech disorder
- amnesia
- tremor
- sleep disorder
- headache
- visual disturbances such as lazy eye (reduced vision in one eye caused by abnormal visual development early in life) and double vision
- sensation of feeling off balance
- high blood pressure, widening of blood vessels
- difficulty breathing
- bronchitis (inflammation of the lining of lungs)
- Pharyngitis (sore throat)
- Cough
- Rhinitis (hay fever)
- Vomiting
- Nausea
- Dental abnormalities
- Gingivitis (inflammation of the gums)
- Diarrhea
- Abdominal pain
- Indigestion
- Constipation
- Dry mouth or throat
- Farting
- Facial swelling
- Purpura most often described as bruises resulting from physical trauma, rash, itch and acne
- incontinence (the loss of bladder control)
- Joint pain
- Muscle pain

- Back pain
- Twitching (small muscle contractions in the body)
- impotence (inability for a man to get and maintain an erection)

Rare:

- low levels of platelets
- allergic reactions
- hallucinations
- movement disorder
- ringing in the ears
- pancreatitis (inflammation of the pancreas)
- Stevens-Johnson syndrome (rare, serious disorder of the skin and mucous membranes),
- Angioedema (severe swelling beneath the skin),
- Erythema multiforme (skin condition),
- Alopecia (spot baldness)
- acute kidney failure
- Hepatitis (inflammation of the liver)
- Jaundice (condition that causes skin and the whites of the eyes to turn yellow)
- feelings or sensations that your heart is pounding or racing

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Center for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of ONEGABA

- Storage

Do not store above 30°C.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

ONEGABA 300

Yellow hard gelatin capsule. It has a 'D' imprinted on the yellow cap and '03' on the yellow body. The capsule is filled with a white to off-white crystalline powder.

ONEGABA 400

Orange hard gelatin capsule. It has a 'D' imprinted on the orange cap and '04' on

the orange body. The capsule is filled with a white to off-white crystalline powder.

- Ingredients

- Active ingredient

Gabapentin

- Inactive ingredients

Maize starch, Purified Talc, Empty Hard Gelatin Capsule

- MAL number(s):

MAL09020987AZ

(Onegaba 300mg Capsules)

MAL09020988AZ

(Onegaba 400mg Capsules)

Manufacturer

AUROBINDO PHARMA LIMITED,
Unit III, Sy. No. 313 & 314,
Bachupally Mandal, Medchal -
Malkajgiri District, Telangana State,
India

Product Registration Holder

UNIMED SDN BHD

No. 53, Jalan Tembaga SD5/2B

Bandar Sri Damansara

52200 Kuala Lumpur

Date of revision

27/10/2025

Serial Number

NPRA(R4/1)100925/00069