

FENO-TG 160 MG TABLET

Fenofibrate 160 mg

What is in this leaflet

1. What Feno-TG is used for
2. How Feno-TG works
3. Before you use Feno-TG
4. How to use Feno-TG
5. While you are using it
6. Side effects
7. Storage and Disposal of Feno-TG
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Feno-TG is used for

FENO-TG is used to lower the level of fats (lipids) in the blood. For example the fats known as triglycerides. It is used, alongside a low fat diet and other nonmedical treatments such as exercise and weight loss, to lower levels of fats in the blood.

How works

FENO-TG works through the activation of a cell nuclear receptor called peroxisome proliferator-activated receptor alpha, which reduces the amount of triglycerides and bad cholesterol made in the body and increases the good cholesterol.

Before you use Feno-TG

- When you must not use it

Do not take Feno-TG if:

- You have an allergy to FENO-TG or any ingredients of it and Gemfibrozil

- Before you start use it

Tell your doctor if:

- You have any allergies to any other medicines or any other substances, such as food, preservatives or dyes.
- You are pregnant or intend to become pregnant and breast-feeding or intend to breast-feed.
- You have or have had any other medical conditions, including: Kidney problems and muscular aching, tenderness or weakness not caused by exercise.

Pregnancy and lactation

Ask your doctor or pharmacist for advice before taking this medicine if you are pregnant or breast-feeding.

- Taking other medicines

Please tell your doctor if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

In particular please let your doctor know if you already take:

- Anti-coagulants to thin your blood , other drugs used to control lipid (fat) levels in the blood and Ciclosporin (an immunosuppressant)

How to use Feno-TG

- How much to use

Always take FENO-TG 160 mg exactly as your doctor has told you. The usual dose for adults is one tablet of FENO-TG 160 mg a day. You should check with your doctor if you are not sure. Tablet should be swallowed during a meal with water.

- When to use it

FENO-TG can be taken at any time of the day. Any dietary measures started before treatment with FENO-TG should be continued. Take FENO-TG at about the same time each day. Taking tablets at the same time each day will have the best effect. It will also help you remember when to take the tablets.

- How long to use it

Do not forget that fenofibrate to be effective, it needs to be taken very regularly and for as long as your doctor has recommended, even if this duration is very long time. Do not stop taking this medicine unless your doctor has told you.

- If you forget to use it

Do not take a double dose to make up for a forgotten tablet. Take the next dose with your next meal and then carry on taking your tablets as usual. If you are worried about this talk to your doctor.

- If you use too much (overdose)

If you accidentally take more dose than you should or if someone else has taken your medicine, contact your nearest hospital casualty department or tell your doctor immediately.

While you are using it

- Things you must do

- Have your blood fats checked when requested by your doctor to make sure FENO-TG is working.
- If you become pregnant while you are taking FENO-TG stop taking it and contact your doctor immediately.
- If you are about to be started on any new medicine tell your doctor and pharmacist that you are taking FENO-TG

- Things you must not do

Do not give FENO-TG to anyone else, even if they have the same condition as you and do not use FENO-TG to treat any other complaints unless your doctor tells you to.

- Things to be careful of

Discontinue use if irritation or sensitivity reaction develops.

Avoid drinking large quantities of alcohol. Drinking large quantities of alcohol may increase your chance of FENO-TG causing liver problems.

Side effects

Like all medicines, FENO-TG can cause side effects although not everybody gets them. Tell your doctor immediately if you experience any of the following symptoms after taking this medicine as the symptoms can be severe.

- Abdominal pain
- Back pain
- Headache
- Diarrhea
- Nausea
- Constipation
- Rhinitis
- Joint disorder
- Leg cramp

FENO-TG

Fenofibrate 160 mg

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.moh.gov.my (Public → Reporting Medicinal Problems/ Sides Effects / AEFI / Vaccine Safety).

Storage and Disposal of Feno-TG

- Storage

Store below 30°C, Protected from light.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Orange colored, round shape, biconvex film coated tablet.

- Ingredients

- Active ingredient
Fenofibrate 160 mg

- Inactive ingredients

Disodium Edetate, Microcrystalline Cellulose, Sodium Starch Glycollate, Starch, Isopropyl Alcohol, Polyvinylpyrrolidone K-30, Crospovidone, Sodium Lauryl Sulphate, Magnesium Stearate, Talc, Ethyl Cellulose, Hydroxypropylmethyl Cellulose, Lake of Sunset Yellow (Dye Content 12 – 17%), Methylene Chloride, Propylene Glycol, Titanium Dioxide.

- MAL number(s):

MAL11045012AZ

Manufacturer

Troikaa Pharmaceuticals Ltd.
C-1, Sara Industrial Estate, Selaqui,
Dehradun-248197, Uttarakhand, India.

Product Registration Holder

Unimed Sdn Bhd
53, Jalan Tembaga SD 5/2B,
Bandar Sri Damansara,
52200, Kuala Lumpur, Malaysia.

Date of revision

17/07/2022

Serial Number: NPRA(R2/1)220519/891