


<p>Mfg. Lic. No. : MB/06/338</p> <p>B. No.</p> <p>Mfg. Date</p> <p>Expiry Date</p> <p>Keep all medicines out of reach of children. Jauhkan dari kanak-kanak "Controlled Medicine / Ubat Terkawal"</p> <p>MAL No.: MAL11030101AZ</p> <p>Holder & Imported by: Unimed Sdn Bhd 53, Jalan Tembaga SD 5/2B, Bandar Sri Damansara, 52200 Kuala Lumpur.</p> <p>Manufactured in India by: FDC Limited Village: Khol-Bhood, Tehsil: Nalagarh, Baddi-173 205, Dist.: Solan, H.P.</p>	<p>60 ml</p> <p>ZEFU POWDER FOR ORAL SUSPENSION</p> <p>125 mg/5ml</p> <p>Cefuroxime Axetil for Oral Suspension</p>  <p>2000007391</p> <p>FDC</p>	<p>COMPOSITION: Each 5 ml (one teaspoonful) of reconstituted suspension contains Cefuroxime Axetil equivalent to Cefuroxime125 mg Excipients q.s.</p> <p>DOSAGE: As directed by the Physician.</p> <p>MAKE UP THE VOLUME UP TO THE ARROW MARK ON THE LABEL →</p> <p>DO NOT FREEZE SHAKE WELL BEFORE EACH DOSE Store in a dry place at a temperature not exceeding 25°C.</p> <p>DIRECTIONS FOR USE: Tap the bottle to loosen the powder. Slowly add boiled and cooled water upto arrow mark on the label and shake well. Add water if necessary to adjust the volume upto the arrow mark.</p> <p>The reconstituted suspension should be stored in a refrigerator and should be used within 7 days.</p>
---	---	--

90 mm

Arrow Distance from Bottom 35 mm

<p>Mfg. Lic. No. : MB/06/338</p> <p>B. No.</p> <p>Mfg. Date</p> <p>Expiry Date</p> <p>Keep all medicines out of reach of children. Jauhkan dari kanak-kanak "Controlled Medicine / Ubat Terkawal"</p> <p>MAL No.: MAL11030101AZ</p> <p>Holder & Imported by: Unimed Sdn Bhd 53, Jalan Tembaga SD 5/2B, Bandar Sri Damansara, 52200 Kuala Lumpur.</p> <p>Manufactured in India by: FDC Limited Village: Khol-Bhood, Tehsil: Nalagarh, Baddi-173 205, Dist.: Solan, H.P.</p>	<p>60 ml</p> <p>ZEFU POWDER FOR ORAL SUSPENSION</p> <p>125 mg/5ml</p> <p>Cefuroxime Axetil for Oral Suspension</p>  <p>2000007391</p> <p>FDC</p>	<p>COMPOSITION: Each 5 ml (one teaspoonful) of reconstituted suspension contains Cefuroxime Axetil equivalent to Cefuroxime125 mg Excipients q.s.</p> <p>DOSAGE: As directed by the Physician.</p> <p>MAKE UP THE VOLUME UP TO THE ARROW MARK ON THE LABEL →</p> <p>DO NOT FREEZE SHAKE WELL BEFORE EACH DOSE Store in a dry place at a temperature not exceeding 25°C.</p> <p>DIRECTIONS FOR USE: Tap the bottle to loosen the powder. Slowly add boiled and cooled water upto arrow mark on the label and shake well. Add water if necessary to adjust the volume upto the arrow mark.</p> <p>The reconstituted suspension should be stored in a refrigerator and should be used within 7 days.</p>
---	---	--

Enlarged