

# IOBRIM 0.15% w/v EYE DROPS

Consumer Medication Information Leaflet (RiMUP)

(Brimonidine Tartrate)

## What is in this leaflet

1. What **IOBRIM** is used for
2. How **IOBRIM** works
3. Before you use **IOBRIM**
4. How to use **IOBRIM**
5. While you are using it
6. Side effects
7. Storage and Disposal of **IOBRIM**
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial No.

## What IOBRIM is used for

IOBRIM is used for lowering pressure in the eye.

## How IOBRIM works

It can be used alone, or with other eye drops, if single use does not work to reduce the pressure in the eye for the treatment of open angle glaucoma or ocular hypertension.

## Before you use IOBRIM

- When you must not use it

### Do not take IOBRIM:

- If you are hypersensitive to IOBRIM or to any of the ingredients in IOBRIM
- If you are breastfeeding.

### Before you start to use it

- Before you start to use it, please inform your doctor

If you are suffering or experiencing depression, mental retardation, less blood flow to the brain, heart disease, less blood flow to the rest of the body or high blood pressure.

If you are having or had kidney failure or liver problem

If given to children between the of 2 and 12 years old because IOBRIM is not recommended for this age group

If you are pregnant or breastfeeding

Ask your doctor or pharmacist for advise before taking this medication Inform your doctor if you are pregnant or

planning for pregnancy. IOBRIM cannot be used in breastfeeding.

### Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines or have just started new medications, including non-prescription medicines.

Tell your doctor if you are particularly taking:

- Anaesthetic medication, sedatives, opioids, barbiturates or frequent alcoholic drinker
- Drugs that blocks beta receptors (eye drops and consumables), blood pressure lowering medication and/or cardiac glycosides.
- Tricyclic antidepressants
- If change of dose of any medication, this can affect your treatment with IOBRIM

## How to use IOBRIM

### How much to use

Always use IOBRIM as instructed by the doctor. You have to check with your doctor or your pharmacist if you are not sure.

Instill one drop in the affected eye 3 times daily, approximately 8 hours apart.

### When to use it

For effectiveness, please use it everyday.

### How long to use it

Do not stop using it unless your doctor says so.

If you have any further queries about the use of this product, please consult with your doctor or pharmacist.

### If you forget to use it

If you forget to take a dose, take the missed dose as soon as you remember. If it is almost time for your next dose anyway, forget about the missed dose. Wait and take the next dose at the regular time.

### If you use too much (overdose)

Adult

No cases have been reported in too much use of IOBRIM among adults. This is not possible when given as eye drop.

If IOBRIM has been accidentally swallowed or if you have used IOBRIM more than you should, please contact your doctor immediately.

## While you are using it

### Things you must do

### Instructions For Use

IOBRIM is a eye drop. Wash your hands before using the eye drop. Your prescription will state how much to use at each dose. If you are using IOBRIM with other eye drops, wait for 5 minutes before instilling the second eye drop.

Use the eye drop as follows:

- Tilt your head back, gaze upward looking at the ceiling.
- Slowly pull down the lower eyelid to make a pouch.
- Place the dropper directly over eye and gently squeeze the bottle to release a drop into your eye.
- Look downward and gently close your eye for 1 to 2 minutes. Press your finger against the inside corner of the closed eye, and hold for 1 minute.
- Try not to blink and do not rub the eye.
- Repeat the above steps for the other eye if necessary.
- Replace and tighten the cap immediately after use.

Avoid touching the bottle tip against the eye. Screw tightly once after use.

### Things you must not do

### Driving or using machines

- IOBRIM can caused blurry eye or unusual. This effect may get worst at night or in low light area.
- IOBRIM may cause drowsiness or fatigue in some patients
- If you experience any of these symptoms, do not drive or use any machines until the effects goes off.

### -Things to be careful of

Driving or using machines

This medication can affect your ability to drive or use any machines. If the medication gives you pain, headache, please do not drive or use any machines and consult your doctor immediately.

**Side effects**

Side effects: Allergic conjunctivitis, hyperemic conjunctiva and eye pruritus.

Worst Side effect: Burning sensation, presence of folliculosis conjunctiva, high blood pressure, dry mouth and blurred vision.

Incident: Allergic reaction, asthenia, blepharitis, bronchitis, oedema conjunctiva, conjunctiva bleeding, conjunctivitis, cough, headache, digestion problem, shortness of breath, excessive watering of the eye, dry eye, eye pain, eyelid swelling, eyelid erythema, flu syndrome, follicle conjunctivitis, foreign body sensation, headache, sore throat, photophobia, rashes, rhinitis, sinus infection, sinus, burning sensation, superficial punctate keratopathy, visual disruption, floater vitreous and worsening of the visual.

Following worst reactions: corneal erosion, insomnia, dry nose, change in sleeping pattern, loss of taste.

Like all medicines, IOBRIM EYE DROP can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects listed above or even if it is not listed here.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my) (Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

**Storage and Disposal of IOBRIM EYE DROP**

-Storage

Store below 30°C. To avoid contamination, store in a closed container. Do not touch the tip of the bottle dropper. The content is sterile unopened.

KEEP OUT OF REACH OF CHILDREN

After first opening: Store below 30°C. Discard after 28 days of opening

-Disposal


Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

**Product Description**

What it looks like

IOBRIM EYE DROPS is a clear, greenish yellow 5 mL solution.

Mode of use:



Ingredients

Active ingredients  
Brimonidine Tartrate Ph. Eur.

Inactive ingredients

Benzalkonium Chloride Ph. Eur.  
(As preservative)  
Aqueous Buffered (As Vehicle)

Number:

MAL\*\*\*\*\*

**Manufacturer**

FDC Limited  
B-8, MIDC Industrial Area, Waluj,  
Aurangabad- 431 136, Maharashtra

**Product Registration Holder**

Unimed Sdn Bhd No 53, Jalan Tembaga  
SD 5/2B, Bandar Sri Damansara, 52200  
Kuala Lumpur.

**Date of revision**

19/11/2021

**Serial Number:**NPRA(R2)21/061