

In Table 4, adverse reactions are presented by frequency category based on incidence in clinical trials or epidemiological studies, when known. The frequency category “not known” is used for adverse reactions for which no valid estimate of the incidence rate can be derived from clinical trials.

Table 4. Adverse Reactions Identified During Postmarketing Experience with RAPIFEN by Frequency Category Estimated from Clinical Trials or Epidemiologic Studies	
Immune System Disorders	
Not known	Hypersensitivity (including anaphylactic reaction, anaphylactoid reaction, and urticaria)
Psychiatric Disorders	
Not known	Disorientation
Nervous System Disorders	
Not known	Loss of consciousness ^a , Convulsion, Myoclonus
Eye Disorders	
Not known	Miosis
Cardiac Disorders	
Not known	Cardiac arrest
Respiratory, Thoracic and Mediastinal Disorders	
Not known	Respiratory arrest, Cough
Uncommon	Respiratory depression ^b
Skin and Subcutaneous Tissue Disorders	
Not known	Erythema, Rash
General Disorders and Administration Site Conditions	
Not known	Pyrexia
^a Postoperative period	
^b Including fatal outcome	

Serotonin syndrome (see *Warnings and Precautions*)
Adrenal insufficiency (see *Warnings and Precautions*)
Androgen deficiency
Cases of androgen deficiency have occurred with chronic use of opioids. Chronic use of opioids may influence the hypothalamic-pituitary-gonadal axis, leading to androgen deficiency that may manifest as low libido, impotence, erectile dysfunction, amenorrhea, or infertility. The casual role of opioids in the clinical syndrome of hypogonadism is unknown because the various medical, physical, lifestyle, and psychological stressors that may influence gonadal hormone levels have not been adequately controlled for in studies conducted to date. Patients presenting with symptoms of androgen deficiency should undergo laboratory evaluation.

Infertility
Chronic use of opioids may cause reduced fertility in females and males of reproductive potential. It is not known whether these effects on fertility are reversible.

Overdose
Signs and Symptoms
The manifestations of RAPIFEN overdose are an extension of its pharmacologic actions. Respiratory depression which can vary in severity from bradypnea to apnea may occur.

Treatment
In the presence of hypoventilation or apnea, oxygen should be administered and respiration should be assisted or controlled as indicated. A specific opioid antagonist should be used as indicated to control respiratory depression. This does not exclude the use of more immediate countermeasures. The respiratory depression may last longer than the effect of the antagonist; additional doses of the latter may therefore be required.

If depressed respiration is associated with muscular rigidity, an intravenous neuromuscular blocking agent might be required to facilitate assisted or controlled respiration.

The patient should be carefully observed; body warmth and adequate fluid intake should be maintained. If hypotension is severe or if it persists, the possibility of hypovolemia should be considered, and if present, it should be controlled with appropriate parenteral fluid administration.

PHARMACOLOGICAL PROPERTIES

Pharmacodynamic Properties
Pharmacotherapeutic group: opioid anesthetics, ATC code: N01AH02.

Mechanism of action
Alfentanil is a potent fast and short acting opioid analgesic, chemically related to fentanyl.

Pharmacodynamic effects
After intravenous administration of alfentanil action sets in almost instantly, the onset of action amounts to only one quarter of that of an equianalgesic dose of fentanyl. The maximum analgesic and respiratory depressant effect occurs within 1-2 minutes (30 minutes with morphine).

The duration of action of alfentanil is approximately one-third of that of an equianalgesic dose of fentanyl and is clearly dose-related. For analgesia lasting longer than 60 minutes, an infusion is preferable. Alfentanil's depressant effects on respiratory rate and alveolar ventilation are of shorter duration than those of fentanyl; in most cases the duration of analgesia exceeds that of the respiratory depression. The duration and degree of respiratory depression tend to be dose-related.

High doses (> 120 mcg/kg) of alfentanil induce sleep and can be used for induction of anesthesia. The induction is smooth, pain-free and devoid of cardiovascular and hormonal stress responses to intubation.

In common with other opioid analgesics, alfentanil can, depending upon the dose and speed of administration, cause muscle rigidity, as well as euphoria, miosis and bradycardia.

At doses up to 200 mcg/kg, alfentanil failed to produce a significant increase in histamine levels or clinical evidence of histamine release.

Recovery after alfentanil administration is typically rapid and smooth with a low incidence of post-operative nausea and vomiting.

All actions of alfentanil are reversed by a specific opioid antagonist.

Pharmacokinetic Properties
Alfentanil is a synthetic opioid with μ -agonist pharmacologic effects, used only intravenously.

Distribution
The sequential distribution half-lives of alfentanil are 0.4-2.2 min and 8-32 min. The low degree of ionization (11% at pH = 7.4) contributes to a rapid but limited tissue distribution. Reported volumes of distribution are 1.27-4.81 L (volume of distribution of the central compartment) and 12.1-98.2 L (volume of distribution at steady-state). Plasma protein binding of alfentanil is about 92%.

Metabolism
Alfentanil is mainly metabolized in the liver. Only 1% of unchanged alfentanil is found in urine. Metabolites are inactive and 70-80% of them are eliminated via the urine.

Elimination
Alfentanil is rapidly eliminated after intravenous administration. Terminal elimination half-lives of 83- 223 min have been reported. The plasma clearance in subjects below 40 years of age averages 356 mL/min, and decreases approximately 8% per decade increase above 40 years of age. Only 1% of unchanged alfentanil is found in urine. Once steady-state has been reached after infusion, the elimination half-life remains unaltered. When the administration is discontinued, the patient awakens rapidly without opioid after effects.

Special Populations
Pediatrics

The data in children are limited. The values for the pharmacokinetic parameters are shown in the table below.

	t1/2 β (hr)	CL (mL/kg/min)	Vdss (L/kg)
Preterm Neonates (0-27 days)			
Gestational age 25-40 weeks; <i>n</i> = 68	0.7-8.8	0.9-8.4	0.3-1.2
Term Neonates (0-27 days) Gestational age: 35-41 weeks; <i>n</i> = 18	4.1-5.5	1.7-3.2	0.5-0.8
Infants			
28 days - 23 months; <i>n</i> = 34	0.9-1.2	7.7-13.1	0.4-1.1
Children			
2-11 years; <i>n</i> = 32	0.7-1.3	4.7-10.2	0.2-1.0
Adolescents			
12-14 years; <i>n</i> = 3	1.1-1.9	5.5-7.4	0.3-0.6

- Note: Data for neonates, infants, and children are given as range of mean values.
- CL = clearance, Vdss = volume of distribution at steady state, t1/2 β = half-life in the elimination phase.

Protein binding in newborns is 75% and increases in children to 85%. Pharmacokinetic information on the use of alfentanil in children is limited. Alfentanil is metabolized by CYP3A4. CYP3A4 activity is low in neonates and increases after birth to reach 30 to 40% of adult levels at 1 month of age. Activity of CYP3A4 increases further to 45% at 6 months, 80% at 12 months and reaches adult levels at 6 years of age.

Hepatic Impairment
After administration of a single intravenous dose of 50 mcg/kg, the terminal half-life in cirrhotic patients is significantly longer than in controls. The volume of distribution remains unchanged. The free fraction of alfentanil increases in cirrhotic patients to 18.5% compared with 11.5% in controls. This increase in free fraction together with a reduction in clearance from 3.06 mL/min/kg in controls to 1.60 mL/min/kg in cirrhotic patients will result in a more prolonged and pronounced effect (see *Warnings and Precautions*).

Renal Impairment
The volume of distribution and clearance of the free fraction is similar in renal failure patients and healthy controls. The free fraction of alfentanil in patients with renal failure is increased to 12.4 to 19% compared with 10.3 to 11% in controls. This may result in an increase in clinical effect of alfentanil (see *Warnings and Precautions*).

NON-CLINICAL INFORMATION
Preclinical effects observed were only at exposures considered sufficiently in excess of the maximum human exposure indicating little relevance to clinical use.

Alfentanil has a wide safety margin. In rats the ratio of LD₅₀/ED₅₀ for the lowest level of analgesia, for alfentanil is 1080 compared with 4.6, 69.5 and 277 for pethidine, morphine and fentanyl respectively.

PHARMACEUTICAL PARTICULARS

List of Excipients
Sodium chloride
Water for injection

Incompatibilities
The injectable solution must not be mixed with other products.

If desired, RAPIFEN may be mixed with sodium chloride or glucose intravenous infusions. Such dilutions are compatible with plastic infusion sets. These should be used within 24 hours of preparation.

Shelf Life
36 months


Storage Conditions
Store below 30° C.
Keep out of the sight and reach of children.

Nature and Contents of Container
2 mL x 5 ampules

Instructions for Use/Handling

Wear gloves while opening ampule.

1. Maintain the ampule between thumb and index finger, leaving the tip of the ampule free.
2. With the other hand, hold the tip of ampule putting the index finger against the neck of ampule, and the thumb on the colored point in parallel to the identification colored ring(s).
3. Keeping the thumb on the point, sharply break the tip of ampule while holding firmly the other part of the ampule in the hand.



Accidental dermal exposure should be treated by rinsing the affected area with water. Avoid usage of soap, alcohol, and other cleaning materials that may cause chemical or physical abrasions to the skin.

MANUFACTURER
DEMO S.A Pharmaceuticals
21st km National Road,
Athens - Lamia, Krioneri Attiki, 14568,
Greece

DATE OF REVISION OF THE TEXT
17 June 2022

Parameters	Remarks	Parameters	Remarks
Product name/Generic name	Rapifen 5 x 2 ml PIL Malaysia	Market/Country	Malaysia
Strength	NA	Barcode (PZN/CIP/EAN/KD/SKUL/VNR)	NA
Component	Pack Insert	Material code	38886330
Dimensions	696 x 332 mm	Superseded Material code	NA
Specification	NA	Artwork number	NA
Font size & type	Helvetica World	MAH	NA
Pantone number	As shown	GTIN	NA
Cut margins/Peel off Margins	NA	Special instructions	NA
Unvarnished zone	NA	Anticounterfiet features	NA
Mfg site	NA	Other observation	NA
Reason for revision:	New Artwork	Pharma Code	440
Date: 17/06/2022			

Black

