

ONEGABA 600

Gabapentin 600 mg

What is in this leaflet

1. What ONEGABA 600 is used for
2. How ONEGABA 600 works
3. Before you use ONEGABA 600
4. How to use ONEGABA 600
5. While you are using it
6. Side effects
7. Storage and Disposal of ONEGABA 600
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What ONEGABA 600 is used for

ONEGABA 600 is used:

- together with other medicines to help treat partial seizures (fits that start in one part of the brain) in adults and children who are 3 years old or older.
- to treat nerve pain in adults (18 years and older), such as pain from diabetes, or shingles.

How ONEGABA 600 works

ONEGABA 600 contains gabapentin as active ingredient, which is an anticonvulsant. Gabapentin works by calming overactive nerves. It reduces the extra electrical signals in the brain that trigger seizures. In nerve pain, it stops damaged nerves from sending the wrong pain messages to the brain.

Before you use ONEGABA 600

- When you must not use it

Do not take ONEGABA 600:

- If you are allergic (hypersensitive) to gabapentin or any other ingredients of this medicine (Listed in Product Description).

- Before you start to use it

Tell your doctor if you have:

- breathing problem
- kidney disease
- other nerve/brain conditions

Pregnancy and lactation

- Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast feeding before using any medicine.
- Gabapentin must only be used during pregnancy if the benefits is more than the risks to the fetus.
- Gabapentin is secreted into breast milk and may affect your baby. Your doctor will discuss the risks and benefits of breast feeding while you're taking gabapentin.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Tell your doctor if you are taking, have recently taken alcohol, or recreational drugs, as these may affect how ONEGABA 600 works or increase the risk of side effects.

Inform your doctor or pharmacist if you are taking:

- Antacids containing aluminium or magnesium (for heartburn/indigestion). Take Gabapentin at least 2 hours after antacids.
- Cimetidine (for stomach ulcers/acid reflux).
- Other epilepsy medications such as phenobarbital, phenytoin, valproic acid, carbamazepine.
- Morphine (opioid type painkiller).
- Oral contraceptives containing norethindrone and/or ethinyl estradiol.
- Probenecid (medication for gout).

Your doctor may need to monitor you more closely or adjust your treatment. If you are taking morphine together with gabapentin, you will be monitored for excessive drowsiness or other signs of central nervous system depression, and the dose of either medicine may need adjustment.

How to use ONEGABA 600

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- How much to use

For epilepsy.

- Adults and children aged 12 and above: The starting dose will generally be between 300 mg and 900 mg each day. Thereafter, the dose may be increased as instructed by your doctor, up to a maximum of 3600 mg each day.
- Children aged 3 – 12 years: The dose will be determined by the doctor based on the child's body weight.

For nerve pain in adults:

- The starting dose is usually 300 mg three times a day (total 900 mg/day).
- The doctor may increase the dose step-by-step if needed, up to 3,600 mg per day (in 3 divided doses).

For people with kidney issues:

Your doctor will adjust your dose according to your condition if you are undergoing hemodialysis or have kidney impairment.

- When to use it

- ONEGABA 600 can be taken with or without food.
- Take three times per day or as directed by your doctor or pharmacist.
- Take each dose on time and do not leave more than 12 hours between dose

- How long to use it

Take this medicine for the duration that your doctor recommends.

- If you forget to use it

If you forget to take a dose, take it as soon as you remember unless it is time for your next dose. Do not take a double dose to make up for a forgotten dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may cause dizziness, double vision, slurred speech, drowsiness, lethargy, and mild diarrhea.

While you are using it

- Things you must do

- Take your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are taking ONEGABA 600.
- Tell your doctor immediately if you become pregnant while using this medication.
- Take ONEGABA 600 only with a doctor's prescription.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give ONEGABA 600 to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Driving and using machines

- This medicine may affect your ability to drive or use machines. If the tablets make you feel drowsy, dizzy or affect your vision, do not drive or use

ONEGABA 600

Gabapentin (600mg)

machines and contact your doctor immediately.

- Do not stop taking ONEGABA 600 suddenly, as stopping too quickly may cause seizures to return, including severe continuous seizures. If treatment needs to be stopped, your doctor will reduce the dose gradually over at least one week.
- Watch out for any breathing problems that occurred while taking this medicine, especially if you are an elderly or patient taking CNS depressants. If this happens, see your doctor as soon as possible.

Side effects

Like all medicines, ONEGABA 600 can cause side effects, although not everybody gets them.

Possible side effects include:

Commonly reported

- Dizziness
- Vertigo
- Drowsiness
- tingling or numbness (paraesthesia)
- changes in reflexes (increased, decreased or absent reflexes)
- increased body movements (hyperkinesia)
- anxiety
- hostility
- weakness (asthenia)
- feeling unwell (malaise)
- swelling of the face
- high blood pressure
- loss of appetite
- flatulence
- inflammation of the gums (gingivitis)
- joint pain
- pneumonia
- urinary tract infection
- bruising (purpura)
- visual disturbances

Children (3–12 years):

- viral infection
- fever
- nausea and/or vomiting
- drowsiness

Rare but serious

Seek medical attention immediately if you experience:

- severe breathing difficulties (respiratory depression).
- difficulty swallowing (dysphagia).

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website nptra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of ONEGABA 600

Storage

Do not store above 30°C.

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

What it looks like

ONEGABA 600

White, biconvex, elliptical, film-coated tablets, with deep break line on both sides and debossed with 'D' and '24' on either side of the break line on one side and plain on other side.

Ingredients

- Active ingredient

Gabapentin

- Inactive ingredients

Maize starch, Crospovidone, Copovidone, Isopropyl Alcohol, Cellulose, microcrystalline (Avicel PH102 & Ceolus KG1000), Magnesium stearate, Opadry White YS-1-18111 and Methylene Chloride.

MAL number(s):

MAL13025038AZ

Manufacturer

Aurobindo Pharma Limited, Unit-VII (SEZ),

Plot No. S1 Survey No. 411/P, 425/P, 434/P, 435/P & 458/P, (Part) Green Industrial Park, Polepally Village, Jedcherla Mandal, Mahaboobnagar District, Telangana State, India.

Product Registration Holder

UNIMED SDN BHD

No. 53, Jalan Tembaga SD 5/2B, Bandar Sri Damansara, 52200 Kuala Lumpur, Malaysia

Date of revision

26/08/2026

Serial Number

NPRA(R4/1)090626/00027