OLODIN

(Olopatadine HCL Opthalmic Solution USP 0.1% w/v)

What is in this leaflet

1. What OLODIN is used for
   How OLODIN works
2. Before you use OLODIN
3. How to use OLODIN
4. While you are using it
5. Side effects
6. Storage and Disposal of OLODIN
7. Product Description
8. Manufacturer and Product Registration Holder
9. Date of revision

What OLODIN is used for

OLODIN is used for the treatment of signs and symptoms of seasonal allergic conjunctivitis.
Allergic conjunctivitis. Some materials (Allergens) like pollens, house dust or animal fur may cause allergic reactions resulting in itching, redness as well as swelling of the surface of your eye.
OLODIN is a medicine for treatment of allergic conditions of the eye.

How OLODIN works

It works by reducing the intensity of the allergic reaction.

Before you use OLODIN

- *When you must not use it*

Pregnancy and lactation

Do not take OLODIN if you are pregnant, trying to get pregnant or think you may be pregnant.
Do not take OLODIN if you are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

- *Before you start use it*

Talk to your doctor or pharmacist before using OLODIN.
You should remove contact lenses that are in your eyes before using OLODIN.

Children
• Do not use OLODIN in children under the age of 3 years. Do not give this medicine to children under the ages of 3 years because there is no data to indicate that it is safe and work in children under 3 years.

- *Taking other medicines*
  Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.
  If you are using other eye drops or eye ointment medicines, leave at least 5 minutes between each medicine. Eye ointments should be administered last.

How to use OLODIN

- *How much to use*

  Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- *When to use it*

  Use as directed by your doctor or pharmacist.

- *How long to use it*

  Continue taking OLODIN for as long as your doctor recommends.

- *If you forget to use it*

  Consult your doctor or pharmacist on what you should do if you forget to use it.
  Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- *If you use too much (overdose)*

  Rinse it all out with warm water. Do not put in any more drops until it's time for your next regular dose.

While you are using it

- *Things you must do*

  Take your medicine exactly as your doctor has told you.
  Tell all the doctors, dentists and pharmacists treating you that you are taking OLODIN.
  Tell your doctor immediately if you become pregnant while taking this medication.

- *Things you must not do*

  Do not stop taking the medicine unless advised by your doctor.
  Do not take any new medicines without consulting your doctor.
  Do not give OLODIN to anyone else, even if they have the same symptoms or condition as you.

- *Things to be careful of*

  Driving and using machines

  This medicine may affect your ability to drive or use machines. If the OLODIN make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

Side effects

Like all medicines, OLODIN can cause side effects, although not everybody gets them.
Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.
Common: Headaches, lack of energy (asthenia), blurred vision, burning or stinging, cold syndrome, dry eye, foreign object were scratching the eye (foreign body sensation), an excess of blood in the vessels supplying an organ or other part of the body (hyperaemia), hypersensitivity, inflammation of the cornea of the eye. (Keratitis), lid edema, nausea, pharyngitis, pruritus, rhinitis, sinusitis and taste perversion

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug

Reaction Monitoring by visiting the website npra.moh.gov.my [Consumers→Reporting Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of OLODIN

- *Storage*

  Keep out of the reach and sight of children.
  Store at temperature not exceeding 30ºC.
  Protect from light

- *Disposal*

You should throw away the bottle four weeks after you first opened it to prevent infections, and use a new bottle. Write down the date you opened it in the space on each bottle label and box
Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- *What it looks like*
  Clear, colourless solution

- *Ingredients*
  - Active ingredient

    Olopatadine Hydrochloride

  - Inactive ingredients

    Benzalkonium Chloride, Sterile Aqueous.

- *MAL number:*
  MAL17025026AZ

Manufacturer
FDC limited
B-8 MIDC Industrial Area
Waluj, Aurangabad-431136
Maharashtra, India

Product Registration Holder
Unimed Sdn Bhd
53, Jalan Tembaga SD 5/2B, Bandar Sri Damansara 52200 Kuala Lumpur.

Serial number:
NPRA(R2/1)200210/354

Date of revision
18/09/2020