

MOSI (Moxifloxacin Ophthalmic Solution 0.5% w/v)

Moxifloxacin HCl (0.5 % w/v)

What is in this leaflet

1. What **MOSI** is used for
2. How **MOSI** works
3. Before you use **MOSI**
4. How to use **MOSI**
5. While you are using it
6. Side effects
7. Storage and Disposal of **MOSI**
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What **MOSI** is used for

MOSI is a sterile topical ophthalmic solution used to treat patients one year of age and older with bacterial conjunctivitis, which is an infection of the outer surface of the eye.

How **MOSI** works

MOSI contain Moxifloxacin which is often an antibacterial agent against a broad spectrum of Gram-positive and Gram-negative organisms.

Before you use **MOSI**

When you must not use it

- If you have ever an allergic (hypersensitivity) reaction to moxifloxacin or any other ingredient of this medicine.
- If you know you will drive, use machinery, or do any other activity requires clear vision.

Pregnancy and lactation

Do not take **MOSI** if you are pregnant, trying to get pregnant or think you may be pregnant.

Do not take if you are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

Before you start use it

Please tell your doctor or pharmacist if you are wearing contact lenses.

Please tell your doctor or pharmacist if you are applying any other type of eye drops or eye ointment before you start to use this medicine. Your medicine may affect their action and could alter their effect.

Please tell your doctor if you are taking any other medicines, including herbal products or any that you buy without a prescription

from a pharmacy, supermarket or health food shop.

How to use **MOSI**

How much to use

Instill one drop in the affected eye 3 times a day for 7 days.

If you are using other kinds of eye drops, wait at least five minutes before applying the other medications.

Always use **MOSI** exactly as your doctor has told you. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

Instructions for use:

(Please also refer to pictograms at the end of the leaflet)

- First wash your hands.
- Avoid touching the eye (or any other surface) with the dropper tip.
- Remove the soft contact lenses before using the eye drops.
- Shake the drops gently to be sure the medicine is well mixed.
- Tilt your head back, gaze upward and pull down the lower eyelid to make a pouch.
- Place the dropper directly over eye and gently squeeze the bottle to release a drop into your eye.
- Look downward and gently close your eye for 1 to 2 minutes. Press your finger against the inside corner of the closed eye, and hold for 1 minute.
- Try not to blink and do not rub the eye.
- Repeat the above steps for the other eye if necessary.
- Replace and tighten the cap immediately after use.

Be careful not to touch the dropper tip onto any surface, including the eyes and hands to avoid contamination. Do not use any eye drop that has particle in it.

When to use it

Use as directed by your doctor or pharmacist.

How long to use it

Continue taking **MOSI** for as long as your doctor recommends.

If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Apply the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

If you use too much (overdose)

If you used too much of **MOSI**, flush your eye(s) with tap water. Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking **MOSI**.

Tell your doctor immediately if you become pregnant while taking this medication.

Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor.

Do not give to anyone else, even if they have the same symptoms or condition as you.

Things to be careful of

Pregnancy and lactation

MOSI should be used during pregnancy only if it benefits both mother and foetus.

Caution should be exercise when **MOSI** is administered to nursing mother.

MOSI (Moxifloxacin Ophthalmic Solution 0.5% w/v)

Moxifloxacin HCl (0.5 % w/v)

Driving and using machines

This medicine may affect your ability to drive or use machines. The medication may make your vision temporarily blurred or unstable. Do not drive, use machines, or do any activity that required clear vision until you are sure you can perform the activities safely.

Side effects

Like all medicines, **MOSI** can cause side effects, although not everybody gets them. Tell your doctor immediately if you notice any of the following side effects:

- Burning/stinging
- Headache
- Inflammation of cornea
- Itchy eyes
- Redness of eyes
- Inflammation at the back of the throat
- Bleeding from eyes

Stop using this medication if you experience an allergic reaction include rash, swelling, dizziness, trouble breathing after applying this medication and visit your doctor or pharmacist immediately.

You may report any side effects or adverse drug reactions directly to The National Centre for Adverse Drug Reaction Monitoring by visiting the website nprra.moh.gov.my (Consumers → Reporting → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

Storage and Disposal of MOSI

- Storage

Keep out of the reach and sight of children.

Preserve in tight containers. Store below 30°C.

After first opening: preserve in tight container and store below 30°C.

Discard after 28 days of opening

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description



- What it looks like

Each bottle contains 5ml of

MOSI.

Your medication is a clear, yellow coloured solution.

Mode of Use:



MAL number:

MAL18026179AZ

Manufacturer

FDC Limited

Plot No.B-8, M.I.D.C Industrial Area,
Waluj- 431136, Dist: Aurangabad
INDIA.

Product Registration Holder

UNIMED SDN BHD

53, JalanTembaga SD 5/2B,
Bandar Sri Damansara,
52200, Kuala Lumpur, Malaysia.

Date of revision

17/03/2020

Serial Number

NPRA(R2/1)190429/121

Ingredients

Active ingredient

0.545% w/v Moxifloxacin
Hydrochloride equivalent to
Moxifloxacin 0.5%w/v

Inactive ingredients

Disodium Edetate, Sodium Chloride,
Boric Acid, Sodium Hydrochloride, and
Water for Injection.