

ANTIHIIST SYRUP

Chlorpheniramine maleate (4mg/5ml)

What is in this leaflet

1. What Antihist Syrup is used for
2. How Antihist Syrup works
3. Before you take Antihist Syrup
4. How to take Antihist Syrup
5. While you are using Antihist Syrup
6. Side effects
7. Storage and Disposal of Antihist Syrup
8. Product Description
9. Manufacturer and Product registration holder
10. Date of Revision

What Antihist Syrup is used for

Antihist Syrup is used to treat problems caused by allergies, such as hay fever, itchy skin, swelling, food allergy, reactions to medicines, or insect bites.

How Antihist Syrup works

Antihist Syrup contains the active ingredient chlorpheniramine maleate. Chlorpheniramine is an antihistamine, which blocks the action of histamine in the body and relieves swelling, itching, mucus production, and other symptoms of allergy.

Before you take Antihist Syrup

Do not give this medicine: to children below 2 years old. For children between 2 to 6 years old, this medicine should be used caution and following the advice given by your doctor or pharmacist.

- When you must not take it

Do not take this medicine if :

- you are allergic to chlorpheniramine or any other ingredients of this medicine (listed in Product Description)
- you are taking, or have taken in the past two weeks, medicines for depression called Monoamine oxidase inhibitors (MAOIs).

- Before you start to take it

Pregnancy and breast-feeding

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy, or breast-feeding before using any medicine.

Chlorpheniramine may be secreted into breast milk and may affect your baby. Your doctor will discuss the risks and benefits of breast feeding while you're taking Antihist Syrup.

- Taking other medicines

Before you take this medicine, make sure that you tell your doctor or pharmacist about any other medicines you might be using at the same time, particularly the following:

- medicines to help you relax (hypnotics or anxiolytics)
- Phenytoin (for epilepsy)

If you are unsure about interactions with any other medicines, talk to your pharmacist. This includes medicines prescribed by your doctor and medicine you have bought for your child, including herbal and homeopathic remedies.

How to take Antihist Syrup

- How much to use

Adults, and children over 12 years:

5 ml, to be taken three or four times a day

Children 6-12 years:

2.5 to 5 ml, to be taken three or four times a day

Children 2-5 years:

1.25 to 2.5 ml, to be taken three or four times a day

- When to use it

To be taken 3 or 4 times a day as directed by your doctor or pharmacist.

- How long to take it

Use Antihist Syrup for the duration that the doctor recommends. If the symptoms do not go away, talk to your doctor or pharmacist.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it. Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (Overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too much of this medicine may cause you to have seizures, difficulty breathing and low blood pressure.

While you are using Antihist Syrup

- Things you must do

Use your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are using Antihist Syrup

Tell your doctor immediately if you become pregnant while using this medication.

- Things you must not do

Do not stop using the medicine unless advised by your doctor.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Antihist Syrup to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

If you have an intolerance to some sugars do not take this medicine, unless if your doctor allowed you to do so. This medicine contains refined sugar.

Driving and using machines

This medicine may cause drowsiness and dizziness. If affected, do not drive or operate machines.

Do not take this medication together with alcohol or other sedating medication.

ANTI HIST SYRUP

Chlorpheniramine maleate (4mg/5ml)

Side effects

Like all medicines, Antihist Syrup can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

Possible side effects include:

- drowsiness
- dizziness

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npa.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Lot 5030, Jalan Teratai,
5 ½ Mile Off Jalan Meru,
41050 Klang,
Selangor Darul Ehsan, Malaysia

Product Registration Holder

Pahang Pharmacy Sdn Bhd
Lot 5979, Jalan Teratai,
5 ½ Mile Off Jalan Meru,
41050 Klang,
Selangor Darul Ehsan, Malaysia

Date of Revision

20/12/ 2023

Serial Number

NPRA(R4/1)181223/00092

Storage and Disposal of Antihist Syrup**-Storage**

Store below 30°C

Keep container tightly closed

Protect from light

-Disposal

Medicines should not be disposed of via wastewater or household waste.

Ask your pharmacist how to dispose of medicines no longer required.

These measures will help to protect the environment.

Product Description**- What it looks like**

A clear orange liquid.

-Ingredients**- Active ingredient**

Chlorpheniramine maleate

- Inactive ingredient

raspberry flavor, sunset yellow lake,
refined sugar carmoisine lake,
Methylparaben, Chloroform spirit.

- MAL Number.

MAL 19861155AZ

Manufacturer

Noripharma Sdn. Bhd