

# CORIFED SYRUP

Triprolidine Hydrochloride/Pseudoephedrine Hydrochloride (1.25mg/30mg per 5ml)

## What is in this leaflet

1. What CORIFED SYRUP is used for
2. How CORIFED SYRUP works
3. Before you use CORIFED SYRUP
4. How to use CORIFED SYRUP
5. While you are using it
6. Side effects
7. Storage and Disposal of CORIFED SYRUP
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What CORIFED SYRUP is used for

Corifed Syrup is used to treat symptoms of stuffy nose, sneezing, itching, watery eyes, and sinus congestion caused by allergies or the common cold.

## How CORIFED SYRUP works

Corifed Syrup contains two active ingredients which are triprolidine hydrochloride and pseudoephedrine hydrochloride.

Triprolidine is an antihistamine, which blocks the action of histamine in the body and relief sneezing, runny nose, watery eyes and other allergy symptoms. Pseudoephedrine is a decongestant that relieves nasal swelling and congestion.

## Before you use CORIFED SYRUP

### - When you must not use it

Do not use CORIFED SYRUP if:

- You are allergic or hypersensitive to triprolidine, pseudoephedrine, or any other ingredients of this medicine (listed in Product Description).
- You are being treated with or are recently taking (within 2 weeks) monoamine oxidase inhibitors (MAOIs) for depression.

### Before you start to use it

Tell your doctor if :

- you have prostate enlargement or bladder dysfunction
- severe liver and kidney problems

When used for treatment of cough and cold:

(a) Not to be used if your child is below 2 years old

(b) To be used with caution under doctor's or pharmacist's advice if your child is 2 to 6 years old

### Pregnancy and lactation

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine

### Taking other medicines

Tell your doctor if you are taking any other medicines, including any medicine that you buy without a prescription from a pharmacy, supermarket or health food shop.

Tell your doctor or pharmacist if you are taking :

- decongestants, appetite suppressants, amphetamines
- medications used for heart problems such as bretylium
- medications use for sedatives or tranquilizers
- blood pressure lowering medications such as bethanidine, guanethidine, debrisoquine, methyldopa, and beta blockers.
- other antihistamines

## How to use CORIFED SYRUP

### How much to use

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

Dosage:

- Adults and children over 12 years: 10ml (2 teaspoonfuls) every 4-6 hours
- Children 6-12 years: 5ml (1 teaspoonful) every 4-6 hours
- Children 2-5 years: 2.5ml (1/2 teaspoonful) every 4-6 hours
- Children under 2 years: Not to be used for cough and cold. For other indications, it can only be use as directed by your doctor

### When to use it

Take Corifed Syrup every 4-6 hours,

or as directed by your doctor.

### How long to use it

Take Corifed Syrup for the duration that the doctor recommends.

### If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too much of this medication could lead to problems like feeling very sleepy, weak, uncoordinated, difficulty urinating, slowed breathing, low blood pressure, agitation, irritability, seizures, high blood pressure, and fast or irregular heartbeats.

## While you are using it

### -Things you must do

Tell all the doctors, dentists and pharmacists treating you that you are taking CORIFED SYRUP.

Tell your doctor immediately if you become pregnant while taking this medication.

### -Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give CORIFED SYRUP to anyone else, even if they have the same symptoms or condition as you.

### -Things to be careful of

#### Driving and using machines

This medicine may affect your ability to drive or use machines. If the medication make you feel

# CORIFED SYRUP

Triprolidine Hydrochloride/Pseudoephedrine Hydrochloride (1.25mg/30mg per 5ml)

---

drowsy, or affect your hearing, do not drive or use machines and contact your doctor immediately.

Avoid taking together with alcohol as it may increase risk of sedative effects.

## Side effects

Like all medicines, CORIFED SYRUP can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

The side effects that was reported include feeling sleepy or difficulty sleeping.

Other side effects (rare) are urinary retention, especially in males with prostate enlargement, and skin reactions (red circular patches, and skin eruptions)

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my) [Consumers → Reporting Side Effects to Medicines ConSERF) or Vaccines (AEFI)].

## Storage and Disposal of CORIFED SYRUP

### Storage

Keep out of reach of children  
Store below 30°C.  
Keep container tightly closed.  
Protect from light.

### Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## Product Description

### What it looks like

A clear yellow liquid

### Ingredients

Active ingredient(s)

Each 5ml contains:

Triprolidine Hydrochloride 1.25mg

Pseudoephedrine Hydrochloride 30mg

### Inactive ingredients

Refined sugar, Methylparaben  
Chloroform spirit,  
Carboxymethylcellulose sodium,  
Quinoline yellow, Raspberry flavour,  
Purified water

### MAL number:

MAL19860260AZ

### Manufacturer

Noripharma Sdn. Bhd  
Lot 5030, Jalan Teratai, 5 ½ Mile Off  
Jalan Meru, 41050 Klang, Selangor  
Darul Ehsan, Malaysia.

### Product Registration Holder

Pahang Pharmacy Sdn. Bhd  
Lot 5979, Jalan Teratai, 5 ½ Mile Off  
Jalan Meru, 41050 Klang, Selangor  
Darul Ehsan, Malaysia.

### Date of revision

14/03/2024

### Serial No.:

NPRA(R4/1)160124/00008