

# CORIFED DM SYRUP

Tripolidine Hydrochloride/ Pseudoephedrine Hydrochloride / Dextromethorphan Hydrobromide  
(1.25mg/30mg/10mg per 5ml)

## What is in this leaflet

1. What is Corifed DM used for
2. How Corifed DM
3. Before you use Corifed DM
4. How to use Corifed DM
5. While you are using it
6. Side effects
7. Storage and Disposal of Corifed DM
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

## What is Corifed DM used for

Corifed DM Syrup is used to relieve dry cough with congestion in the upper respiratory tract, including congestion related to allergies.

## How Corifed DM works.

Corifed DM syrup contains tripolidine, pseudoephedrine and dextromethorphan as active ingredients.

Tripolidine an antihistamine which blocks the effects of histamine, and relieves sneezing, runny nose, watery eyes and other allergy symptoms..

Pseudoephedrine is a decongestant. relieves nasal swelling and congestion.

Dextromethorphan is a cough suppressant. , which reduces the cough reflex in the brain.

## Before you use Corifed DM

- When you must not take it  
Do not take this syrup if you :

- are allergic to pseudoephedrine, tripolidine, or dextromethorphan or any other ingredients of this medicine (listed in Product Description)
- have very high blood pressure or severe heart disease
- are taking or have taken in last two weeks, medicines for depression called Monoamine oxidase

inhibitors (MAOIs).

- had a cough with a lot of phlegm or that may be caused by asthma.

- Before you start to take it

Tell your doctor or pharmacist if you:

- suffer from liver or kidney problems
- have breathing problems
- have uncontrolled diabetes
- have increased pressure in the eye
- have overactive thyroid
- have prostate enlargement

When used for treatment of cough and cold:

- (a) Not to be used if your child is below 2 years old
- (b) To be used with caution under doctor's or pharmacist's advice if your child is 2 to 6 years old

## Pregnancy and lactation

If you are pregnant, planning to get pregnant or think you may be pregnant, discuss with your doctor before starting any medicine.

Corifed DM Syrup must only be used during pregnancy if the benefits is more than the risks to the fetus.

Small amount Corifed DM Syrup is secreted into breast milk and may affect your baby. Your doctor will discuss the risks and benefits of breast feeding while you're taking Corifed DM Syrup.

## -Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Tell your doctor or pharmacist if you are taking :

- medications to lower high blood pressure including bethanidine.
- Monoamine Oxidase Inhibitors

(MAOIs)for example furazolidone (to treat bacterial infections)

- Sympathomimetics medications such as decongestants, antidepressants, amphetamine-like stimulants
- bretylium (to treat irregular heart rhythm)

## How to take Corifed DM syrup

- How much to take

Adults and children over 12 years:  
10ml (2 teaspoonfuls), to be taken 3 times a day

Children 6-12 years:  
5ml (1 teaspoonful), to be taken 3 times a day

Children 2-5 years:  
2.5ml (1/2 teaspoonful), to be taken 3 times a day

- When to take it

Take 3 times a day or as directed by doctor or pharmacist

- How long to take it

Take Corifed DM for the duration that the doctor recommends.

- If you forget to take it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take extra medicine to make up the missed dose.

- If you take too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have use too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too much of this medicine may cause feeling sleepy, tired, dizzy, lack of coordination, abnormal eye

# CORIFED DM SYRUP

Tripolidine Hydrochloride/ Pseudoephedrine Hydrochloride / Dextromethorphan Hydrobromide  
(1.25mg/30mg/10mg per 5ml)

movements, weakness, poor muscle strength, slowed breathing, dry skin, fast heart rate, high blood pressure, hyperactive, very high body temperature, irritability, seeing or hearing things that aren't there, seizures, trouble urinating, and feelings of nausea or vomiting.

## While you are using Corifed DM syrup

### - Things you must do

Use your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are using Corifed DM.

Tell your doctor immediately if you become pregnant while using this medication.

### - Things you must not do

Do not stop using the medicine unless advised by your doctor.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Corifed DM to anyone else, even if they have the same symptoms or condition as you.

### - Things to be careful of Driving and using machines

This medicine may affect your ability to drive or use machines.

If the medicine make you drowsy or impair your thinking abilities, do not drive or use machines and contact your doctor immediately.

Do not take this medication together with alcohol or other sedating medication.

## Side effects

Like all medicines, Corifed DM can cause side effects, although not

everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

The side effects include:

- sleep disturbances
- hallucinations
- drowsiness
- fixed drug eruptions (skin reactions)
- urinary retention, mainly in males with prostate enlargement
- nausea, vomiting, or other gastrointestinal disturbances

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my) [Consumers —> Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

## Storage and Disposal of Corifed DMsyrup

### - Storage

Store in a cool dry place below 30°C. Keep the container tightly closed. Protect from light.

Keep medicine out of reach of children.

### - Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## Product description

### - What it looks like

Red coloured syrup having raspberry flavour.

### - Ingredients

- Active ingredient  
Pseudoephedrine Hydrochloride  
Tripolidine hydrochloride  
Dextromethorphan Hydrobromide

- Inactive ingredients

- Refined sugar
- Methylparaben
- Propylparaben
- Chloroform spirit
- Carbohymethylcellulose sodium
- Ponceau 4R
- Citric acid
- Raspberry flavour
- Purified water

### MAL Number.

MAL19992586AZ

### **Manufacturer:**

Noripharma Sdn. Bhd  
Lot 5030, Jalan Teratai, 5 ½ Mile Off  
Jalan Meru, 41050 Klang, Selangor  
Darul Ehsan, Malaysia.

### **Product Registration Holder**

Pahang Pharmacy sdn. Bhd  
Lot 5979, Jalan Teratai, 5 ½ Mile Off  
Jalan Meru, 41050 Klang, Selangor  
Darul Ehsan, Malaysia.

### **Date of revision**

11/03/2024

### **Serial Number**

NPRA(R4/1)160124/00009