

DOMIL SUSPENSION

Domperidone (5mg/5ml)

What is in this leaflet

1. What is Domil used for
2. How Domil works
3. Before you use Domil
4. How to use Domil
5. While you are using it
6. Side effects
7. Storage and Disposal of Domil
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What is Domil used for

For the relief of the symptoms of nausea and vomiting.

How Domil works

Domperidone works by blocking the action of a chemical messenger in the brain which causes the feeling of nausea and vomiting.

Before you use Domil

- When you must not use it
 - If you have ever had an allergic to domperidone or any of the ingredients listed at the end of this leaflet.
 - If you have a tumor of the pituitary gland (prolactinoma).
 - If you have ever had heart problems including heart failure, or heart rhythm disorders including a rapid or slow or irregular heartbeat.
 - If you are taking medications for heart problem
 - If you have an underlying illness affecting the digestive system, for example:
 - A blocked or perforated gut.
 - Your bowel motions (stools) are often black.
 - If you have moderate to severe liver problems.

- Before you start to use it

Domperidone should only be used during the first trimester of pregnancy when really necessary.

A small amount of domperidone is secreted in breast milk. Thus, do not take this medicine if you are breast-feeding unless your doctor or another health care professional has told you to.

Inform your doctor if you have kidney or liver problems.

- Taking other medicines

Inform your doctor if you are taking any medicines including

- an “anticholinergic” medicine
- an antifungal (e.g. ketoconazole)
- an antibiotic (e.g. erythromycin)
- an antiviral also known as protease inhibitors
- nefazodone, use for depression

How to use Domil

- How much to use

Adults and children ≥ 12 years of age and weighing ≥ 35 kg

10 mL 3 to 4 times per day

Children < 12 years of age and weighing ≥ 35 kg

Calculation by doctor based on body weight

Children < 12 years of age and weighing < 35 kg

The efficacy of Domil suspension in this group of people has not been established.

If you have severe kidney impairment, dosage and dosing frequency may need to be reduced to once or twice daily depending on the severity of the impairment.

- When to use it

Domil should be taken 3 or 4 times daily, 15-30 minutes before meals.

- How long to use it

Usually, the maximum treatment duration should not exceed one week. If nausea and vomiting persists for longer than one week, consult your doctor.

- If you forget to use it

If you forget to take a dose, take it as soon as you remember it. However, if it is almost time for your next dose, skip the missed dose. Do not take a double dose to make up for a forgotten dose

- If you use too much (overdosage)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have use too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

Go easy on snacks. Snack foods often contain a lot of fat, and can make your symptoms worse.

Eat regularly. Have three or four meals a day, at the same times, and don't rush your meals.

Shake the bottle before pouring.

- Things you must not do

Do not take Domil after meal as the absorption of the medicine may be delayed.

- Things to be careful of

Cut down or avoid from your 'triggers' such as coffee, fatty foods or alcohol.

Side effects

Side effects occur rarely and may include transient stomach cramps. Domperidone can also cause allergic reaction, or discharge from breasts, and swollen breasts (in men).

Side effects such as irregular heart beat or heart rhythm and increase heart beat are rarely reported.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Domil

- Storage

Store below 30°C. Protect from light. Keep container tightly closed. Keep out of reach of children

- Disposal

DOMIL SUSPENSION

Domperidone (5mg/5ml)

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

White coloured suspension with raspberry flavour.

- Ingredients

- Active ingredient

Domperidone

- Inactive ingredients

Carboxymethylcellulose sodium, Sorbitol 70% solution, Methylparaben, Sodium citrate, Colloidal silicone dioxide, Dimethylpolysiloxane, Raspberry flavor, Polysorbate 80, Propylparaben, Purified water, Sodium saccharin.

- MAL number:

MAL20034660AZ

Manufacturer

Noripharma Sdn. Bhd.
Lot 5030, Jalan Teratai.
5½ Mile Off Jalan Meru,
41050 Klang, Selangor, Malaysia

Product Registration Holder

Pahang Pharmacy Sdn. Bhd.
Lot 5979, Jalan Teratai.
5½ Mile Off Jalan Meru,
41050 Klang, Selangor, Malaysia.

Date of Revision

11/03/2024

Serial Number

NPRA(R4/1)070224/00018