

SULBUMIN 200 CAPSULE

Sulbutiamine (200mg)

What is in this leaflet

1. What Sulbumin is used for
2. How Sulbumin works
3. Before you use Sulbumin
4. How to use Sulbumin
5. While you are using it
6. Side effects
7. Storage and Disposal of Sulbumin
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Sulbumin is used for
Sulbumin is used for treatment of all form of function asthenia (lack of strength)

How Sulbumin works

Sulbutiamine is a thiamine derivative. Sulbutiamine is a lipophilic molecule that crosses the blood brain barrier more easily than vitamin B1 and leads to an increase in the level of Thiamine triphosphate. It gives better physical resistance, improved intellectual performance and psychological balance.

Before you use Sulbumin

- When you must not use it

Do not take Sulbumin if:

- You are allergic (hypersensitive) to Sulbutiamine or any of the other ingredients.

If you are pregnant, trying to get pregnant or think you may be pregnant, consult your doctor before taking this medicine.

If you are breastfeeding, consult your doctor before taking this medicine. This is because Thiamine is widely distributed to most body tissues and appears in breast milk.

- Before you start to use it

You must consult a doctor or pharmacist if you:

- Allergic (hypersensitivity) to Sulbutiamine or other ingredients in this product.

- Taking other medicines

Consult a doctor or pharmacist if you are taking:

- Loop diuretics medicines such as furosemide, ethacrynic acid and bumetanide.
- Intake of thiamine with foods that contain sulfites (such as dried fruits, potato chips, jams and grape juice).
- Intake of thiamin tea, coffee and decaffeinated coffee.

How to use Sulbumin

- How much to use

Follow all directions given in the insert carefully. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

For adult use only. 2-3 capsules daily. Capsule should be swallowed whole with a large glass of water, dividing the doses between the morning and midday meals. Duration of treatment is limited to 4 weeks.

- When to use it

Use as directed as in this leaflet.

- How long to use it

As per the dosage recommendation

- If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double

dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor or pharmacist for advice immediately.

Signs and symptoms of overdose include nervous excitement (agitation) with intense feelings of well-being (euphoria) and tremor in the extremities.

While you are using it

- Things you must do

Take your dose exactly as given in the leaflet.

Tell all the doctors, dentists and pharmacists that are treating you that you are taking Sulbumin.

Consult a doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not give Sulbumin to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of Driving and using machines

This product does not affect your ability to drive or use machines.

Side effects

Like all medicines, Sulbumin can cause side effects, although not everybody gets them.

The side effects are:

- Mild nervous excitement (agitation) in elderly person.
- Allergic skin reaction
- Unpleasant smell of urine.

SULBUMIN 200 CAPSULE

Sulbutiamine (200mg)

Visit your doctor or pharmacist immediately if you experience any side effects after taking this Sulbumin.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Sulbumin

- Storage

Store in a dry place below 30°C. Protect from light and moisture.

- Disposal

Products should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of Sulbumin that is no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Two pieces, size '0' maroon color hard vegetable capsule contain white color powder.

- Ingredients

-Active ingredient(s)
Sulbutiamine

- Inactive ingredients
Sodium Starch Glycolate
Corn Starch
Colloidal Silicon Dioxide
Microcrystalline Cellulose
Magnesium Stearate

- MAL number(s):

MAL06111366X

Manufacturer

Pahang Pharmacy Sdn. Bhd.
Lot 5979, Jalan Teratai, 5 1/2
Miles Off, Jalan Meru, 41050
Klang, Selangor, Malaysia.

Product Registration Holder

Pahang Pharmacy Sdn. Bhd.
Lot 5979, Jalan Teratai, 5 1/2
Miles Off, Jalan Meru, 41050
Klang, Selangor, Malaysia.

Date of Revision

October 2021

Serial Number

BPFK(R4/1)190417/00085