

SITAKRAFT SITAGLIPTIN FILM-COATED TABLETS

Sitagliptin Phosphate (50mg and 100mg)

What is in this leaflet

1. What **SITAKRAFT** is taken for
2. How **SITAKRAFT** works
3. Before you take **SITAKRAFT**
4. How to take **SITAKRAFT**
5. While you are taking **SITAKRAFT**
6. Side effects
7. Storage and disposal of **SITAKRAFT**
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial Number

Please read this information carefully before you start to take your medicine, even if you have just refilled your prescription. Some of the information may have changed.

Remember that your doctor has prescribed this medicine only for you. Never give it to anyone else.

What SITAKRAFT is taken for

SITAKRAFT (sitagliptin phosphate) is a tablet that contains 50, or 100 mg of sitagliptin as the active ingredient.

- **SITAKRAFT** helps to improve the levels of insulin after a meal.
- **SITAKRAFT** decreases the amount of sugar made by the body.
- **SITAKRAFT** works when blood sugar is high, especially after a meal. This is when the body needs the greatest amount of help in lowering blood sugar. **SITAKRAFT** also lowers blood sugar between meals.
- **SITAKRAFT** by itself is unlikely to cause low blood sugar (hypoglycemia) because it does not work when your blood sugar is low.

How SITAKRAFT works

SITAKRAFT is a member of a class of medicines you take by mouth called DPP-4 inhibitors (dipeptidyl peptidase-4 inhibitors) that lowers blood sugar levels in patients with type 2 diabetes mellitus. Type 2 diabetes is also called non-insulin-dependent diabetes mellitus, or NIDDM.

Why has my doctor prescribed SITAKRAFT ?

Your doctor has prescribed **SITAKRAFT** to help lower your blood sugar, which is too high because of your type 2 diabetes. **SITAKRAFT** can be used alone or in combination with certain other medicines that lower blood sugar, along with a recommended diet and exercise program.

What is type 2 diabetes?

Type 2 diabetes is a condition in which your body does not make enough insulin, and the insulin that your body produces does not work as well as it should. Your body can also make too much sugar. When this happens, sugar (glucose) builds up in the blood. This can lead to serious medical problems.

The main goal of treating diabetes is to lower your blood sugar to a normal level. Lowering and controlling blood sugar may help prevent or delay complications of diabetes, such as heart disease, kidney disease, blindness, and amputation.

High blood sugar can be lowered by diet and exercise, and by certain medicines.

Before you take SITAKRAFT

- When you must not take it
Do not take **SITAKRAFT** if you are allergic to any of the ingredients in **SITAKRAFT**.

- Before you start to take it
Tell your doctor if you have or have had:

- type 1 diabetes
- diabetic ketoacidosis (increased ketones in the blood or urine)
- any kidney problems, or any past or present medical problems
- an allergic reaction to **SITAKRAFT**
- gallstones, alcohol dependence or very high levels of triglycerides (a form of fat) in your blood. These medical conditions can increase your chance of getting pancreatitis.

- Taking other medicines
SITAKRAFT may be taken with most medicines. Tell your doctor about all the medicines you take. This includes prescription and non-prescription medicines, and herbal supplements.

In particular, tell your doctor:

- if you are taking digoxin (a medicine used to treat irregular heart beat and other heart problems). The level of digoxin in your blood may need to be checked if you are taking it with **SITAKRAFT**.

How to take SITAKRAFT

- How much to take and When to take it

Take **SITAKRAFT** exactly as your doctor has prescribed. The recommended dose is to take:

SITAKRAFT SITAGLIPTIN FILM-COATED TABLETS

Sitagliptin Phosphate (50mg and 100mg)

- one 100 mg tablet
- once a day
- by mouth, with or without food

If you have kidney problems, your doctor may prescribe lower doses.

Your doctor may prescribe **SITAKRAFT** along with certain other medicines that lower blood sugar.

- How long to take it

Continue to take **SITAKRAFT** as long as your doctor prescribes it so you can continue to help control your blood sugar.

Diet and exercise can help your body use its blood sugar better. It is important to stay on your doctor recommended diet, exercise and weight loss program while taking **SITAKRAFT**.

- If you forget to take it

If you miss a dose, take it as soon as you remember. If you do not remember until it is time for your next dose, skip the missed dose and go back to your regular schedule. Do not take a double dose of **SITAKRAFT**.

- If you take too much (overdose)

If you take more than the prescribed dosage of **SITAKRAFT**, contact your doctor immediately.

While you are taking SITAKRAFT

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking **SITAKRAFT**.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor. Do not give **SITAKRAFT** to anyone else, even if they have the

same symptoms or condition as you.

- Things to be careful of

Cases of inflammation of the pancreas (pancreatitis) have been reported in patients receiving **SITAKRAFT**. Pancreatitis can be a serious, potentially life-threatening medical condition. Stop taking **SITAKRAFT** and call your doctor if you experience severe and persistent stomach pain, with or without vomiting, because you could have pancreatitis.

Cases of a skin reaction called bullous pemphigoid that can require treatment in a hospital have been reported in patients receiving **SITAKRAFT**. Tell your doctor if you develop blisters or the breakdown of your skin (erosion). Your doctor may tell you to stop taking **SITAKRAFT**.

Use in children

SITAKRAFT is not effective in children and adolescents 10 to 17 years of age with type 2 diabetes. **SITAKRAFT** has not been studied in children younger than 10 years of age.

Use in the elderly

In studies, **SITAKRAFT** worked well in and was well-tolerated by older adult patients. No dosage adjustment is necessary based on age.

Use in pregnancy and breast-feeding

Women who are pregnant or plan to become pregnant should consult with their doctor before taking **SITAKRAFT**.

SITAKRAFT is not recommended for use during pregnancy.

It is not known if **SITAKRAFT** passes into breast milk. You should not use **SITAKRAFT** if you are breast-feeding or plan to breast-feed.

Can I drive or operate machinery while using SITAKRAFT?

There is no information to suggest that **SITAKRAFT** affects your ability to drive a car or operate machinery.

Side effects

Like all prescription medicines, **SITAKRAFT** may cause side effects. In studies, side effects usually were mild and did not cause patients to stop taking **SITAKRAFT**. The side effects reported in patients treated with **SITAKRAFT** were similar to side effects in patients treated with a tablet containing no medication (a placebo).

When **SITAKRAFT** is used in combination with a sulfonylurea medicine or with insulin, low blood sugar with symptoms (hypoglycemia) due to the sulfonylurea or insulin can occur. Lower doses of the sulfonylurea medicine or insulin may be required.

When **SITAKRAFT** was used in combination with insulin, the following additional side effects were reported:

- Flu
- Headache

When **SITAKRAFT** and metformin were started together, the following side effects were reported:

- Diarrhea
- Indigestion

SITAKRAFT SITAGLIPTIN FILM-COATED TABLETS

Sitagliptin Phosphate (50mg and 100mg)

- Flatulence
- Vomiting
- Headache

When **SITAKRAFT** and pioglitazone were started together, the side effect of decreased blood sugar measurements without symptoms of hypoglycemia was reported.

When **SITAKRAFT** was used in combination with metformin and rosiglitazone, the following side effects were reported:

- Headache
- Low blood sugar with symptoms (hypoglycemia)
- Diarrhea
- Upper respiratory infection
- Nausea
- Cough
- Fungal skin infection
- Swelling of the hands or legs
- Vomiting

Additional side effects have been reported in general use with **SITAKRAFT**, by itself and/or with other diabetes medicines:

- Allergic reactions, which may be serious, including rash, hives, and swelling of the face, lips, tongue, and throat that may cause difficulty in breathing or swallowing. If you have an allergic reaction, stop taking **SITAKRAFT** and call your doctor right away. Your doctor may prescribe a medication to treat your allergic reaction and a different medication for your diabetes.
- Inflammation of the pancreas.
- Kidney problems (sometimes requiring dialysis).
- Upper respiratory infection.
- Stuffy or runny nose and sore throat.
- Constipation.
- Vomiting.
- Headache.
- Joint pain.
- Muscle aches.

- Arm or leg pain.
- Back pain.
- Itching.
- Blisters.
- Reduced number of platelets.

Tell your doctor or pharmacist if you develop any unusual side effect, or if any known side effect does not go away or gets worse.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

How can I learn more about SITAKRAFT and diabetes?

You may obtain further information from your doctor or pharmacist, who has more detailed information.

Storage and disposal of SITAKRAFT

- Storage

Do not store above 30°C. Protect from light and Moisture.

- Disposal

How long should I keep my medicine?

Do not use this medicine after the month and year shown by the four numbers following EX (or EXP) on the container. The first two numbers indicate the month; the last two numbers indicate the year.

Product description

- What it looks like

SITAKRAFT 50mg: Light yellow round, biconvex film coated tablets, debossed with "L" one side and "5" on other side.

. Available in packs of 30s.

SITAKRAFT 100mg: Light brown round, biconvex film coated tablets, debossed with "L" one side and "7" on other side. Available in packs of 30s.

- Ingredients

Active ingredient: Sitagliptin phosphate

Inactive ingredient:

Core: Dicalcium Phosphate Anhydrous (A-Tab), Talc (Luzanac M), Croscarmellose sodium (Ac di sol), Sodium Stearyl Fumarate (Pruv), Colloidal silicon dioxide (Aerosil 200)

Tablets Coating : Opadry Yellow 03F520488, Opadry Brown 03F565259 Isoprophyl Alcohol, Methylene Chloride.

- MAL numbers:

MAL *****

Manufacturer

MANKIND PHARMA LTD.
Unit-III, Village Kunja, Rampur Ghat Road, Opposite Dental College, Tehsil Paonta Sahib, District Sirmaur 173025 (INDIA)

Registered Office :

208, Okhla Industrial Estate, Phase-3, New Delhi-110 020 (INDIA)

Product Registration Holder

Pahang Pharmacy Sdn. Bhd.
Lot 5979, Jalan Teratai, 5 1/2 Mile Off Jalan Meru, 41050 Klang, Selangor, Malaysia

Date of revision

09/03/2026

Serial Number:

NPRA (R2) 26/021