

NUFEN SUSPENSION

Ibuprofen (100mg in every 5ml)

What is in this leaflet

1. What Nufen Suspension is used for
2. How Nufen Suspension works
3. Before you use Nufen Suspension
4. How to use Nufen Suspension
5. While you are using it
6. Side Effects
7. Storage and Disposal of Nufen Suspension
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What Nufen Suspension is used for

Nufen Suspension is used to relief fever associated with cold as well as pain from teething, toothache, earache, sore throats, headache, and sprains.

Also for the relief of inflammation of muscles and joints.

How Nufen Suspension works

Nufen Suspension contains ibuprofen, a type of Non steroidal Anti-inflammatory Drugs (NSAIDs). It works by reducing prostaglandins which is natural substance that causes pain and inflammation.

Before you use Nufen Suspension

- When you must not use it

Do not take Nufen Suspension if you are:

- allergic to ibuprofen, aspirin, other NSAIDs or any other ingredients of this medicine (listed in Product Description)
- you have or had history of stomach or intestinal ulcer

- Before you start to use it

Tell your doctor if you have or history of:

- asthma or allergies
- any gastrointestinal problem
- kidney, liver or heart problem
- high blood pressure

- developing gastrointestinal bleeding (heavy or frequent alcohol drinking, smoking or taking medications or conditions that may increase chance of bleeding)

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Tell your doctor if you are taking:

- Lithium (mood disorders medication)
- Methotrexate (cancer and autoimmune medication)
- blood thinning medication
- high blood pressure medications
- diuretics (medication to get rid of extra fluid in the body)

How to use Nufen Suspension

- How much to use

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

Children age 1 to 12 years old with body weight more than 7kg:

The usual dose for children are as follows:

- 1 - 2 years: 2.5 ml
- 3 - 7 years: 5 ml (one teaspoon)
- 8 - 12 years: 10ml (two teaspoon)

Do not exceed the stated dose

- When to use it

Take 3 or 4 times a day or as directed by your doctor or pharmacist.

- How long to use it

Take Nufen Suspension for the duration your doctor

recommends. If the symptom still persists or worsen, please consult your doctor.

- If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have use too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too much of this medicine may cause headache, nausea, vomiting, drowsiness, low blood pressure and loss of consciousness.

While you are using it

- Things you must do

Tell all the doctors, dentists and pharmacists treating you that you are using Nufen Suspension

- Things you must not do

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Nufen Suspension to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Driving and using machines

This medicine may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

Side Effects

Like all medicines, Nufen Suspension can cause side effects, although not everybody gets them.

NUFEN SUSPENSION

Ibuprofen (100mg in every 5ml)

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

Ibuprofen may cause hypersensitivity with symptoms such as:

- severe allergic reactions
- difficulty breathing, especially if you have asthma
- skin reactions such as rashes, itching, swelling, or even blisters
- stomach pain, nausea, or indigestion
- bleeding in the stomach or low platelet count

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Nufen Suspension

- Storage

Keep all medicines out of the reach of children.

Store below 30°C.

Do not take this medicine after the expiry date stamped on the pack/bottle.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

An orange colored suspension with fruity flavour.

-Ingredients

- Active Ingredient

Ibuprofen

- Inactive ingredients

Colour sunset yellow, Water Purified, Sodium Citrate, citric acid monohydrate, Sodium Benzoate, Xanthan gum, Polysorbate 20, Propylparaben, orange flavour, Glycerin, refined sugar, Methylparaben and Gum tragacanth

- MAL Number

MAL07010961AZ

Manufacturer

Noripharma Sdn. Bhd.,
Lot 5030, Jalan Teratai,
5 1/2 Mile Off Jalan Meru,
41050 Klang,
Selangor Darul Ehsan, Malaysia

Product Registration Holder

Steriline Sdn. Bhd.,
Lot 5979, Jalan Teratai,
5 1/2 Mile Off Jalan Meru,
41050 Klang,
Selangor Darul Ehsan, Malaysia

Date of revision

14/08/2024

Serial Number

NPRA(R4/1)310724/00097