

ELOMET[®] CREAM

Mometasone Furoate (0.1% w/w)

What is in this leaflet

1. What *ELOMET CREAM* is used for
2. How *ELOMET CREAM* works
3. Before you take *ELOMET CREAM*
4. How to take *ELOMET CREAM*
5. While you are taking *ELOMET CREAM*
6. Side effects
7. Storage and Disposal of *ELOMET CREAM*
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What *ELOMET CREAM* is used for

ELOMET CREAM is used to reduce inflammation and itchiness caused by certain skin problems that are responsive to corticosteroids.

How *ELOMET CREAM* works

ELOMET CREAM is one of a group of medicines called topical corticosteroids.

Mometasone furoate have anti-inflammatory, anti itching and blood vessel narrowing action.

Before you use *ELOMET CREAM*

- When you must not use it

If you are allergic (hypersensitive) to mometasone furoate, or to any of the other ingredients of *ELOMET CREAM*

Side effects that may happen with oral or injectable corticosteroids may also occur with topical corticosteroids, especially in infants and children.

Ask your doctor or pharmacist if you are not sure.

- Before you start to use it

Pregnancy and breast-feeding

You should tell your doctor if you are pregnant or breast-feeding, before you start using *ELOMET CREAM*.

- Taking other medicines

Tell your doctor or pharmacist if you are using other creams, ointments or lotions or taking any other medicines. This includes any that you get without a prescription from your pharmacy, supermarket or health food shop.

How to use *ELOMET CREAM*

- How much to use

Always use *ELOMET CREAM* exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

A thin layer of *ELOMET CREAM* should be applied to the affected area of skin.

- When to use it

ELOMET CREAM should be applied once a day.

- How long to use it

Use *ELOMET CREAM* on the affected area of skin until the lesion heals or for a duration of three weeks whichever is sooner.

- If you forget to use it

If you forget to use your cream at the right time, use it as soon as you remember, then carry on as before.

- If you use too much (overdose)

You should tell your doctor if you use *ELOMET CREAM* more often than you should, or on large areas of the body, or for a long time, the chance of absorption through the skin and the chance of side effects increases and can affect some of your hormones or sugar in blood or urine. Infants and children are more susceptible to these effects due to their larger skin surface to body mass ratios.

While you are using *ELOMET CREAM*

- Things you must do

Tell all doctors and pharmacists who are treating you that you are using *ELOMET CREAM*.

If your skin becomes irritated or sensitive after using *ELOMET CREAM*, you should stop using it and tell your doctor

Contact your doctor if you experience blurred vision or other visual disturbances.

Tell your doctor if you feel that *ELOMET CREAM* is not helping your condition or if your skin condition worsens or appear to be infected.

Tell your doctor if, for any reason, you have not used *ELOMET CREAM* exactly as prescribed.

Otherwise, your doctor may think that it was not effective and change your treatment unnecessarily.

- Things you must not do

Do not use *ELOMET CREAM* under tight dressings or on large areas of skin unless your doctor tells you.

Do not use plastic pants or tight fitting nappies if *ELOMET CREAM* is to be used on the nappy area of young children.
Do not use *ELOMET CREAM* in or near the eyes.

Do not give *ELOMET CREAM* to anyone else even if their symptoms seem similar to yours.

Do not use *ELOMET CREAM* to treat other conditions unless your doctor tells you.

Your doctor has prescribed *ELOMET CREAM* especially for you and your condition. If you use it for another condition, it may not work or make the condition worse.

ELOMET[®] CREAM

Mometasone Furoate (0.1% w/w)

- Things to be careful of

Only use *ELOMET CREAM* on skin occlusive areas that rub together such as under the arm or in the groin area if your doctor tells you.

Only use *ELOMET CREAM* on the face if your doctor tells you.

If improvement does not occur within one week, tell your doctor.

Side effects

Like all medicines, *ELOMET CREAM* can cause side effects, although not everybody gets them.

A few people may find that they suffer from some of the following side effects after using *ELOMET CREAM*:

- allergic skin reactions
- secondary infections
- thinning of the skin
- red marks with associated prickly heat
- loss of skin colour
- burning
- stinging
- itching
- tingling
- excessive hair growth
- softening of the skin and stretch marks
- blurred vision

Other side effects that may occur with topical corticosteroids are dry skin, skin irritation, allergic contact skin inflammation, and rashes around the mouth.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website nptra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI)

Storage and disposal of *ELOMET CREAM*

- Storage

Store below 30°C.

Keep out of the reach of children.

Do not use *ELOMET CREAM* after the expiry date which is stamped on the pack.

- Disposal

Medicines should not be disposed of via wastewater or household waste.

Ask your pharmacist how to dispose of medicines no longer required.

These measures will help to protect the environment.

Product description

- What it looks like

ELOMET CREAM: A smooth, white to off-white homogenous cream, free from foreign matter. The cream is available in a tube of 15g.

- Ingredients

- Active ingredient: Each gram of *ELOMET CREAM* contains 1 mgmometasone furoate

- Inactive ingredients: White petrolatum, white wax, hydrogenated phosphatidylcholine, hexylene glycol, titanium dioxide, aluminium starch octenylsuccinate, purified water and phosphoric acid to adjust the pH.

- MAL number:

MAL19992562AZ

Manufacturer

Organon Heist bv
Industriepark 30,
2220 Heist-
op-den-Berg,
Belgium

Product Registration Holder

Organon Malaysia Sdn. Bhd.
3A-08-02, Level 8,
Corporate Tower 3A,
Pavilion Damansara Heights,
No. 3, Jalan Damansara,
50490 Kuala Lumpur,
Malaysia.

Date of revision

13/11/2025 (Version 082017)

Serial Number:

NPRA (R1/3) 28082023 156