

ATOZET® (ezetimibe/atorvastatin)

LPC-OG0653C-T-062023 [Tablet]

1. NAME OF THE DRUG

ATOZET® 10MG/10MG FILM-COATED TABLETS

(EZETIMIBE 10 MG/ATORVASTATIN 10 MG)

ATOZET® 10MG/20MG FILM-COATED TABLETS

(EZETIMIBE 10 MG/ATORVASTATIN 20 MG)

ATOZET® 10MG/40MG FILM-COATED TABLETS

(EZETIMIBE 10 MG/ATORVASTATIN 40 MG)

ATOZET® 10MG/80MG FILM-COATED TABLETS

(EZETIMIBE 10 MG/ATORVASTATIN 80 MG)

2. PHARMACEUTICAL FORM

Film-coated tablets

3. PHARMACEUTICAL PARTICULARS


3.1 Chemistry

ATOZET contains ezetimibe, a selective inhibitor of intestinal cholesterol and related phytosterol absorption, and atorvastatin, a 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase inhibitor.

Ezetimibe

The chemical name of ezetimibe is 1-(4-fluorophenyl)-3(R)-[3-(4-fluorophenyl)-3(S)-hydroxypropyl]-4(S)-(4-hydroxyphenyl)-2-azetidinone. The empirical formula is C₂₄H₂₁F₂NO₃. Its molecular weight is 409.4.


Ezetimibe is a white, crystalline powder that is freely to very soluble in ethanol, methanol, and acetone and practically insoluble in water. Its structural formula is:



Atorvastatin

Atorvastatin is [R-(R*, R*)]-2-(4-fluorophenyl)-β, δ , -dihydroxy-5-(1-methylethyl)-3-phenyl-4-[(phenylamino) carbonyl]-1H-pyrrole-1-heptanoic acid, calcium salt (2:1) trihydrate.

Atorvastatin calcium is a white or almost white powder that is soluble in dimethyl sulfoxide. The degree of solubility in water, ethanol, and methylene chloride is very slightly soluble to practically insoluble. The molecular formula of atorvastatin calcium is $C_{66}H_{68}CaF_2N_4O_{10} \cdot 3H_2O$. The molecular weight of atorvastatin calcium is 1209.36. Its structural formula is:



3.2 Composition

Active Ingredients

ATOZET is available for oral use as tablets containing 10 mg of ezetimibe and: 10.9 mg of atorvastatin calcium (crystalline trihydrate), equivalent to 10 mg of atorvastatin (ATOZET 10 mg/10 mg); 21.7 mg of atorvastatin calcium (crystalline trihydrate), equivalent to 20 mg of atorvastatin (ATOZET 10 mg/20 mg); 43.4 mg of atorvastatin calcium (crystalline trihydrate), equivalent to 40 mg of atorvastatin (ATOZET 10 mg/40 mg); or 86.8 mg of atorvastatin calcium (crystalline trihydrate), equivalent to 80 mg of atorvastatin (ATOZET 10 mg/80 mg).

Inactive Ingredients (List of excipients)

Each film-coated tablet of ATOZET contains the following inactive ingredients: calcium carbonate, colloidal silicon dioxide, croscarmellose sodium, hydroxypropyl cellulose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polysorbate, povidone, and sodium lauryl sulfate.

The film coating contains: hydroxypropyl methylcellulose/hypromellose, macrogol/polyethylene glycol, titanium dioxide, and talc.

3.3 Product Description

ATOZET are bilayer, capsule shaped, biconvex, film coated tablets, white to off white in color and with the following size and debossing.

10 mg/ 10 mg: with "257" debossed on one side of the tablet

10 mg/ 20 mg: with "333" debossed on one side of the tablet

10 mg/ 40 mg: with "337" debossed on one side of the tablet

10 mg/ 80 mg: with "357" debossed on one side of the tablet

4. CLINICAL PHARMACOLOGY

4.1 Therapeutic Class

ATOZET (ezetimibe/atorvastatin) is a lipid-lowering product that selectively inhibits the intestinal absorption of cholesterol and related plant sterols and inhibits the endogenous synthesis of cholesterol.

4.2 Mechanism of Action

ATOZET

Plasma cholesterol is derived from intestinal absorption and endogenous synthesis. ATOZET contains ezetimibe and atorvastatin, two lipid-lowering compounds with complementary mechanisms of action. ATOZET reduces elevated total-C, LDL-C, Apo B, TG, and non-HDL-C, and increases HDL-C through dual inhibition of cholesterol absorption and synthesis.

Ezetimibe

Ezetimibe inhibits the intestinal absorption of cholesterol. Ezetimibe is orally active and has a mechanism of action that differs from other classes of cholesterol-reducing compounds (e.g., statins, bile acid sequestrants [resins], fibric acid derivatives, and plant stanols). The molecular target of ezetimibe is the sterol transporter, Niemann-Pick C1-Like 1 (NPC1L1), which is responsible for the intestinal uptake of cholesterol and phytosterols.

Ezetimibe localizes at the brush border of the small intestine and inhibits the absorption of cholesterol, leading to a decrease in the delivery of intestinal cholesterol to the liver; statins reduce cholesterol

synthesis in the liver and together these distinct mechanisms provide complementary cholesterol reduction.

In a 2-week clinical study in 18 hypercholesterolemic patients, ezetimibe inhibited intestinal cholesterol absorption by 54%, compared with placebo.

A series of preclinical studies was performed to determine the selectivity of ezetimibe for inhibiting cholesterol absorption. Ezetimibe inhibited the absorption of [^{14}C]-cholesterol with no effect on the absorption of triglycerides, fatty acids, bile acids, progesterone, ethinyl estradiol, or the fat-soluble vitamins A and D.

Atorvastatin

Atorvastatin is a selective, competitive inhibitor of HMG-CoA reductase, the rate-limiting enzyme that converts 3-hydroxy-3-methyl-glutaryl-coenzyme A to mevalonate, a precursor of sterols, including cholesterol.

In animal models, atorvastatin lowers plasma cholesterol and lipoprotein levels by inhibiting HMG-CoA reductase and cholesterol synthesis in the liver and by increasing the number of hepatic LDL receptors on the cell surface to enhance uptake and catabolism of LDL; atorvastatin also reduces LDL production and the number of LDL particles.

4.3 Pharmacokinetics

General Introduction

ATOZET

ATOZET has been shown to be bioequivalent to coadministration of corresponding doses of ezetimibe and atorvastatin tablets.

Absorption

ATOZET

The effects of a high-fat meal on the pharmacokinetics of ezetimibe and atorvastatin when administered as ATOZET tablets are comparable to those reported for the individual tablets.

Ezetimibe

After oral administration, ezetimibe is rapidly absorbed and extensively conjugated to a pharmacologically active phenolic glucuronide (ezetimibe-glucuronide). Mean maximum plasma concentrations (C_{max}) occur within 1 to 2 hours for ezetimibe-glucuronide and 4 to 12 hours for ezetimibe. The absolute bioavailability of ezetimibe cannot be determined as the compound is virtually insoluble in aqueous media suitable for injection.

Concomitant food administration (high fat or non-fat meals) had no effect on the oral bioavailability of ezetimibe when administered as ezetimibe 10-mg tablets.

Atorvastatin

Atorvastatin is rapidly absorbed after oral administration; maximum plasma concentrations (C_{max}) occur within 1 to 2 hours. Extent of absorption increases in proportion to atorvastatin dose. After oral administration, atorvastatin film-coated tablets are 95% to 99% bioavailable compared to the oral solution. The absolute bioavailability of atorvastatin is approximately 12% and the systemic availability of HMG-CoA reductase inhibitory activity is approximately 30%. The low systemic availability is attributed to presystemic clearance in gastrointestinal mucosa and/or hepatic first-pass metabolism.

Distribution

Ezetimibe

Ezetimibe and ezetimibe-glucuronide are bound 99.7% and 88 to 92% to human plasma proteins, respectively.

Atorvastatin

Mean volume of distribution of atorvastatin is approximately 381 l. Atorvastatin is $\geq 98\%$ bound to plasma proteins.

Metabolism

Ezetimibe

Ezetimibe is metabolized primarily in the small intestine and liver via glucuronide conjugation (a phase II reaction) with subsequent biliary excretion. Minimal oxidative metabolism (a phase I reaction) has been observed in all species evaluated. Ezetimibe and ezetimibe-glucuronide are the major drug-derived compounds detected in plasma, constituting approximately 10 to 20 % and 80 to 90 % of the total drug in plasma, respectively. Both ezetimibe and ezetimibe-glucuronide are slowly eliminated from plasma with evidence of significant enterohepatic recycling. The half-life for ezetimibe and ezetimibe-glucuronide is approximately 22 hours.

Atorvastatin

Atorvastatin is metabolized by cytochrome P450 3A4 to ortho- and parahydroxylated derivatives and various beta-oxidation products. Apart from other pathways these products are further metabolized via glucuronidation. *In vitro*, inhibition of HMG-CoA reductase by ortho- and parahydroxylated metabolites is equivalent to that of atorvastatin. Approximately 70% of circulating inhibitory activity for HMG-CoA reductase is attributed to active metabolites.

Elimination

Ezetimibe

Following oral administration of ^{14}C -ezetimibe (20 mg) to human subjects, total ezetimibe accounted for approximately 93% of the total radioactivity in plasma. Approximately 78% and 11% of the administered radioactivity were recovered in the feces and urine, respectively, over a 10-day collection period. After 48 hours, there were no detectable levels of radioactivity in the plasma.

Atorvastatin

Atorvastatin is eliminated primarily in bile following hepatic and/or extrahepatic metabolism. However, the medicinal product does not appear to undergo significant enterohepatic recirculation. Mean plasma elimination half-life of atorvastatin in humans is approximately 14 hours. The half-life of inhibitory activity for HMG-CoA reductase is approximately 20 to 30 hours due to the contribution of active metabolites.

Special Populations

Renal Impairment

Ezetimibe

After a single 10-mg dose of ezetimibe in patients with severe renal disease ($n=8$; mean $\text{CrCl} \leq 30 \text{ ml/min/1.73 m}^2$), the mean AUC for total ezetimibe was increased approximately 1.5-fold, compared to healthy subjects ($n=9$).

An additional patient in this study (post-renal transplant and receiving multiple medications, including cyclosporine) had a 12-fold greater exposure to total ezetimibe.

Atorvastatin

Renal disease has no influence on the plasma concentrations or lipid effects of atorvastatin and its active metabolites.

Hepatic Impairment

Ezetimibe

After a single 10-mg dose of ezetimibe, the mean area under the curve (AUC) for total ezetimibe was increased approximately 1.7-fold in patients with mild hepatic impairment (Child-Pugh score 5 or 6), compared to healthy subjects. In a 14-day, multiple-dose study (10 mg daily) in patients with moderate hepatic impairment (Child-Pugh score 7 to 9), the mean AUC for total ezetimibe was increased approximately 4-fold on Day 1 and Day 14 compared to healthy subjects. No dosage adjustment is necessary for patients with mild hepatic impairment. Due to the unknown effects of the

increased exposure to ezetimibe in patients with moderate or severe (Child-Pugh score >9) hepatic impairment, ezetimibe is not recommended in these patients [see 8. WARNINGS AND PRECAUTIONS, 8.5 Hepatic Impairment].

Atorvastatin

Plasma concentrations of atorvastatin and its active metabolites are markedly increased (approx. 16-fold in C_{max} and approx. 11-fold in AUC) in patients with chronic alcoholic liver disease (Child-PughB).

Pediatric

Ezetimibe

The absorption and metabolism of ezetimibe are similar between children and adolescents (10 to 18 years) and adults. Based on total ezetimibe, there are no pharmacokinetic differences between adolescents and adults. Pharmacokinetic data in the pediatric population < 10 years of age are not available.

Atorvastatin

Apparent oral clearance of atorvastatin in pediatric subjects appeared similar to that of adults when scaled allometrically by body weight as the body weight was the only significant covariate in atorvastatin population PK model with data including pediatric HeFH patients (ages 10 years to 17 years of age, n=29) in an open-label, 8-week study.

Geriatric

Ezetimibe

Plasma concentrations for total ezetimibe are about 2-fold higher in the elderly (≥ 65 years) than in the young (18 to 45 years). LDL-C reduction and safety profile are comparable between elderly and young subjects treated with ezetimibe.

Atorvastatin

Plasma concentrations of atorvastatin and its active metabolites are higher in healthy elderly subjects than in young adults while the lipid effects were comparable to those seen in younger patient populations.

Race

Based on a meta analysis of pharmacokinetic studies with ezetimibe, there were no pharmacokinetic differences between Blacks and Caucasians.

Gender

Ezetimibe

Plasma concentrations for total ezetimibe are slightly higher (<20 %) in women than in men. LDL-C reduction and safety profile are comparable between men and women treated with ezetimibe.

Atorvastatin

Concentrations of atorvastatin and its active metabolites in women differ from those in men (women: approx. 20% higher for C_{max} and approx. 10% lower for AUC). These differences were of no clinical significance, resulting in no clinically significant differences in lipid effects among men and women.

Hemodialysis

Atorvastatin

While studies have not been conducted in patients with end-stage renal disease, hemodialysis is not expected to significantly enhance clearance of atorvastatin since the drug is extensively bound to plasma proteins.

5. INDICATIONS AND USAGE

Prevention of Cardiovascular Events

ATOZET is indicated to reduce the risk of cardiovascular events [cardiovascular death, nonfatal myocardial infarction, nonfatal stroke, hospitalization for unstable angina, or need for revascularization] in patients with coronary heart disease (CHD) and a history of acute coronary syndrome (ACS), either previously treated with a statin or not.

Hypercholesterolaemia

ATOZET is indicated as adjunctive therapy to diet for use in adults with primary (heterozygous familial and non-familial) hypercholesterolaemia or mixed hyperlipidaemia where use of a combination product is appropriate

- patients not appropriately controlled with a statin alone
- patients already treated with a statin and ezetimibe

Homozygous Familial Hypercholesterolaemia (HoFH)

ATOZET is indicated as adjunctive therapy to diet for use in adults with HoFH. Patients may also receive adjunctive treatments (e.g., low-density lipoprotein [LDL] apheresis).

6. DOSAGE AND ADMINISTRATION

Hypercholesterolaemia and/or Coronary Heart Disease (with ACS History)

The patient should be on an appropriate lipid lowering diet and should continue on this diet during treatment with ATOZET.

The dose range of ATOZET is 10/10 mg/day through 10/80 mg/day. The typical dose is 10/10 mg once a day. The patient's low-density lipoprotein cholesterol (LDL-C) level, coronary heart disease risk status, and response to current cholesterol-lowering therapy should be considered when starting therapy or adjusting the dose.

The dose of ATOZET should be individualised based on the known efficacy of the various dose strengths of ATOZET (see section 14, Table 2) and the response to the current cholesterol-lowering therapy. Adjustment of dose should be made at intervals of 4 weeks or more.

Homozygous Familial Hypercholesterolaemia

The dose of ATOZET in patients with homozygous FH is 10/10 to 10/80 mg daily. ATOZET may be used as an adjunct to other lipid-lowering treatments (e.g., LDL apheresis) in these patients or if such treatments are unavailable.

Coadministration with bile acid sequestrants

Dosing of ATOZET should occur either ≥ 2 hours before or ≥ 4 hours after administration of a bile acid sequestrant.

Geriatric patients

No dose adjustment is required for older patients [see 4. CLINICAL PHARMACOLOGY, 4.3 Pharmacokinetics].

Paediatric population

The safety and efficacy of ATOZET in children has not been established [see 4. CLINICAL PHARMACOLOGY, 4.3 Pharmacokinetics]. No data are available.

Patients with hepatic impairment

ATOZET should be used with caution in patients with hepatic impairment [see 8. WARNINGS AND PRECAUTIONS and 4. CLINICAL PHARMACOLOGY, 4.3 Pharmacokinetics]. ATOZET is contraindicated in patients with active liver disease [see 7. CONTRAINDICATIONS].

Patients with renal impairment

No dose adjustment is required for renally impaired patients [see 4. *CLINICAL PHARMACOLOGY*, 4.3 *Pharmacokinetics*].

Cyclosporine, Clarithromycin, Itraconazole, or Certain HIV/HCV Antiviral Agents

In patients taking cyclosporine or the HIV protease inhibitors tipranavir plus ritonavir or the hepatitis C protease inhibitor telaprevir, therapy with ATOZET should be avoided. In patients with HIV taking lopinavir plus ritonavir, caution should be used when prescribing ATOZET and the lowest dose necessary employed. In patients taking clarithromycin, itraconazole, the hepatitis C antiviral agents boceprevir, elbasvir, grazoprevir, or in patients with HIV taking a combination of saquinavir plus ritonavir, darunavir plus ritonavir, fosamprenavir, or fosamprenavir plus ritonavir, therapy with ATOZET should be limited to 10/20 mg, and appropriate clinical assessment is recommended to ensure that the lowest dose necessary of atorvastatin is employed. In patients taking the HIV protease inhibitor nelfinavir therapy with ATOZET should be limited to 10/40 mg, and appropriate clinical assessment is recommended to ensure that the lowest dose necessary of atorvastatin is employed. [See 8. *WARNINGS AND PRECAUTIONS*, 8.1 *Myopathy/Rhabdomyolysis*.]

Other Concomitant Lipid-Lowering Therapy

The combination of ATOZET and fibrates is not recommended.

Method of administration

ATOZET is for oral administration. ATOZET can be administered as a single dose at any time of the day, with or without food.

7. CONTRAINDICATIONS

- ATOZET is contraindicated in patients with hypersensitivity to ezetimibe, atorvastatin, or any of its inactive ingredients.
- Active liver disease or unexplained persistent elevations of serum transaminases.
- Pregnancy and nursing [see 10. *USE IN SPECIFIC POPULATIONS*, 10.1. *Pregnancy* and 10.2. *Nursing Mothers*].

8. WARNINGS AND PRECAUTIONS

8.1 Myopathy/Rhabdomyolysis

Rare cases of rhabdomyolysis with acute renal failure secondary to myoglobinuria have been reported with atorvastatin and with other drugs in this class. A history of renal impairment may be a risk factor for the development of rhabdomyolysis. Such patients merit closer monitoring for skeletal muscle effects.

Atorvastatin, like other statins, occasionally causes myopathy, defined as muscle aches or muscle weakness in conjunction with increases in CPK values >10 times ULN. Myopathy should be considered in any patient with diffuse myalgias, muscle tenderness or weakness, and/or marked elevation of CPK. Patients should be advised to report promptly unexplained muscle pain, tenderness or weakness, particularly if accompanied by malaise or fever or if muscle signs and symptoms persist after discontinuing ATOZET. ATOZET therapy should be discontinued if markedly elevated CPK levels occur or myopathy is diagnosed or suspected [see 12. ADVERSE REACTIONS].

The risk of myopathy during treatment with statins is increased with concurrent administration of cyclosporine, fibric acid derivatives, erythromycin, clarithromycin, the hepatitis C antiviral agents telaprevir, elbasvir, grazoprevir, combinations of HIV protease inhibitors, including saquinavir plus ritonavir, lopinavir plus ritonavir, tipranavir plus ritonavir, darunavir plus ritonavir, fosamprenavir, and fosamprenavir plus ritonavir, niacin, or azole antifungals. Physicians considering combined therapy with ATOZET and fibrates, erythromycin, immunosuppressive drugs, azole antifungals, or lipid-modifying doses of niacin ($\geq 1\text{g/day}$) should carefully weigh the potential benefits and risks and should carefully monitor patients for any signs or symptoms of muscle pain, tenderness, or weakness, particularly during the initial months of therapy and during any periods of upward dosage titration of either drug. Therefore, lower starting and maintenance doses of ATOZET should also be considered when taken concomitantly with the aforementioned drugs. Temporary suspension of ATOZET may be appropriate during fusidic acid therapy. [See 9. DRUG INTERACTIONS AND OTHER FORMS OF INTERACTIONS, 9.2 CYP3A4 Interactions.] Periodic CPK determinations may be considered in such situations, but there is no assurance that such monitoring will prevent the occurrence of severe myopathy.

Prescribing recommendations for interacting agents are summarized in Table 1 [see 6. DOSAGE AND ADMINISTRATION, 6.8 Cyclosporine, Clarithromycin, Itraconazole, or Certain Protease Inhibitors, and 9. DRUG INTERACTIONS AND OTHER FORMS OF INTERACTIONS, 9.2 CYP3A4 Interactions].

Table 1

**Atorvastatin Drug Interactions Associated with Increased Risk of
Myopathy/Rhabdomyolysis**

Interacting Agents	Prescribing Recommendations for ATOZET
Cyclosporine, HIV protease inhibitors (tipranavir plus ritonavir), hepatitis C protease inhibitor (telaprevir), gemfibrozil	Avoid ATOZET.
Other fibrates (except fenofibrate), fusidic acid	Not recommended with ATOZET
HIV protease inhibitor (lopinavir plus ritonavir)	Use with caution and lowest dose necessary.
Clarithromycin, itraconazole, HIV protease inhibitors (saquinavir plus ritonavir*, darunavir plus ritonavir, fosamprenavir, fosamprenavir plus ritonavir), hepatitis C antiviral agents (boceprevir, elbasvir, grazoprevir)	Do not exceed 10/20 mg ATOZET daily.
HIV protease inhibitor (nelfinavir)	Do not exceed 10/40 mg ATOZET daily.

* Use with caution and with the lowest dose necessary

Cases of myopathy, including rhabdomyolysis, have been reported with atorvastatin coadministered with colchicine, and caution should be exercised when prescribing ATOZET with colchicine [see 9. DRUG INTERACTIONS AND OTHER FORMS OF INTERACTIONS, 9.3 Other Interactions].

Reports of myopathy and/or rhabdomyolysis have been observed with HMG-CoA reductase inhibitors coadministered with daptomycin. Caution should be used when prescribing HMG-CoA reductase inhibitors with daptomycin, as either agent can cause myopathy and/or rhabdomyolysis when given alone. Consideration should be given to suspending ATOZET temporarily in patients taking daptomycin [see 9. DRUG INTERACTIONS AND OTHER FORMS OF INTERACTIONS, 9.3 Other Interactions].

ATOZET therapy should be temporarily withheld or discontinued in any patient with an acute, serious condition suggestive of a myopathy or having a risk factor predisposing to the development of renal failure secondary to rhabdomyolysis (e.g., severe acute infection, hypotension, major surgery, trauma, severe metabolic, endocrine and electrolyte disorders, and uncontrolled seizures).

There have been very rare reports of an immune-mediated necrotizing myopathy (IMNM) during or after treatment with some statins. IMNM is clinically characterized by:

- persistent proximal muscle weakness and elevated serum creatine kinase, which persist despite discontinuation of statin treatment;
- muscle biopsy showing necrotizing myopathy without significant inflammation;
- improvement with immunosuppressive agents.

8.2 Myasthenia Gravis/Ocular Myasthenia

In few cases, statins have been reported to induce de novo or aggravate pre-existing myasthenia gravis or ocular myasthenia [See 12. *ADVERSE REACTIONS*]. ATOZET should be discontinued in case of aggravation of symptoms. Recurrences when the same or a different statin was (re-) administered have been reported.

8.3 Liver Enzymes

In controlled coadministration trials in patients receiving ezetimibe with atorvastatin, consecutive transaminase elevations ($\geq 3 \times$ the upper limit of normal [ULN]) have been observed. [See 12. *ADVERSE REACTIONS*.]

Atorvastatin, like some other lipid-lowering therapies, has been associated with biochemical abnormalities of liver function.

It is recommended that liver enzyme tests be obtained prior to initiating therapy with ATOZET and repeated as clinically indicated. There have been rare postmarketing reports of fatal and non-fatal hepatic failure in patients taking statins, including atorvastatin. If serious liver injury with clinical symptoms and/or hyperbilirubinemia or jaundice occurs during treatment with ATOZET, promptly interrupt therapy. If an alternate etiology is not found, do not restart ATOZET.

ATOZET should be used with caution in patients who consume substantial quantities of alcohol and/or have a history of liver disease. Active liver disease or unexplained persistent transaminase elevations are contraindications to the use of atorvastatin [see 7. *CONTRAINDICATIONS*].

8.4 Endocrine Function

Statins interfere with cholesterol synthesis and theoretically might blunt adrenal and/or gonadal steroid production. Clinical studies have shown that atorvastatin does not reduce basal plasma cortisol concentration or impair adrenal reserve. The effects of statins on male fertility have not been studied in adequate numbers of patients. The effects, if any, on the pituitary-gonadal axis in premenopausal women are unknown. Caution should be exercised if ATOZET is administered

concomitantly with drugs that may decrease the levels or activity of endogenous steroid hormones, such as ketoconazole, spironolactone, and cimetidine.

8.5 Hepatic Impairment

Due to the unknown effects of the increased exposure to ezetimibe in patients with moderate or severe hepatic impairment, ATOZET is not recommended in these patients.

8.6 Fibrates [see 9. DRUG INTERACTIONS AND OTHER FORMS OF INTERACTIONS, 9.3 Other Interactions]

Gemfibrozil: Concomitant administration of ATOZET with gemfibrozil should be avoided.

Fenofibrate: Caution should be used when prescribing ATOZET and fenofibrate, as fenofibrate can cause myopathy when give alone.

If cholelithiasis is suspected in a patient receiving ATOZET and fenofibrate, gallbladder studies are indicated and alternative lipid-lowering therapy should be considered [see the product labeling for fenofibrate and fenofibric acid].

Other fibrates: The coadministration of ezetimibe with other fibrates has not been studied. Therefore, coadministration of ATOZET and other fibrates is not recommended.

8.7 Fusidic acid

Patients on fusidic acid treated concomitantly with ATOZET may have an increased risk of myopathy/rhabdomyolysis [see 9. DRUG INTERACTIONS AND OTHER FORMS OF INTERACTIONS, 9.3 Other Interactions]. Coadministration with fusidic acid is not recommended. In patients where the use of systemic fusidic acid is considered essential, ATOZET should be discontinued throughout the duration of fusidic acid treatment. In exceptional circumstances, where prolonged systemic fusidic acid is needed, e.g. for the treatment of severe infections, the need for coadministration of ATOZET and fusidic acid should only be considered on a case-by-case basis under close medical supervision.

8.8 Anticoagulants

If ATOZET is added to warfarin, another coumarin anticoagulant, or fluindione, the International Normalized Ratio (INR) should be appropriately monitored [see 9. DRUG INTERACTIONS AND OTHER FORMS OF INTERACTIONS, 9.3 Other Interactions].

8.9 Use in Patients with Recent Stroke or TIA

In a post-hoc analysis of the Stroke Prevention by Aggressive Reduction in Cholesterol Levels (SPARCL) study where atorvastatin 80 mg vs. placebo was administered in 4,731 subjects without CHD who had a stroke or TIA within the preceding 6 months, a higher incidence of hemorrhagic stroke was seen in the atorvastatin 80 mg group compared to placebo. The incidence of fatal hemorrhagic stroke was similar across treatment groups. The incidence of nonfatal hemorrhagic stroke was significantly higher in the atorvastatin group as compared to the placebo group. Some baseline characteristics, including hemorrhagic and lacunar stroke on study entry, were associated with a higher incidence of hemorrhagic stroke in the atorvastatin group.

9. DRUG INTERACTIONS AND OTHER FORMS OF INTERACTIONS

9.1 ATOZET

No clinically significant pharmacokinetic interaction was seen when ezetimibe was coadministered with atorvastatin.

Multiple mechanisms may contribute to potential interactions with HMG Co-A reductase inhibitors. Drugs or herbal products that inhibit certain enzymes (e.g. CYP3A4) and/or transporter (e.g. OATP1B) pathways may increase atorvastatin plasma concentrations and may lead to an increased risk of myopathy/rhabdomyolysis.

Consult the prescribing information of all concomitantly used drugs to obtain further information about their potential interactions with atorvastatin and/or the potential for enzyme or transporter alterations and possible adjustments to dose and regimens.

9.2 CYP3A4 Interactions

In preclinical studies, it has been shown that ezetimibe does not induce cytochrome P450 drug metabolizing enzymes. No clinically significant pharmacokinetic interactions have been observed between ezetimibe and drugs known to be metabolized by cytochromes P450 1A2, 2D6, 2C8, 2C9, and 3A4, or N-acetyltransferase. Atorvastatin is metabolized by cytochrome P450 3A4. Concomitant administration of atorvastatin with inhibitors of cytochrome P450 3A4 can lead to increases in plasma concentrations of atorvastatin. The extent of interaction and potentiation of effects depends on the variability of effect on cytochrome P450 3A4.

Inhibitors of cytochrome P3A4 increase the risk of myopathy by reducing the elimination of the atorvastatin component of ATOZET [see 8. WARNINGS AND PRECAUTIONS, 8.1

Myopathy/Rhabdomyolysis]:

Clarithromycin: Atorvastatin AUC was significantly increased with concomitant administration of 80 mg atorvastatin with clarithromycin (500 mg twice daily) compared to that of atorvastatin alone. Therefore, in patients taking clarithromycin, caution should be used when the ATOZET dose exceeds 10/20 mg [see 8. WARNINGS AND PRECAUTIONS, 8.1 *Myopathy/Rhabdomyolysis*, and 6. DOSAGE AND ADMINISTRATION, 6.8 *Cyclosporine, Clarithromycin, Itraconazole, or Certain Protease Inhibitors*].

Combination of Protease Inhibitors: Atorvastatin AUC was significantly increased with concomitant administration of atorvastatin with several combinations of HIV protease inhibitors, as well as with the hepatitis C protease inhibitor telaprevir, compared to that of atorvastatin alone. Therefore, in patients taking the HIV protease inhibitor tipranavir plus ritonavir, or the hepatitis C protease inhibitor telaprevir, concomitant use of ATOZET should be avoided. In patients taking the HIV protease inhibitor lopinavir plus ritonavir, caution should be used when prescribing ATOZET, and the lowest dose necessary should be used. In patients taking the HIV protease inhibitors saquinavir plus ritonavir, darunavir plus ritonavir, fosamprenavir, or fosamprenavir plus ritonavir, or the hepatitis C protease inhibitor boceprevir, the dose of ATOZET should not exceed 10/20 mg and should be used with caution [see 8. WARNINGS AND PRECAUTIONS, 8.1 *Myopathy/Rhabdomyolysis*, and 6. DOSAGE AND ADMINISTRATION, 6.8 *Cyclosporine, Clarithromycin, Itraconazole, or Certain Protease Inhibitors*]. In patients taking the HIV protease inhibitor nelfinavir the dose of ATOZET should not exceed 10/40 mg and close clinical monitoring is recommended.

Itraconazole: Atorvastatin AUC was significantly increased with concomitant administration of atorvastatin 40 mg and itraconazole 200 mg. Therefore, in patients taking itraconazole, caution should be used when the ATOZET dose exceeds 10/20 mg.

Grapefruit Juice: Contains one or more components that inhibit CYP 3A4 and can increase plasma concentrations of atorvastatin, especially with excessive grapefruit juice consumption (>1.2 liters per day).

Cyclosporine: In a study of eight post-renal transplant patients with creatinine clearance of >50 mL/min on a stable dose of cyclosporine, a single 10-mg dose of ezetimibe resulted in a 3.4- fold (range 2.3- to 7.9-fold) increase in the mean AUC for total ezetimibe compared to a healthy control population from another study (n=17). In a different study, a renal transplant patient with severe renal insufficiency (creatinine clearance of 13.2 mL/min/1.73 m²) who was receiving multiple medications, including cyclosporine,

demonstrated a 12-fold greater exposure to total ezetimibe compared to concurrent controls. In a two-period crossover study in twelve healthy subjects, daily administration of 20 mg ezetimibe for 8 days with a single 100-mg dose of cyclosporine on Day 7 resulted in a mean 15% increase in cyclosporine AUC (range 10% decrease to 51% increase) compared to a single 100-mg dose of cyclosporine alone [see 8. WARNINGS AND PRECAUTIONS, 8.1 Myopathy/Rhabdomyolysis].

Atorvastatin and atorvastatin-metabolites are substrates of the OATP1B1 transporter. Inhibitors of the OATP1B1 (e.g., cyclosporine) can increase the bioavailability of atorvastatin. Atorvastatin AUC was significantly increased with concomitant administration of atorvastatin 10 mg and cyclosporine 5.2 mg/kg/day compared to that of atorvastatin alone. The coadministration of ATOZET with cyclosporine should be avoided [see 8. WARNINGS AND PRECAUTIONS, 8.1 Myopathy/Rhabdomyolysis].

9.3 Other Interactions

Antacids: Concomitant antacid administration decreased the rate of absorption of ezetimibe but had no effect on the bioavailability of ezetimibe. This decreased rate of absorption is not considered clinically significant.

Coadministration of atorvastatin with an oral antacid suspension containing magnesium and aluminium hydroxides decreased plasma concentrations of atorvastatin and its active metabolites approximately 35%; however, LDL-C reduction was not altered.

Cholestyramine: Concomitant cholestyramine administration decreased the mean AUC of total ezetimibe (ezetimibe + ezetimibe glucuronide) approximately 55%. The incremental LDL-C reduction due to adding ezetimibe to cholestyramine may be lessened by this interaction.

Fibrates [see 8. WARNINGS AND PRECAUTIONS, 8.6 Fibrates]:

Gemfibrozil: Due to an increased risk of myopathy/rhabdomyolysis when HMG-CoA reductase inhibitors are coadministered with gemfibrozil, concomitant administration of ATOZET with gemfibrozil should be avoided.

In a pharmacokinetic study, concomitant gemfibrozil administration increased total ezetimibe concentrations approximately 1.7-fold. This increase is not considered clinically significant. No clinical data are available.

Fenofibrate: Because it is known that the risk of myopathy during treatment with HMG-CoA

reductase inhibitors is increased with concurrent administration of fenofibrate, ATOZET should be administered with caution when used concomitantly with fenofibrate.

In a pharmacokinetic study, concomitant fenofibrate administration increased total ezetimibe concentrations approximately 1.5-fold. This increase is not considered clinically significant.

Other fibrates: The safety and effectiveness of ezetimibe administered with other fibrates have not been established. Fibrates may increase cholesterol excretion into the bile, leading to cholelithiasis. In a preclinical study in dogs, ezetimibe increased cholesterol in the gallbladder bile. Although the relevance of this preclinical finding to humans is unknown, coadministration of ATOZET with other fibrates is not recommended until use in patients is studied.

Fusidic Acid: The risk of myopathy/rhabdomyolysis may be increased by concomitant administration of fusidic acid [see 8. WARNINGS AND PRECAUTIONS, 8.1 Myopathy/Rhabdomyolysis].

Anticoagulants: Concomitant administration of ezetimibe (10 mg once daily) had no significant effect on bioavailability of warfarin and prothrombin time in a study of twelve healthy adult males. There have been post-marketing reports of increased INR in patients who had ezetimibe added to warfarin or fluindione. Most of these patients were also on other medications. [See 8. WARNINGS AND PRECAUTIONS, 8.8 Anticoagulants.]

Atorvastatin had no clinically significant effect on prothrombin time when administered to patients receiving chronic warfarin treatment.

The effect of ATOZET on the prothrombin time has not been studied.

Inhibitors of Breast Cancer Resistant Protein (BCRP): Atorvastatin is a substrate of the efflux transporter BCRP. Concomitant administration of products that are inhibitors of BCRP (e.g., elbasvir and grazoprevir) may lead to increased plasma concentrations of atorvastatin and an increased risk of myopathy; therefore, a dose adjustment of atorvastatin may be necessary. Coadministration of elbasvir and grazoprevir with atorvastatin increases plasma concentrations of atorvastatin by 1.9-fold due in part to CYP3A and/or BCRP inhibition; therefore, the dose of ATOZET should not exceed 10/20 mg daily in patients receiving concomitant medication with products containing elbasvir or grazoprevir [see 8. WARNINGS AND PRECAUTIONS, 8.1 Myopathy/Rhabdomyolysis].

Inducers of Cytochrome P450 3A4: Concomitant administration of atorvastatin with inducers of cytochrome P450 3A4 (e.g., efavirenz, rifampin) can lead to variable reductions in plasma concentrations of atorvastatin. Due to the dual interaction mechanism of rifampin, simultaneous

coadministration of atorvastatin with rifampin is recommended, as delayed administration of atorvastatin after administration of rifampin has been associated with a significant reduction in atorvastatin plasma concentrations.

Antipyrine: Because atorvastatin does not affect the pharmacokinetics of antipyrine, interactions with other drugs metabolized via the same cytochrome isozymes are not expected.

Colestipol: Plasma concentrations of atorvastatin decreased approximately 25% when colestipol and atorvastatin were coadministered. However, LDL-C reduction was greater when atorvastatin and colestipol were coadministered than when either drug was given alone.

Digoxin: When multiple doses of atorvastatin and digoxin were coadministered, steady-state plasma digoxin concentrations increased by approximately 20%. Patients taking digoxin should be monitored appropriately.

Oral Contraceptives: Coadministration of atorvastatin and an oral contraceptive increased AUC values for norethindrone and ethinyl estradiol by approximately 30% and 20%. These increases should be considered when selecting an oral contraceptive for a woman taking atorvastatin.

Amlodipine: In a drug-drug interaction study in healthy subjects, coadministration of atorvastatin 80 mg and amlodipine 10 mg resulted in an 18% increase in exposure to atorvastatin that was not clinically meaningful.

Niacin: The risk of skeletal muscle effects may be enhanced when ATOZET is used in combination with niacin; a reduction in ATOZET dosage should be considered in this setting [*see 8. WARNINGS AND PRECAUTIONS, 8.1 Myopathy/Rhabdomyolysis*].

Colchicine: Cases of myopathy, including rhabdomyolysis, have been reported with atorvastatin coadministered with colchicine, and caution should be exercised when prescribing ATOZET with colchicine.

Daptomycin: The risk of myopathy and/or rhabdomyolysis may be increased by concomitant administration of HMG-CoA reductase inhibitors and daptomycin [*see 8. WARNINGS AND PRECAUTIONS, 8.1 Myopathy/Rhabdomyolysis*].

10. USE IN SPECIFIC POPULATIONS

10.1 Pregnancy

Atherosclerosis is a chronic process and discontinuation of lipid-lowering drugs during pregnancy should have little impact on the outcome of long-term therapy of primary hypercholesterolemia.

ATOZET

ATOZET is contraindicated during pregnancy. Because HMG-CoA reductase inhibitors decrease cholesterol synthesis and possibly the synthesis of other biologically active substances derived from cholesterol, ATOZET may cause fetal harm when administered to pregnant women. ATOZET should be discontinued as soon as pregnancy is recognized [see 7. CONTRAINDICATIONS]. Advise females of reproductive potential to use effective contraception during treatment of ATOZET.

Ezetimibe

No clinical data on exposed pregnancies are available. Animal studies of ezetimibe administered alone do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/fetal development, parturition or postnatal development.

When ezetimibe was given with statins, no teratogenic effects were observed in embryo-fetal development studies in pregnant rats. In pregnant rabbits, a low incidence of skeletal malformations was observed.

Atorvastatin

Limited published data on atorvastatin from observational studies, meta-analyses and case reports have not shown an increased risk of major congenital malformations or miscarriage. Rare reports of congenital anomalies have been received following intrauterine exposure to other HMG-CoA reductase inhibitors. In a review of approximately 100 prospectively followed pregnancies in women exposed to simvastatin or lovastatin, the incidences of congenital anomalies, spontaneous abortions, and fetal deaths/stillbirths did not exceed what would be expected in the general population. The number of cases is adequate to exclude a greater than or equal to three-to-four-fold increase in congenital anomalies over background incidence. In 89% of the prospectively followed pregnancies, drug treatment was initiated prior to pregnancy and was discontinued at some point in the first trimester when pregnancy was identified.

In animal reproduction studies in rats and rabbits there was no evidence of embryo-fetal toxicity or

congenital malformations at doses up to 30 and 20 times, respectively, the human exposure at the maximum recommended human dose (MRHD) of 80 mg, based on body surface area (mg/m²). In rats administered atorvastatin during gestation and lactation, decreased postnatal growth and development was observed at doses $\geq$ 6 times the MRHD.

10.2 Nursing Mothers

ATOZET

ATOZET is contraindicated in nursing mothers. Because of the potential for serious adverse reactions in a breastfed infant, women who are nursing should not take ATOZET.

Ezetimibe

Studies in rats have shown that ezetimibe and atorvastatin are excreted in milk. It is not known whether ezetimibe is excreted into human breast milk.

Atorvastatin

There is no available information on the effects of atorvastatin on the breastfed infant or the effects of atorvastatin on milk production. It is not known whether atorvastatin is present in human milk, but it has been shown that another drug in this class passes into human milk and atorvastatin is present in rat milk.

10.3 Pediatric Use

There are insufficient data for the safe and effective use of ATOZET in pediatric patients.

11. EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

No studies of the effects on the ability to drive and use of machines have been performed. However, certain side effects that have been reported with ATOZET may affect some patients' ability to drive or operate machinery. Individual responses to ATOZET may vary. [See 12. ADVERSE REACTIONS.]

12. ADVERSE REACTIONS

12.1 Clinical Trials Experience

Adults

ATOZET

ATOZET (or coadministration of ezetimibe and atorvastatin equivalent to ATOZET) has been evaluated for safety in more than 2400 patients in 7 clinical trials. ATOZET was generally well tolerated.

The following common ($\geq 1/100$, $<1/10$) or uncommon ($\geq 1/1000$, $<1/100$) drug-related adverse experiences were reported in patients taking ATOZET:

Infections and Infestations:

Uncommon: influenza

Psychiatric disorders:

Uncommon: depression; insomnia; sleep disorder

Nervous system disorders:

Uncommon: dizziness; dysgeusia; headache; paresthesia

Cardiac disorders:

Uncommon: sinus bradycardia

Vascular disorders:

Uncommon: hot flush

Respiratory, thoracic and mediastinal disorders:

Uncommon: dyspnea

Gastrointestinal disorders:

Common: diarrhea

Uncommon: abdominal discomfort; abdominal distension; abdominal pain; abdominal pain lower; abdominal pain upper; constipation; dyspepsia; flatulence; frequent bowel movements; gastritis; nausea; stomach discomfort

Skin and subcutaneous tissue disorders:

Uncommon: acne; urticaria

Musculoskeletal and connective tissue disorders:

Common: myalgia

Uncommon: arthralgia; back pain; muscle fatigue; muscle spasms; muscular weakness; pain in extremity

General disorders and administration site conditions:

Uncommon: asthenia; fatigue; malaise; edema

Investigations:

Uncommon: ALT and/or AST increased; alkaline phosphatase increased; blood CK increased; gamma-glutamyltransferase increased; hepatic enzyme increased; liver function test abnormal; weight increased

In controlled clinical trials, the incidence of clinically important elevations in serum transaminases (ALT and/or AST $\geq 3 \times$ ULN, consecutive) was 0.6% for patients treated with ATOZET. These elevations in transaminases were generally asymptomatic, not associated with cholestasis, and returned to baseline spontaneously or after discontinuation of therapy. [See 8. WARNINGS AND PRECAUTIONS, 8.3 Liver Enzymes.]

None of the patients treated with ATOZET had CK levels $\geq 10 \times$ ULN.

12.2 Post-marketing Experience and Other Clinical Trial Experience

The following additional adverse reactions have been reported in post-marketing use with ATOZET or in clinical studies or post-marketing use with ezetimibe or atorvastatin:

Infections and infestations: nasopharyngitis

Blood and lymphatic system disorders: thrombocytopenia

Immune system disorders: hypersensitivity reactions, including anaphylaxis, angioedema, rash, and urticaria

Metabolism and nutrition disorders: decreased appetite; anorexia; hyperglycemia; hypoglycemia

Psychiatric disorders: nightmares

Nervous system disorders: hypesthesia; peripheral neuropathy;
Frequency 'not known': myasthenia gravis [see 8. WARNINGS AND PRECAUTIONS, 8.2 Myasthenia Gravis/Ocular Myasthenia].

There have been rare post-marketing reports of cognitive impairment (e.g., memory loss, forgetfulness, amnesia, memory impairment, confusion) associated with statin use. These cognitive

issues have been reported for all statins. The reports are generally nonserious, and reversible upon statin discontinuation, with variable times to symptom onset (1 day to years) and symptom resolution (median of 3 weeks).

Eye disorders: vision blurred; visual disturbance;
Frequency 'not known': ocular myasthenia [see 8. WARNINGS AND PRECAUTIONS, 8.2 Myasthenia Gravis/Ocular Myasthenia].

Ear and labyrinth disorders: tinnitus; hearing loss

Vascular disorders: hypertension

Respiratory, thoracic, and mediastinal disorders: cough; pharyngolaryngeal pain; epistaxis

Gastrointestinal disorders: dry mouth; pancreatitis; gastroesophageal reflux disease; eructation; vomiting

Hepatobiliary disorders: hepatitis; cholelithiasis; cholecystitis; cholestasis; drug-induced liver injury

Skin and subcutaneous tissue disorders: alopecia; pruritus; skin rash; angioneurotic oedema; severe cutaneous adverse reactions (SCARs), including Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN), drug reaction with eosinophilic and systemic symptoms (DRESS), dermatitis bullous and erythema multiforme

Musculoskeletal and connective tissue disorders: myopathy/rhabdomyolysis [See 8. WARNINGS AND PRECAUTIONS, 8.1 Myopathy/Rhabdomyolysis.]; neck pain; joint swelling; myositis tendonopathy, sometimes complicated by rupture

There have been very rare reports of immune-mediated necrotizing myopathy (IMNM), an autoimmune myopathy, associated with statin use. IMNM is characterized by: proximal muscle weakness and elevated serum creatine kinase, which persist despite discontinuation of statin treatment; muscle biopsy showing necrotizing myopathy without significant inflammation; improvement with immunosuppressive agents [see 8. WARNINGS AND PRECAUTIONS, 8.1 Myopathy/Rhabdomyolysis].

Reproductive system and breast disorders: gynecomastia

General disorders and administration site conditions: chest pain; pain; edema peripheral; pyrexia

Investigations: white blood cells urine positive

Increases in HbA1c and fasting serum glucose levels have been reported with HMG-CoA reductase inhibitors, including atorvastatin. The risk of hyperglycemia, however, is outweighed by the reduction in vascular risk with statins.

The following adverse events have been reported with some statins:

- sexual dysfunction
- depression
- exceptional cases of interstitial lung disease, especially with long term therapy
- diabetes mellitus: frequency will depend on the presence or absence of risk factors (fasting blood glucose ≥ 5.6 mmol/L, BMI >30 kg/m², raised triglycerides, history of hypertension)

13. OVERDOSAGE

ATOZET

No specific treatment of overdosage with ATOZET can be recommended. In the event of an overdose, symptomatic and supportive measures should be employed.

Ezetimibe

In clinical studies, administration of ezetimibe, 50 mg/day to 15 healthy subjects for up to 14 days, 40 mg/day to 18 patients with primary hyperlipidemia for up to 56 days, and 40 mg/day to 27 patients with homozygous sitosterolemia for 26 weeks, was generally well tolerated.

A few cases of overdosage have been reported; most have not been associated with adverse experiences. Reported adverse experiences have not been serious.

Atorvastatin

Due to extensive drug binding to plasma proteins, hemodialysis is not expected to significantly enhance atorvastatin clearance.

14. CLINICAL STUDIES

In controlled clinical studies, ATOZET significantly reduced total-C, LDL-C, Apo B, and TG, and increased HDL-C in patients with hypercholesterolaemia.

Primary Hypercholesterolaemia

In a placebo-controlled study, 628 patients with hyperlipidaemia were randomized to receive placebo, ezetimibe (10 mg), atorvastatin (10 mg, 20 mg, 40 mg, or 80 mg), or coadministered ezetimibe and atorvastatin equivalent to ATOZET (10/10, 10/20, 10/40, and 10/80) for up to 12 weeks.

Patients receiving all doses of ATOZET were compared to those receiving all doses of atorvastatin. ATOZET lowered total-C, LDL-C, Apo B, TG, and non-HDL-C, and increased HDL-C significantly more than atorvastatin alone. (See Table 2.)

Table 2
Response to ATOZET in Patients with Primary Hyperlipidaemia
(Mean^a % Change from Untreated Baseline^b at 12 weeks)

Treatment (Daily Dose)	N	Total-C	LDL-C	Apo B	TG ^a	HDL-C	Non-HDL-C
Pooled data (All ATOZET doses) ^c	255	-41	-56	-45	-33	+7	-52
Pooled data (All atorvastatin doses) ^c	248	-32	-44	-36	-24	+4	-41
Ezetimibe 10 mg	65	-14	-20	-15	-5	+4	-18
Placebo	60	+4	+4	+3	-6	+4	+4
ATOZET by dose							
10/10	65	-38	-53	-43	-31	+9	-49
10/20	62	-39	-54	-44	-30	+9	-50
10/40	65	-42	-56	-45	-34	+5	-52
10/80	63	-46	-61	-50	-40	+7	-58
Atorvastatin by dose							
10 mg	60	-26	-37	-28	-21	+6	-34
20 mg	60	-30	-42	-34	-23	+4	-39
40 mg	66	-32	-45	-37	-24	+4	-41
80 mg	62	-40	-54	-46	-31	+3	-51

^a For triglycerides, median % change from baseline

^b Baseline - on no lipid-lowering medicinal product

^c ATOZET pooled (10/10-10/80 mg) significantly reduced total-C, LDL-C, Apo B, TG, non-HDL-C, and significantly increased HDL-C compared to all doses of atorvastatin pooled (10-80 mg).

In a controlled study, the Titration of Atorvastatin Versus Ezetimibe Add-On to Atorvastatin in Patients with Hypercholesterolaemia (TEMPO) study, 184 patients, with an LDL-C level ≥ 2.6 mmol/L and ≤ 4.1 mmol/L and at moderate high risk for CHD, received atorvastatin 20 mg for a minimum of 4 weeks prior to randomization. Patients not at an LDL-C level <2.6 mmol/L were randomized to receive either coadministered ezetimibe and atorvastatin (equivalent to ATOZET 10/20) or atorvastatin 40 mg

for 6 weeks.

ATOZET 10/20 was significantly more effective than doubling the dose of atorvastatin to 40 mg in further reducing total-C (-20% vs. -7%), LDL-C (-31% vs. -11%), Apo B (-21% vs. -8%), and non-HDL-C (-27% vs. -10%). Results for HDL-C and TG between the two treatment groups were not significantly different. Also, significantly more patients receiving ATOZET 10/20 attained LDL-C <2.6 mmol/L compared to those receiving atorvastatin 40 mg, 84% vs. 49%.

In a controlled study, The Ezetimibe Plus Atorvastatin Versus Atorvastatin Titration in Achieving Lower LDL-C Targets in Hypercholesterolaemic Patients (EZ-PATH) study, 556 high-cardiovascular-risk patients with a LDL-C level ≥ 1.8 mmol/L and ≤ 4.1 mmol/L received atorvastatin 40 mg for a minimum of 4 weeks prior to randomization. Patients not at a LDL-C level <1.8 mmol/L were randomized to receive either coadministered ezetimibe and atorvastatin (equivalent to ATOZET 10/40) or atorvastatin 80 mg for 6 weeks.

ATOZET 10/40 was significantly more effective than doubling the dose of atorvastatin to 80 mg in further reducing total-C (-17% vs. -7%), LDL-C (-27% vs. -11%), Apo B (-18% vs. -8%), TG (-12% vs. -6%), and non-HDL-C (-23% vs. -9%). Results for HDL-C between the two treatment groups were not significantly different. Also, significantly more patients receiving ATOZET 10/40 attained LDL-C <1.8 mmol/L compared to those receiving atorvastatin 80 mg, 74% vs. 32%.

In a placebo-controlled, 8-week study, 308 hypercholesterolemic patients receiving atorvastatin and not at National Cholesterol Education Program (NCEP) LDL-C goal (LDL-C goal based upon baseline LDL-C and CHD risk status) were randomized to receive either ezetimibe 10 mg or placebo in addition to their on-going atorvastatin therapy.

Among patients not at LDL-C goal at baseline (~83%), significantly more patients receiving ezetimibe coadministered with atorvastatin achieved their LDL-C goal compared to patients receiving placebo coadministered with atorvastatin, 67% vs. 19%. Ezetimibe added to atorvastatin therapy lowered LDL-C significantly more than placebo added to atorvastatin therapy, 25% vs. 4%. Ezetimibe added to atorvastatin therapy also significantly decreased total-C, Apo B, and TG compared with placebo added to atorvastatin therapy.

In a controlled, 12-week, 2-phase study, 1539 high-cardiovascular-risk patients, with a LDL-C level between 2.6 and 4.1 mmol/L, on atorvastatin 10 mg daily were randomized to receive: atorvastatin 20 mg, rosuvastatin 10 mg, or ATOZET 10/10. After 6 weeks of treatment (Phase I), patients taking atorvastatin 20 mg who failed to achieve a LDL-C level <2.6 mmol/L were switched to either

atorvastatin 40 mg or ATOZET 10/20 for 6 weeks (Phase II), and similar patients taking rosuvastatin 10 mg during Phase I were switched to either rosuvastatin 20 mg or ATOZET 10/20. Reductions in LDL-C and comparisons between the ATOZET group and other treatment groups studied are shown in Table 3.

Table 3
Response to ATOZET* in High-Risk Patients with a LDL-C Level
Between 2.6 and 4.1 mmol/L on Atorvastatin 10 mg Daily at Baseline

Treatment	N	Percent Change from Baseline†					
		Total-C	LDL-C	Apo B	TG‡	HDL-C	Non-HDL-C
Phase I							
Switched from atorvastatin 10 mg							
ATOZET 10/10	120	-13.5	-22.2	-11.3	-6.0	+0.6	-18.3
Atorvastatin 20 mg	480	-6.4§	-9.5§	-6.0¶	-3.9	-1.1	-8.1§
Rosuvastatin 10 mg	939	-7.7§	-13.0§	-6.9#	-1.1	+1.1	-10.6§
Phase II							
Switched from atorvastatin 20 mg							
ATOZET 10/20	124	-10.7	-17.4	-9.8	-5.9	+0.7	-15.1
Atorvastatin 40 mg	124	-3.8 ^b	-6.9 ^b	-5.4	-3.1	+1.7	-5.8 ^b
Switched from rosuvastatin 10 mg							
ATOZET 10/20	231	-11.8	-17.1	-11.9	-10.2	+0.1	-16.2
Rosuvastatin 20 mg	205	-4.5 ^b	-7.5 ^b	-4.1 ^b	-3.2 [§]	+0.8	-6.4 ^b

* Coadministered ezetimibe and atorvastatin equivalent to ATOZET 10/10 or ATOZET 10/20

† M-Estimates (based on the method of Huber; 95% CI and p-value were obtained from fitting a robust Regression model with terms for treatment and baseline)

‡ Geometric mean percent changes from baseline in TG were calculated based on back-transformation via exponentiation of the model-based least square (LS) means and expressed as (geometric mean – 1) multiplied by 100

§ p<0.001 versus ATOZET 10/10

¶ p<0.01 versus ATOZET 10/10

p<0.05 versus ATOZET 10/10

^b p<0.001 versus ATOZET 10/20

^b p<0.05 versus ATOZET 10/20

Table 3 does not contain data comparing the effects of ATOZET 10/10 or 10/20 to doses higher than atorvastatin 40 mg or rosuvastatin 20 mg.

In a placebo-controlled study, the Myocardial Ischaemia Reduction with Aggressive Cholesterol Lowering (MIRACL) study, patients with an acute coronary syndrome (non Q-wave MI or unstable angina) were randomized to receive atorvastatin 80 mg/day (n=1538) or placebo (n=1548). Treatment was initiated during the acute phase after hospital admission and lasted for 16 weeks. Atorvastatin 80 mg/day provided a 16% (p=0.048) reduction in risk of the combined primary endpoint: death from any cause, nonfatal MI, resuscitated cardiac arrest, or angina pectoris with evidence of myocardial

ischaemia requiring hospitalisation. This was mainly due to a 26% reduction in re-hospitalisation for angina pectoris with evidence of myocardial ischaemia ($p=0.018$).

ATOZET contains atorvastatin. In a placebo-controlled study, the Anglo-Scandinavian Cardiac Outcomes Trial Lipid Lowering Arm (ASCOT-LLA), the effect of atorvastatin 10 mg on fatal and non-fatal CHD was assessed in 10,305 hypertensive patients, 40-80 years old, with TC levels ≤ 6.5 mmol/L and at least three cardiovascular risk factors. Patients were followed for a median duration of 3.3 years. Atorvastatin 10 mg significantly ($p<0.001$) reduced the relative risk for: fatal CHD plus nonfatal MI by 36% (absolute risk reduction = 1.1%); total cardiovascular events and revascularization procedures by 20% (absolute risk reduction = 1.9%); and total coronary events by 29% (absolute risk reduction = 1.4%).

In a placebo-controlled study, the Collaborative Atorvastatin Diabetes Study (CARDS), the effect of atorvastatin 10 mg on cardiovascular disease (CVD) endpoints was assessed in 2838 patients, 40-75 years old, with type 2 diabetes, one or more cardiovascular risk factors, LDL ≤ 4.1 mmol/L, and TG ≤ 6.8 mmol/L. Patients were followed for a median duration of 3.9 years. Atorvastatin 10 mg significantly ($p<0.05$) reduced: the rate of major cardiovascular events by 37% (absolute risk reduction = 3.2%); the risk of stroke by 48% (absolute risk reduction = 1.3%); and the risk of MI by 42% (absolute risk reduction = 1.9%).

ATOZET

Prevention of Cardiovascular Events

In an ezetimibe/simvastatin, multicenter, randomized, double-blind, active-control study, 18,144 patients enrolled within 10 days of hospitalisation for acute coronary syndrome (ACS; either acute myocardial infarction [MI] or unstable angina [UA]). All patients were randomized in a 1:1 ratio to receive either ezetimibe/simvastatin 10/40 mg ($n=9067$) or simvastatin 40 mg ($n=9077$) and followed for a median of 6.0 years.

Patients had a mean age of 63.6 years; 76% were male, 84% were Caucasian, and 27% were diabetic. The average LDL-C value at the time of study qualifying event was 80 mg/dL (2.1 mmol/L) for those on lipid-lowering therapy ($n=6390$) and 101 mg/dL (2.6 mmol/L) for those not on previous lipid-lowering therapy ($n=11594$). Prior to the hospitalisation for the qualifying ACS event, 34% of the patients were on statin therapy. At one-year, the average LDL-C for patients continuing on therapy was 53.2 mg/dL (1.4 mmol/L) for the ezetimibe/simvastatin group and 69.9 mg/dL (1.8 mmol/L) for the simvastatin monotherapy group.


The primary endpoint was a composite consisting of cardiovascular death, major coronary events (MCE; defined as non-fatal myocardial infarction, documented unstable angina that required

hospitalisation, or any coronary revascularization procedure occurring at least 30 days after randomized treatment assignment) and non-fatal stroke. The study demonstrated that treatment with ezetimibe/simvastatin provided incremental benefit in reducing the primary composite endpoint of cardiovascular death, MCE, and non-fatal stroke compared with simvastatin alone (relative risk reduction of 6.4%, $p=0.016$). The primary endpoint occurred in 2572 of 9067 patients (7-year Kaplan-Meier [KM] rate 32.72%) in the ezetimibe/simvastatin group and 2742 of 9077 patients (7-year KM rate 34.67%) in the simvastatin alone group. (See Figure 1 and Table 4.) This incremental benefit is expected to be similar with coadministration of ezetimibe and atorvastatin. Total mortality was unchanged in this high risk group.

There was an overall benefit for all strokes; however there was a small non-significant increase in haemorrhagic stroke in the ezetimibe-simvastatin group compared with simvastatin alone. The risk of haemorrhagic stroke for ezetimibe coadministered with higher potency statins in long-term outcome studies has not been evaluated.

The treatment effect of ezetimibe/simvastatin was generally consistent with the overall results across many subgroups, including sex, age, race, medical history of diabetes mellitus, baseline lipid levels, prior statin therapy, prior stroke, and hypertension.

Figure 1: Effect of ezetimibe/simvastatin on the Primary Composite Endpoint of Cardiovascular Death, Major Coronary Event, or Non-fatal Stroke



Subjects at risk								
Ezetimibe/Simvastatin	9067	7371	6801	6375	5839	4284	3301	1906
Simvastatin	9077	7455	6799	6327	5729	4206	3284	1857

Table 4

Major Cardiovascular Events by Treatment Group in All Randomized Patients in IMPROVE-IT

<u>Outcome</u>	Ezetimibe/Simvastatin		Simvastatin		Hazard Ratio (95% CI)	p-value
	10/40 mg* (N=9067)		40 mg† (N=9077)			
	n	K-M %‡	n	K-M %‡		
Primary Composite Efficacy Endpoint						
(CV death, Major Coronary Events and non-fatal stroke)	2572	32.72%	2742	34.67%	0.936 (0.887, 0.988)	0.016
Components of Primary Composite Endpoint and Select Efficacy Endpoints (first occurrences of specified event at any time)						
Cardiovascular death	537	6.89%	538	6.84%	1.000 (0.887, 1.127)	0.997
Major Coronary Event:						
Non-fatal MI	945	12.77%	1083	14.41%	0.871 (0.798, 0.950)	0.002
Unstable angina requiring hospitalization	156	2.06%	148	1.92%	1.059 (0.846, 1.326)	0.618
Coronary revascularization after 30 days	1690	21.84%	1793	23.36%	0.947 (0.886, 1.012)	0.107
Non-fatal stroke	245	3.49%	305	4.24%	0.802 (0.678, 0.949)	0.010

* 6% were uptitrated to ezetimibe/simvastatin 10/80 mg.

† 27% were uptitrated to simvastatin 80 mg.

‡ Kaplan-Meier estimate at 7 years.

Homozygous Familial Hypercholesterolaemia (HoFH)

A double-blind, randomized, 12-week study was performed in patients with a clinical and/or genotypic diagnosis of HoFH. Data were analyzed from a subgroup of patients (n=36) receiving atorvastatin 40 mg at baseline. Increasing the dose of atorvastatin from 40 to 80 mg (n=12) produced a reduction of LDL-C of 2% from baseline on atorvastatin 40 mg. Coadministered ezetimibe and atorvastatin equivalent to ATOZET (10/40 and 10/80 pooled, n=24), produced a reduction of LDL-C of 19% from baseline on atorvastatin 40 mg. In those patients coadministered ezetimibe and atorvastatin equivalent to ATOZET (10/80, n=12), a reduction of LDL-C of 25% from baseline on atorvastatin 40 mg was produced.

After completing the 12-week study, eligible patients (n=35), who were receiving atorvastatin 40 mg at baseline, were assigned to coadministered ezetimibe and atorvastatin equivalent to ATOZET 10/40 for up to an additional 24 months. Following at least 4 weeks of treatment, the atorvastatin dose could be doubled to a maximum dose of 80 mg. At the end of the 24 months, ATOZET (10/40 and 10/80 pooled) produced a reduction of LDL-C that was consistent with that seen in the 12-week study.

15. STORAGE

Do not store above 30°C.

Store in original package.

16. SHELF LIFE

Please refer to the expiry date on the outer carton.

17. AVAILABILITY

Trade pack of 30 film-coated tablets

18. MANUFACTURED BY:

MSD International GmbH (Singapore Branch)
70 Tuas West Drive, Singapore 638414,
Singapore.

19. PRODUCT REGISTRATION HOLDER

ORGANON MALAYSIA SDN. BHD.

3A-08-02, Level 8, Corporate Tower 3A,

Pavilion Damansara Heights,

No. 3, Jalan Damanlela

50490 Kuala Lumpur, Malaysia.

20. DATE OF REVISION

12 November 2025

Copyright 2025 Organon group of companies. All rights reserved.