

“For the use only of a Registered Medical Practitioner”

IRBEL HCT
(Irbesartan and Hydrochlorothiazide Tablets)

COMPOSITION

IRBEL HCT TABLETS 150 + 12.5 mg

Each film-coated tablet contains:

Irbesartan Ph. Eur.	150 mg
Hydrochlorothiazide Ph.Eur.	12.5 mg

IRBEL HCT TABLETS 300 + 12.5 mg

Each film-coated tablet contains:

Irbesartan Ph. Eur.	300 mg
Hydrochlorothiazide Ph.Eur.	12.5 mg

IRBEL HCT TABLETS 300 + 25 mg

Each film coated tablet contains:

Irbesartan Ph. Eur.	300 mg
Hydrochlorothiazide Ph.Eur.	25 mg

In-active components:

Lactose monohydrate, Microcrystalline Cellulose (PH 101), Croscarmellose Sodium, Hypromellose, Silica, colloidal anhydrous, Microcrystalline Cellulose (PH 102), Magnesium Stearate, Opadry Pink 03G54386 for Irbel 150+12.5 & 300+25 mg and Opadry yellow 03G52390 for Irbel 300+12.5 mg, Water Purified

Coating composition of Opadry Pink 03G54386

Hypromellose, Titanium Dioxide, Macrogol 400, Talc, Macrogol 4000, Iron Oxide red, Iron oxide black


Coating composition of Opadry yellow 03G52390

Hypromellose, Titanium Dioxide, Macrogol 400, Talc, Macrogol 4000, Iron oxide yellow

DESCRIPTION


IRBEL HCT tablets are a fixed dose combination of an angiotensin II receptor antagonist (AT₁ subtype), irbesartan, and a thiazide diuretic, hydrochlorothiazide (HCTZ).

Irbesartan is a non-peptide compound, chemically described as 2-butyl-3-[p-(o-1H-tetrazol-5-ylphenyl)benzyl]-1,3-diazaspiro[4.4]non-1-en-4-one. It has a molecular weight of 428.5; its empirical formula is C₂₅H₂₈N₆O, and its structural formula is as given below:



Irbesartan

Hydrochlorothiazide is 6-chloro-3, 4-dihydro-2*H*-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide. It has a molecular weight of 297.7, its empirical formula is $C_7H_8ClN_3O_4S_2$ and its structural formula is as given below:



Hydrochlorothiazide

Product Description:

For 150 + 12.5 mg strength: Pink to Red coloured, film coated, oval shaped tablets debossed with 'IH1' on one side and plain on other side.

For 300 + 25 mg strength: Pink to Red coloured, film coated, oval shaped tablets debossed with 'IH2' on one side and plain on other side.

For 300 + 12.5 mg strength: Yellow coloured, film coated, oval shaped tablets debossed with 'IH3' on one side and plain on other side.

PHARMACODYNAMIC AND PHARMACOKINETIC PROPERTIES

Pharmacodynamic

This product is a combination of an angiotensin-II receptor antagonist, irbesartan, and a thiazide diuretic, hydrochlorothiazide. The combination of these ingredients has an additive antihypertensive effect, reducing blood pressure to a greater degree than either component alone.

Irbesartan is a potent, orally active, selective angiotensin-II receptor (AT1 subtype) antagonist. It is expected to block all actions of angiotensin-II mediated by the AT1 receptor, regardless of the source or route of synthesis of angiotensin-II. The selective antagonism of the angiotensin-II (AT1) receptors results in increases in plasma renin levels and angiotensin-II levels, and a decrease in plasma aldosterone concentration. Serum potassium levels are not significantly affected by irbesartan alone at the recommended doses in patients without risk of electrolyte imbalance. Irbesartan does not inhibit

ACE (kininase-II), an enzyme which generates angiotensin-II and also degrades bradykinin into inactive metabolites. Irbesartan does not require metabolic activation for its activity.

Hydrochlorothiazide is a thiazide diuretic. The mechanism of antihypertensive effect of thiazide diuretics is not fully known. Thiazides affect the renal tubular mechanisms of electrolyte reabsorption, directly increasing excretion of sodium and chloride in approximately equivalent amounts. The diuretic action of hydrochlorothiazide reduces plasma volume, increases plasma renin activity, increases aldosterone secretion, with consequent increases in urinary potassium and bicarbonate loss, and decreases in serum potassium. Presumably through blockade of the renin-angiotensin-aldosterone system, co-administration of irbesartan tends to reverse the potassium loss associated with these diuretics. With hydrochlorothiazide, onset of diuresis occurs in 2 hours, and peak effect occurs at about 4 hours, while the action persists for approximately 6-12 hours.

The combination of hydrochlorothiazide and irbesartan produces dose-related additive reductions in blood pressure across their therapeutic dose ranges.

The effect of the combination of irbesartan and hydrochlorothiazide on morbidity and mortality has not been reported. It has been reported that long term treatment with hydrochlorothiazide reduces the risk of cardiovascular mortality and morbidity.

There is no difference in response to of irbesartan and hydrochlorothiazide, regardless of age or gender. As is the case with other medicinal products that affect the renin-angiotensin system, black hypertensive patients have notably less response to irbesartan monotherapy. When irbesartan is administered concomitantly with a low dose of hydrochlorothiazide (e.g. 12.5 mg daily), the antihypertensive response in black patients approaches that of non-black patients.

The types and incidences of adverse events reported for patients treated with the irbesartan and hydrochlorothiazide are similar to the adverse event profile for patients on monotherapy.

Non-melanoma skin cancer: Based on available data from epidemiological studies, cumulative dose-dependent association between HCTZ (hydrochlorothiazide) and non-melanoma skin cancer (NMSC) has been observed. One study included a population comprised of 71,533 cases of basal cell carcinoma (BCC) and of 8,629 cases of sec matched to 1,430,833 and 172,462 population controls, respectively. High HCTZ use ($\geq 50,000$ mg cumulative) was associated with an adjusted OR of 1.29 (95% CI: 1.23-1.35) for BCC and 3.98 (95% CI: 3.68-4.31) for squamous cell carcinoma (SCC). A clear cumulative dose response relationship was observed for both BCC and SCC. Another study showed a possible association between lip cancer (SCC) and exposure to HCTZ: 633 cases of lip-cancer were matched with 63,067 population controls, using a risk-set sampling strategy. A cumulative dose-response relationship was demonstrated with an adjusted OR 2.1 (95% CI: 1.7-2.6) increasing to OR 3.9 (3.0-4.9) for high use ($\sim 25,000$ mg) and OR 7.7 (5.7-10.5) for the highest cumulative dose ($\sim 100,000$ mg).

Pharmacokinetics

Concomitant administration of hydrochlorothiazide and irbesartan has no effect on the pharmacokinetics of either medicinal product.

Irbesartan and hydrochlorothiazide are orally active agents and do not require biotransformation for their activity. Following oral administration of combination of irbesartan and hydrochlorothiazide, the absolute oral bioavailability is 60-80% and 50-80% for irbesartan and hydrochlorothiazide, respectively. Food does not affect the bioavailability of combination of irbesartan and hydrochlorothiazide. Peak plasma concentration occurs at 1.5-2 hours after oral administration for irbesartan and 1-2.5 hours for hydrochlorothiazide.

Plasma protein binding of irbesartan is approximately 96%, with negligible binding to cellular blood components. The volume of distribution for irbesartan is 53-93 litres. Hydrochlorothiazide is 68% protein-bound in the plasma, and its apparent volume of distribution is 0.83-1.14 l/kg.

Irbesartan exhibits linear and dose proportional pharmacokinetics over the dose range of 10 to 600 mg. A less than proportional increase in oral absorption at doses beyond 600 mg has been reported; the mechanism for this is unknown. The total body and renal clearance are 157-176 and 3.0-3.5 ml/min, respectively. The terminal elimination half-life of irbesartan is 11–15 hours. Steady-state plasma concentrations are attained within 3 days after initiation of a once-daily dosing regimen. Limited accumulation of irbesartan (< 20%) has been reported in plasma upon repeated once-daily dosing. Higher plasma concentrations of irbesartan have been reported in female hypertensive patients. However, there was no difference in the half-life and accumulation of irbesartan. No dosage adjustment is necessary in female patients. Irbesartan AUC and C_{max} values have also been reported to be greater in elderly subjects (≥ 65 years) than those of young subjects (18-40 years). However the terminal half-life is not significantly altered. No dosage adjustment is necessary in elderly patients. The mean plasma half-life of hydrochlorothiazide reportedly ranges from 5-15 hours.

Following oral or intravenous administration of ^{14}C irbesartan, 80-85% of the circulating plasma radioactivity is attributable to unchanged irbesartan. Irbesartan is metabolized by the liver via glucuronide conjugation and oxidation. The major circulating metabolite is irbesartan glucuronide (approximately 6%). *In vitro* irbesartan is primarily oxidised by the cytochrome P450 enzyme CYP2C9; isoenzyme CYP3A4 has negligible effect. Irbesartan and its metabolites are eliminated by both biliary and renal pathways. After either oral or intravenous administration of ^{14}C irbesartan, about 20% of the radioactivity is recovered in the urine, and the remainder in the faeces. Less than 2% of the dose is excreted in the urine as unchanged irbesartan. Hydrochlorothiazide is not metabolized but is eliminated rapidly by the kidneys. At least 61% of the oral dose is eliminated unchanged within 24 hours. Hydrochlorothiazide crosses the placental but not the blood-brain barrier, and is excreted in breast milk.

Renal impairment: in patients with renal impairment or those undergoing haemodialysis, the pharmacokinetic parameters of irbesartan are not significantly altered. Irbesartan is not removed by haemodialysis. In patients with creatinine clearance <20 ml/min, the elimination half-life of hydrochlorothiazide was reported to increase to 21 hours.

Hepatic impairment: in patients with mild to moderate cirrhosis, the pharmacokinetic parameters of irbesartan are not significantly altered.

INDICATIONS

Treatment of essential hypertension.

This combination is indicated in adult patients whose blood pressure is not adequately controlled on irbesartan or hydrochlorothiazide alone.

DOSE AND METHOD OF ADMINISTRATION

IRBEL HCT Tablets (Irbesartan and Hydrochlorothiazide tablets) can be taken once daily, with or without food.

Dose titration with the individual components (i.e. irbesartan and hydrochlorothiazide) may be recommended.

When clinically appropriate direct change from monotherapy to the combinations may be considered:

- **IRBEL HCT 150 mg/12.5 mg tablets** may be administered in patients whose blood pressure is not adequately controlled with hydrochlorothiazide or irbesartan 150 mg alone;
- **IRBEL HCT 300 mg/12.5 mg tablets** may be administered in patients insufficiently controlled by irbesartan 300 mg or by **IRBEL HCT 150 mg/12.5 mg tablets**.
- **IRBEL HCT 300 mg/25 mg tablets** may be administered in patients insufficiently controlled by **IRBEL HCT 300 mg/12.5 mg tablets**.

Doses higher than 300 mg irbesartan/25 mg hydrochlorothiazide once daily are not recommended.

When necessary, **IRBEL HCT** tablets may be administered with another antihypertensive medicinal product.

Renal impairment: due to the hydrochlorothiazide component, **IRBEL HCT** tablets are not recommended for patients with severe renal dysfunction (creatinine clearance < 30 ml/min). Loop diuretics are preferred to thiazides in this population. No dosage adjustment is necessary in patients with renal impairment whose renal creatinine clearance is ≥ 30 ml/min.

Hepatic impairment: **IRBEL HCT** tablets are not indicated in patients with severe hepatic impairment. Thiazides should be used with caution in patients with impaired hepatic function. No dosage adjustment of **IRBEL HCT** tablets are necessary in patients with mild to moderate hepatic impairment.

Elderly patients: no dosage adjustment of **IRBEL HCT** tablets are necessary in elderly patients.

Pediatric patients: This product is not recommended for use in children and adolescents due to a lack of data on safety and efficacy of Irbesartan and HCTZ.

CONTRAINDICATIONS

- Patients with a history of hypersensitivity to the active substances, to any of the excipients, or to other sulfonamide-derived substances (hydrochlorothiazide is a sulfonamide-derived substance)
- Second and third trimesters of pregnancy
- Severe renal impairment (creatinine clearance < 30 ml/min)
- Refractory hypokalaemia, hypercalcaemia
- Severe hepatic impairment, biliary cirrhosis and cholestasis
- Lactation

WARNINGS AND PRECAUTIONS

Non-melanoma skin cancer: An increased risk of non-melanoma skin cancer (NMSC) [basal cell carcinoma (BCC) and squamous cell carcinoma (SCC)] with increasing cumulative dose of hydrochlorothiazide (HCTZ) exposure has been observed in two epidemiological studies based on the Danish National Cancer Registry. Photosensitizing actions of HCTZ could act as a possible mechanism for NMSC.

Patients taking HCTZ should be informed of the risk of NMSC and advised to regularly check their skin for any new lesions and promptly report any suspicious skin lesions. Possible preventive measures such as limited exposure to sunlight and UV rays and, in case of exposure, adequate protection should be advised to the patients in order to minimize the risk of skin cancer. Suspicious skin lesions should be promptly examined potentially including histological examinations of biopsies. The use of HCTZ may also need to be reconsidered in patients who have experienced previous NMSC.

Hypotension-Volume-depleted patients: Combination of irbesartan and hydrochlorothiazide has been rarely associated with symptomatic hypotension in hypertensive patients without other risk factors for hypotension. Symptomatic hypotension may be expected to occur in patients who are volume and/or sodium depleted by vigorous diuretic therapy, dietary salt restriction, diarrhoea or vomiting. Such conditions should be corrected before initiating therapy with combination of irbesartan and hydrochlorothiazide.

Renal artery stenosis-Renovascular hypertension: there is an increased risk of severe hypotension and renal insufficiency when patients with bilateral renal artery stenosis or stenosis of the artery to a single functioning kidney are treated with angiotensin converting enzyme inhibitors or angiotensin-II receptor antagonists. While this is not documented with combination of irbesartan and hydrochlorothiazide, a similar effect should be anticipated.

Renal impairment and kidney transplantation: when combination of irbesartan and hydrochlorothiazide is used in patients with impaired renal function, a periodic monitoring of potassium, creatinine and uric acid serum levels is recommended. No experience regarding the administration of combination of irbesartan and hydrochlorothiazide in patients with recent kidney transplantation has been reported.

Combination of irbesartan and hydrochlorothiazide should not be used in patients with severe renal impairment (creatinine clearance < 30 ml/min). Thiazide diuretic-associated azotemia may occur in patients with impaired renal function. No dosage adjustment is necessary in patients with renal impairment whose creatinine clearance is ≥ 30 ml/min. However, in patients with mild to moderate renal impairment (creatinine clearance ≥ 30 ml/min but < 60 ml/min) this combination should be administered with caution.

Hepatic impairment: thiazides should be used with caution in patients with impaired hepatic function or progressive liver disease, since minor alterations of fluid and electrolyte balance may precipitate hepatic coma. There is no clinical experience with combination of irbesartan and hydrochlorothiazide in patients with hepatic impairment.

Aortic and mitral valve stenosis, obstructive hypertrophic cardiomyopathy: as with other vasodilators, special caution is indicated in patients suffering from aortic or mitral stenosis, or obstructive hypertrophic cardiomyopathy.

Primary aldosteronism: patients with primary aldosteronism generally will not respond to anti-hypertensive medicinal products acting through inhibition of the renin-angiotensin system. Therefore, the use of combination of irbesartan and hydrochlorothiazide is not recommended.

Metabolic and endocrine effects: thiazide therapy may impair glucose tolerance. In diabetic patients dosage adjustments of insulin or oral hypoglycemic agents may be required. Latent diabetes mellitus may become manifest during thiazide therapy. Increases in cholesterol and triglyceride levels have been associated with thiazide diuretic therapy; however at the 12.5 mg dose contained in combination of irbesartan and hydrochlorothiazide minimal or no effects have been reported. Hyperuricaemia may occur or frank gout may be precipitated in certain patients receiving thiazide therapy.

Electrolyte imbalance: as for any patient receiving diuretic therapy, periodic determination of serum electrolytes should be performed at appropriate intervals. Thiazides, including hydrochlorothiazide, can cause fluid or electrolyte imbalance (hypokalaemia, hyponatraemia, and hypochloremic alkalosis). Warning signs of fluid or electrolyte imbalance are dryness of mouth, thirst, weakness, lethargy, drowsiness, restlessness, muscle pain or cramps, muscular fatigue, hypotension, oliguria, tachycardia, and gastrointestinal disturbances such as nausea or vomiting. Although hypokalaemia may develop with the use of thiazide diuretics, concurrent therapy with irbesartan may reduce diuretic-induced hypokalaemia. The risk of hypokalaemia is greatest in patients with cirrhosis of the liver, in patients experiencing brisk diuresis, in patients who are receiving inadequate oral intake of electrolytes and in patients receiving concomitant therapy with corticosteroids or ACTH. Conversely, due to the irbesartan component of combination of irbesartan and hydrochlorothiazide hyperkalaemia might occur, especially in the presence of renal impairment.

and/or heart failure, and diabetes mellitus. Adequate monitoring of serum potassium in patients at risk is recommended. Potassium-sparing diuretics, potassium supplements or potassium-containing salts substitutes should be co-administered cautiously with combination of irbesartan and hydrochlorothiazide.

Dilutional hyponatremia may occur in edematous patients in hot weather; appropriate therapy is water restriction, rather than administration of salt except in rare instances when the hyponatremia is life-threatening. In actual salt depletion, appropriate replacement is the therapy of choice. There is no reported evidence that irbesartan would reduce or prevent diuretic-induced hyponatraemia. Chloride deficit is generally mild and usually does not require treatment. Thiazides may decrease urinary calcium excretion and cause an intermittent and slight elevation of serum calcium in the absence of known disorders of calcium metabolism. Marked hypercalcaemia may be evidence of hidden hyperparathyroidism. Thiazides should be discontinued before carrying out tests for parathyroid function. Thiazides have been shown to increase the urinary excretion of magnesium, which may result in hypomagnesaemia.

Lithium: the combination of lithium and combination of irbesartan and hydrochlorothiazide is not recommended.

Anti-doping test: hydrochlorothiazide contained in this medicinal product could produce a positive analytic result in an anti-doping test.

General: in patients whose vascular tone and renal function depend predominantly on the activity of the renin-angiotensin-aldosterone system (e.g. patients with severe congestive heart failure or underlying renal disease, including renal artery stenosis), treatment with angiotensin converting enzyme inhibitors or angiotensin-II receptor antagonists that affect this system has been associated with acute hypotension, azotemia, oliguria, or rarely acute renal failure. As with any anti-hypertensive agent, excessive blood pressure decrease in patients with ischemic cardiopathy or ischemic cardiovascular disease could result in a myocardial infarction or stroke. Hypersensitivity reactions to hydrochlorothiazide may occur in patients with or without a history of allergy or bronchial asthma, but are more likely in patients with such a history. Exacerbation or activation of systemic lupus erythematosus has been reported with the use of thiazide diuretics. Cases of photosensitivity reactions have been reported with thiazides diuretics. If photosensitivity reaction occurs during treatment, it is recommended to stop the treatment. If a re-administration of the diuretic is deemed necessary, it is recommended to protect exposed areas to the sun or to artificial UVA.

Acute Myopia and Secondary Angle-Closure Glaucoma: Sulfonamide or sulfonamide derivative drugs, such as hydrochlorothiazide, can cause an idiosyncratic reaction, resulting in transient myopia and acute angle-closure glaucoma. Cases of acute angle-closure glaucoma have been reported with hydrochlorothiazide. Symptoms include acute onset of decreased visual acuity or ocular pain and typically occur within hours to weeks of drug initiation. Untreated acute angle-closure glaucoma can lead to permanent vision loss. The primary treatment is to discontinue drug intake as rapidly as possible. Prompt medical or surgical treatments may need to be considered if the intraocular pressure remains uncontrolled. Risk factors for developing acute angle-closure glaucoma may include a history of sulfonamide or penicillin allergy.

Pregnancy: Angiotensin II Receptor Antagonists (AIIAs) should not be initiated during pregnancy. Unless continued AIIAs therapy is considered essential, patients planning pregnancy should be changed to alternative anti-hypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with AIIAs should be stopped immediately, and, if appropriate, alternative therapy should be started. This product contains lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this product.

Acute Respiratory Toxicity: Very rare severe cases of acute respiratory toxicity, including acute respiratory distress syndrome (ARDS) have been reported after taking hydrochlorothiazide. Pulmonary oedema typically develops within minutes to hours after hydrochlorothiazide intake. At the onset, symptoms include dyspnoea, fever, pulmonary deterioration and hypotension. If diagnosis of ARDS is suspected, **IRBEL HCT** should be withdrawn and appropriate treatment given. Hydrochlorothiazide should not be administered to patients who previously experienced ARDS following hydrochlorothiazide intake.

Effect on ability to drive and use machine: No studies on the effect on the ability to drive and use machines have been performed.

DRUG INTERACTIONS

Other antihypertensive agents: the antihypertensive effect of combination of irbesartan and hydrochlorothiazide may be increased with the concomitant use of other antihypertensive agents. Irbesartan and hydrochlorothiazide (at doses up to 300 mg irbesartan/25 mg hydrochlorothiazide) have been safely administered with other antihypertensive agents including calcium channel blockers and beta-adrenergic blockers. Prior treatment with high dose diuretics may result in volume depletion and a risk of hypotension when initiating therapy with irbesartan with or without thiazide diuretics unless the volume depletion is corrected first.

Lithium: reversible increases in serum lithium concentrations and toxicity have been reported during concomitant administration of lithium with angiotensin converting enzyme inhibitors. Similar effects have been very rarely reported with irbesartan so far. Furthermore, renal clearance of lithium is reduced by thiazides so the risk of lithium toxicity could be increased with combination of irbesartan and hydrochlorothiazide. Therefore, the combination of lithium and combination of irbesartan and hydrochlorothiazide is not recommended. If the combination proves necessary, careful monitoring of serum lithium levels is recommended.

Medicinal products affecting potassium: the potassium-depleting effect of hydrochlorothiazide is attenuated by the potassium-sparing effect of irbesartan. However, this effect of hydrochlorothiazide on serum potassium is expected to be potentiated by other medicinal products associated with potassium loss and hypokalaemia (e.g. other kaliuretic diuretics, laxatives, amphotericin,

carbenoxolone, penicillin G sodium). Conversely, based on the experience with the use of other medicinal products that blunt the renin-angiotensin system, concomitant use of potassium-sparing diuretics, potassium supplements, salt substitutes containing potassium or other medicinal products that may increase serum potassium levels (e.g. heparin sodium) may lead to increases in serum potassium. Adequate monitoring of serum potassium in patients at risk is recommended.

Medicinal products affected by serum potassium disturbances: periodic monitoring of serum potassium is recommended when combination of irbesartan and hydrochlorothiazide is administered with medicinal products affected by serum potassium disturbances (e.g. digitalis glycosides, antiarrhythmics).

Non-steroidal anti-inflammatory drugs: when angiotensin II antagonists are administered simultaneously with non-steroidal anti-inflammatory drugs (i.e. selective COX-2 inhibitors, acetylsalicylic acid (> 3 g/day) and non-selective NSAIDs), attenuation of the antihypertensive effect may occur. As with ACE inhibitors, concomitant use of angiotensin II antagonists and NSAIDs may lead to an increased risk of worsening of renal function, including possible acute renal failure, and an increase in serum potassium, especially in patients with poor pre-existing renal function. The combination should be administered with caution, especially in the elderly. Patients should be adequately hydrated and consideration should be given to monitoring renal function after initiation of concomitant therapy, and periodically thereafter.

Additional information on irbesartan interactions: the pharmacokinetic of irbesartan is not affected by hydrochlorothiazide. Irbesartan is mainly metabolized by CYP2C9 and to a lesser extent by glucuronidation. No significant pharmacokinetic or pharmacodynamic interactions have been reported when irbesartan was coadministered with warfarin, a medicinal product metabolised by CYP2C9. The effects of CYP2C9 inducers such as rifampicin on the pharmacokinetic of irbesartan have not been reported. The pharmacokinetic of digoxin was not altered by co-administration of irbesartan.

Additional information on hydrochlorothiazide interactions: when administered concurrently, the following medicinal products may interact with thiazide diuretics:

Alcohol, Barbiturates and Narcotics: potentiation of orthostatic hypotension may occur

Antidiabetic medicinal products (oral agents and insulins): dosage adjustment of the antidiabetic medicinal product may be required

Colestyramine and Colestipol resins: absorption of hydrochlorothiazide is impaired in the presence of anionic exchange resins. Combination of irbesartan and hydrochlorothiazide should be taken at least one hour before or four hours after these medications

Corticosteroids, ACTH: electrolyte depletion, particularly hypokalaemia, may be increased

Digitalis glycosides: thiazide induced hypokalaemia or hypomagnesaemia favour the onset of digitalis-induced cardiac arrhythmias

Non-steroidal anti-inflammatory drugs: the administration of a non-steroidal anti-inflammatory drug may reduce the diuretic, natriuretic and antihypertensive effects of thiazide diuretics in some patients

Pressor amines (e.g. noradrenaline): the effect of pressor amines may be decreased, but not sufficiently to preclude their use

Nondepolarizing skeletal muscle relaxants (e.g. tubocurarine): the effect of nondepolarizing skeletal muscle relaxants may be potentiated by hydrochlorothiazide

Antigout medicinal products: dosage adjustments of antigout medicinal products may be necessary as hydrochlorothiazide may raise the level of serum uric acid. Increase in dosage of probenecid or sulfinpyrazone may be necessary. Co-administration of thiazide diuretics may increase the incidence of hypersensitivity reactions to allopurinol

Calcium salts: thiazide diuretics may increase serum calcium levels due to decreased excretion. If calcium supplements or calcium sparing medicinal products (e.g. vitamin D therapy) must be prescribed, serum calcium levels should be monitored and calcium dosage adjusted accordingly

Other interactions: the hyperglycaemic effect of beta-blockers and diazoxide may be enhanced by thiazides. Anticholinergic agents (e.g. atropine, beperiden) may increase the bioavailability of thiazide-type diuretics by decreasing gastrointestinal motility and stomach emptying rate. Thiazides may increase the risk of adverse effects caused by amantadine. Thiazides may reduce the renal excretion of cytotoxic medicinal products (e.g. cyclophosphamide, methotrexate) and potentiate their myelosuppressive effects.

SIDE EFFECTS/ ADVERSE REACTIONS

Combination of irbesartan and hydrochlorothiazide

Adverse reactions reported in hypertensive patients receiving various doses (including spontaneous reports) are presented below.

Table: Adverse Reactions reported in hypertensive patients receiving various doses including Spontaneous Reports*		
Investigations	Common	increases in blood urea nitrogen (BUN), creatinine and creatine kinase
	Uncommon	decreases in serum potassium and sodium
Cardiac disorders	Uncommon	syncope, hypotension, tachycardia, oedema
Nervous system disorders	Common	dizziness
	Uncommon	orthostatic dizziness
	Not known	headache

Table: Adverse Reactions reported in hypertensive patients receiving various doses including Spontaneous Reports*		
Ear and labyrinth disorders	Not known	tinnitus
Respiratory, thoracic and mediastinal disorders	Not known	cough
Gastrointestinal disorders	Common	nausea/vomiting
	Uncommon	diarrhoea
	Not known	dyspepsia, dysgeusia
Renal and urinary disorders	Common	abnormal urination
	Not known	impaired renal function including isolated cases of renal failure in patients at risk
Musculoskeletal and connective tissue disorders	Uncommon	swelling extremity
	Not known	arthralgia, myalgia
Metabolism and nutrition disorders	Not known	hyperkalaemia
Vascular disorders	Uncommon	flushing
General disorders and administration site conditions	Common	fatigue
Immune system disorders	Not known	cases of hypersensitivity reactions such as angioedema, rash, urticaria
Hepatobiliary disorders	Uncommon	jaundice
	Not known	hepatitis, abnormal liver function
Reproductive system and breast disorders	Uncommon	sexual dysfunction, libido changes
* Frequency for adverse reactions detected by spontaneous reports is described as "not known"		

Other reported adverse events with combination of irbesartan and hydrochlorothiazide regardless of drug relationship are listed below:

Influenza, abdominal pain, sinus abnormality, upper respiratory tract infections (URI), pharyngitis, rhinitis, urinary tract infection, anxiety/nervousness, muscle cramps, chest pain, musculoskeletal pain and hypokalemia.

Additional information on individual components

in addition to the adverse reactions listed above for the combination of irbesartan and hydrochlorothiazide, other adverse reactions previously reported with one of the individual components may be potential adverse reactions with combination of irbesartan and hydrochlorothiazide. Tables below provide adverse reactions reported with the individual components of irbesartan and hydrochlorothiazide.

Adverse reactions reported with Irbesartan alone without regards to causality are listed below:

Fever, chills, facial oedema, upper extremity oedema, hypertension, cardiac murmur, myocardial infarction, angina pectoris, arrhythmic/conduction disorder, cardiorespiratory arrest, heart failure, hypertension crisis, pruritis, dermatitis, ecchymosis, erythema face, gout, constipation, gastroenteritis, flatulence, abdominal distention, musculoskeletal trauma, muscle cramp, arthritis, muscle ache, musculoskeletal chest pain, joint stiffness, bursitis, muscle weakness, sleep disturbance, numbness,

somnolence, vertigo, emotional disturbances, depression, paresthesia, tremor, transient ischemic attack, cerebrovascular accidents, prostate disorder, epistaxis, tracheobronchitis, congestion, pulmonary congestion, dyspnoea, wheezing, vision disturbance, hearing abnormality, ear infection, ear pain, conjunctivitis.

Rare cases of rhabdomyolysis have been reported in patients receiving angiotensin II receptor blockers.

Adverse reactions reported with hydrochlorothiazide alone

Table: Adverse reactions (regardless of relationship to medicinal product) reported with the use of <i>hydrochlorothiazide</i> alone		
Investigations	Not known	electrolyte imbalance (including hyponatraemia), hyperuricaemia, glycosuria, hyperglycaemia, increases in cholesterol and triglycerides
Cardiac disorders	Not known	cardiac arrhythmias
Blood and lymphatic system disorders	Not known	aplastic anaemia, bone marrow depression, neutropenia/agranulocytosis, haemolytic anaemia, leucopenia, thrombocytopenia
Nervous system disorders	Not known	vertigo, paraesthesia, light-headedness, restlessness
Eye disorders	Not known	transient blurred vision, xanthopsia
Respiratory, thoracic and mediastinal disorders	Very rare	Acute respiratory distress syndrome (ARDS)
Gastrointestinal disorders	Not known	pancreatitis, anorexia, diarrhoea, constipation, gastric irritation, sialadenitis, loss of appetite
Renal and urinary disorders	Not known	interstitial nephritis, renal dysfunction
Skin and subcutaneous tissue disorders	Not known	anaphylactic reactions, toxic epidermal necrolysis, necrotizing angitis (vasculitis, cutaneous vasculitis), cutaneous lupus erythematosus-like reactions, reactivation of cutaneous lupus erythematosus, photosensitivity reactions, rash, urticaria
Musculoskeletal and connective tissue disorders	Not known	weakness, muscle spasm
Vascular disorders	Not known	postural hypotension
General disorders and administration site conditions	Not known	fever
Hepatobiliary disorders	Not known	jaundice (intrahepatic cholestatic jaundice)
Psychiatric disorders	Not known	depression, sleep disturbances
Neoplasms benign, malignant and unspecified (incl. cysts and polyps)	Not known	Non-melanoma skin cancer (Basal cell carcinoma and Squamous cell carcinoma) <i>Description of selected adverse reactions</i> <i>Non-melanoma skin cancer:</i> Based on available data from epidemiological studies, cumulative dose- dependent association between HCTZ and NMSC has been observed.

The dose dependent adverse events of hydrochlorothiazide (particularly electrolyte disturbances) may increase when titrating the hydrochlorothiazide.

Other adverse events reported with hydrochlorothiazide alone without regards to causality are listed below:

Cramping, purpura, erythema multiforme including Stevens-Johnson syndrome, exfoliative dermatitis.

USE IN SPECIAL POPULATION

- **Pregnancy**

Reported epidemiological evidence regarding the risk of teratogenicity following exposure to ACE inhibitors during the first trimester of pregnancy is not conclusive; however a small increase in risk cannot be excluded. Whilst there is no controlled reported epidemiological information on the risk with Angiotensin II Receptor Antagonists (AIIAs), similar risks may exist for this class of drugs. Unless continued AIIAs therapy is considered essential, patients planning pregnancy should be changed to alternative anti-hypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with AIIAs should be stopped immediately, and, if appropriate, alternative therapy should be started.

Exposure to AIIAs therapy during the second and third trimesters may induce human fetotoxicity (decreased renal function, oligohydramnios, skull ossification retardation) and neonatal toxicity (renal failure, hypotension, hyperkalaemia).

Should exposure to AIIAs have occurred from the second trimester of pregnancy, ultrasound check of renal function and skull is recommended. Infants whose mothers have taken AIIAs should be closely observed for hypotension.

Thiazides cross the placental barrier and appear in cord blood. They may cause a decrease in placental perfusion, foetal electrolyte disturbances and possibly other reactions that have occurred in the adults. Cases of neonatal thrombocytopenia, or foetal or neonatal jaundice have been reported with maternal thiazide therapy. Since combination of irbesartan and hydrochlorothiazide contains hydrochlorothiazide, it is not recommended during the first trimester of pregnancy. A switch to a suitable alternative treatment should be carried out in advance of a planned pregnancy.

- **Lactation**

Combination of irbesartan and hydrochlorothiazide is not recommended during lactation and alternative treatments with better established safety profiles during breast-feeding are preferable, especially while nursing a newborn or preterm infant.

OVERDOSE

No specific information regarding treatment of overdose with combination of irbesartan and hydrochlorothiazide has been reported. The patient should be closely monitored, and the treatment

should be symptomatic and supportive. Management depends on the time since ingestion and the severity of the symptoms. Suggested measures include induction of emesis and/or gastric lavage. Activated charcoal may be useful in the treatment of overdose. Serum electrolytes and creatinine should be monitored frequently. If hypotension occurs, the patient should be placed in a supine position, with salt and volume replacements given quickly.

The most likely manifestations of irbesartan overdose are expected to be hypotension and tachycardia; bradycardia might also occur.

Overdose with hydrochlorothiazide is associated with electrolyte depletion (hypokalaemia, hyponatraemia, hyponatremia) and dehydration resulting from excessive diuresis. The most common signs and symptoms of overdose are nausea and somnolence. Hypokalaemia may result in muscle spasms and/or accentuate cardiac arrhythmias associated with the concomitant use of digitalis glycosides or certain anti-arrhythmic medicinal products.

Irbesartan is not removed by haemodialysis. The degree to which hydrochlorothiazide is removed by haemodialysis has not been reported.

Storage: Store below 30°C

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN

Supply: Blister pack of 4 x 7's

Shelf life: 24 Months

Manufacturer:

SUN PHARMACEUTICAL INDUSTRIES LIMITED

Paonta Sahib, Dist Sirmour

Himachal Pradesh-173025

Product Registration Holder:

RANBAXY (MALAYSIA) SDN, BHD

Lot 23, Bakar Arang Industrial Estate,

08000 Sungai Petani, Kedah, MALAYSIA

Date of revision: August 2024