

Please refer enclosed package insert for more information.
Contains Chlorocresol 0.1% w/w as preservative.
Keep container well closed. Store below 30°C
Protect from light.

Product Registration Holder & Manufactured By :
 **Kotra Pharma (M) Sdn. Bhd.**
No. 1, 2 & 3, Jalan TTC 12, (90082V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

FOR EXTERNAL USE ONLY
UNTUK KEGUNAAN LUAR SAHAJA
KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK

Controlled
Medicine





MICONAZOLE
CREAM

5 Gm

Miconazole Nitrate (micronised) 2% w/w

5 Gm


MICONAZOLE
CREAM

INDICATIONS :

Used topically in the management of superficial candidiasis and fungal skin infection such as mucocutaneous candidiasis, dermatophytosis and pityriasis versicolor.

DIRECTION FOR USE :

Apply cream (according to the size of lesion) to the lesion twice daily. Rub gently until the cream has been completely absorbed by the skin. The treatment should be continued without interruption until the lesion has completely disappeared (usually after 2 to 5 weeks). Continue treatment for at least 1 week after the lesion has completely disappeared to prevent relapse.

MAL19900255AZ SIN10967P
BRU08090129P TZ15H0280
Lao Reg. No. 1113968/13
Zimbabwe Reg No. 2018/14.1.3/5751 PIM

REGN - 010324

120 x 30 x 25

Batch No.:
Exp Date:
Mfg Date:



MICONAZOLE
CREAM

5 Gm

Miconazole Nitrate (micronised) 2% w/w



Please refer enclosed package insert for more information.
Contains Chlorocresol 0.1% w/w as preservative.
Keep container well closed. Store below 30°C
Protect from light.

Product Registration Holder & Manufactured By :
 **Kotra Pharma (M) Sdn. Bhd.**
No. 1, 2 & 3, Jalan TTC 12, (90082V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

FOR EXTERNAL USE ONLY
UNTUK KEGUNAAN LUAR SAHAJA
KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK

Controlled
Medicine





MICONAZOLE
CREAM

15 Gm

Miconazole Nitrate (micronised) 2% w/w

15 Gm



MICONAZOLE
CREAM

INDICATIONS :

Used topically in the management of superficial candidiasis and fungal skin infection such as mucocutaneous candidiasis, dermatophytosis and pityriasis versicolor.

DIRECTION FOR USE :

Apply cream (according to the size of lesion) to the lesion twice daily. Rub gently until the cream has been completely absorbed by the skin. The treatment should be continued without interruption until the lesion has completely disappeared (usually after 2 to 5 weeks). Continue treatment for at least 1 week after the lesion has completely disappeared to prevent relapse.

MAL19900255AZ SIN10967P
BRU08090129P TZ15H0280
Lao Reg. No. 1113968/13
Zimbabwe Reg No. 2018/14.1.3/5751 PIM

REGN - 010324

120 x 30 x 25



MICONAZOLE
CREAM

15 Gm

Miconazole Nitrate (micronised) 2% w/w

Batch No.:
Exp Date:
Mfg Date:

