

SUSPENSION
CEFACLOR-125
Axxcel®

Axxcel®
CEFACLOR-125

SUSPENSION

Each 5ml contains:
Cefaclor Monohydrate
equivalent to
Cefaclor 125mg

60ml

60ml when reconstituted.

For oral administration only.

PRECAUTION:

Should be used with caution in patients with penicillin sensitivity.

Keep container well closed. Store below 30°C until reconstituted. After reconstitution, refrigerate between 2°C to 8°C and use within 7 days.

Please refer to enclosed package insert for more information.

KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK

Controlled
Medicine | SHAKE WELL BEFORE USE
GONCANG SEBELUM DIGUNAKAN

MAL20014012AZ
MYN R1903A6687
CAM R₃ 6791 IP-09
BRU14081447P

 **Kotra Pharma (M) Sdn. Bhd.**
No. 1, 2 & 3, Jalan TTC 12, (90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

Axxcel®
CEFACLOR-125

SUSPENSION

Each 5ml contains:
Cefaclor Monohydrate
equivalent to
Cefaclor 125mg

60ml

HOW TO PREPARE YOUR MEDICINE




TO COMPLETE THE COURSE OF MEDICINE



50 x 62 x 120

Batch No.
Mfg Date
Exp Date

SUSPENSION CEFACLOR-125 Axcel®			
Axcel® CEFACLOR-125 SUSPENSION Each 5ml contains: Cefaclor Monohydrate equivalent to Cefaclor 125mg	120ml when reconstituted.	Axcel® 	HOW TO PREPARE YOUR MEDICINE 1  SHAKE BOTTLE WELL TO LOOSEN GRANULES 2  FIRST REMOVE CAP AND DOSE-MASTER™ ADAPTER. THEN, ADD SOME WATER TO MIX WITH THE GRANULES. AFTER THAT, SHAKE AND TOP UP TO 120ML MARK 3  PLACE THE DOSE-MASTER™ ADAPTER AND CAP 4  SHAKE THE BOTTLE WELL TO MIX THE MEDICINE PROPERLY 5  PLACE THE DOSE-MASTER™ 6  WITHDRAW THE MEDICINE WITH DOSE-MASTER™ FOR ACCURATE DOSE TO COMPLETE THE COURSE OF MEDICINE  Barcode No.
	For oral administration only.	CEFACLOR-125	
	PRECAUTION: Should be used with caution in patients with penicillin sensitivity.	SUSPENSION	
	Keep container well closed. Store below 30°C until reconstituted. After reconstitution, refrigerate between 2°C to 8°C and use within 7 days.	Each 5ml contains: Cefaclor Monohydrate equivalent to Cefaclor 125mg	
	Please refer to enclosed package insert for more information.		
KEEP OUT OF REACH OF CHILDREN JAUHI DARI KANAK-KANAK			
Controlled Medicine SHAKE WELL BEFORE USE GONCANG SEBELUM DIGUNAKAN			
MAL20014012AZ MYN R1903A6687 CAM R: 6791 IP-09 BRU14081447P			
KOTRA PHARMA Kotra Pharma (M) Sdn. Bhd. No. 1, 2 & 3, Jalan TTC 12, (90082-V) Cheng Industrial Estate, 75250 Melaka, Malaysia.			
120ml	120ml		
	Batch No. Mfg Date Exp Date		
	50 x 62 x 120		