

Please refer enclosed package insert for more information. Keep container well closed. Store below 30°C. Protect from light.



Kotra Pharma (M) Sdn. Bhd.
No. 1, 2 & 3, Jalan TTC 12, (90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

FOR EXTERNAL USE ONLY
UNTUK KEGUNAAN LUAR SAHAJA
KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK

Controlled
Medicine



Axcel®

FUSIDIC ACID
OINTMENT

5 Gm

Sodium Fusidate (micronised) 2% w/w

5 Gm

Axcel®
FUSIDIC ACID
OINTMENT

INDICATIONS :

Skin infection caused by *Staphylococci*, *Streptococci* and other susceptible or sodium fusidate sensitive organisms.

DIRECTION FOR USE :

Apply a thin layer of the ointment to the affected area 2-3 times a day, generally for a period of seven days.

MAL20021410AZ
HK-60868
CAM R,0299IP-09
BRU08090127P
MYN R2001AA2327
VN-12967-11
11 I 3963/13

90 x 27 x 20

Axcel®

FUSIDIC ACID
OINTMENT

5 Gm

Sodium Fusidate (micronised) 2% w/w

Batch no. :
Mfg Date :
Exp Date :



Pantone 192



Black



Pantone 2738C



Pantone 555C

Please refer enclosed package insert for more information. Keep container well closed. Store below 30°C. Protect from light.



Kotra Pharma (M) Sdn. Bhd.
No.1, 2 & 3, Jalan TTC 12, (90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

FOR EXTERNAL USE ONLY
UNTUK KEGUNAAN LUAR SAHAJA

KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK

Controlled
Medicine



9 556586 201146

Axcel[®]

FUSIDIC ACID
OINTMENT

15 Gm

Sodium Fusidate (micronised) 2% w/w

15 Gm

Axcel[®]
FUSIDIC ACID
OINTMENT

INDICATIONS:

Skin infection caused by *Staphylococci*, *Streptococci* and other susceptible or sodium fusidate sensitive organisms.

DIRECTION FOR USE:

Apply a thin layer of the ointment to the affected area 2-3 times a day, generally for a period of seven days.

MAL20021410AZ
HK-60868
CAM R,0298IP-09
BRU08090127P
MYN R2001AA2327
VN-12967-11
11 I 3963/13

120 x 30 x 25

Axcel[®]

FUSIDIC ACID
OINTMENT

15 Gm

Sodium Fusidate (micronised) 2% w/w

Batch no. :
Mfg Date :
Exp Date :

