

			SUSPENSION ERYTHROMYCIN ES-400 
HOW TO PREPARE YOUR MEDICINE 1  SHAKE BOTTLE WELL TO LOOSEN GRANULES 2  FIRST REMOVE CAP AND DOSE-MASTER™ ADAPTER. THEN, ADD SOME WATER TO MIX WITH THE GRANULES. AFTER THAT, SHAKE AND TOP UP TO 60ml MARK 3  PLACE THE DOSE-MASTER™ ADAPTER AND CAP 4  SHAKE THE BOTTLE WELL TO MIX THE MEDICINE PROPERLY 5  PLACE THE DOSE-MASTER™ 6  WITHDRAW THE MEDICINE WITH DOSE-MASTER™ FOR ACCURATE DOSE TO COMPLETE THE COURSE OF MEDICINE  9 556586 503066	 ERYTHROMYCIN ES-400 SUSPENSION Each 5ml contains: Erythromycin Ethylsuccinate (micronised) equivalent to Erythromycin 400mg 60ml	DOSAGE: Adults: 400mg (5ml) 6 hourly or 800mg (10ml) 12 hourly up to 4g daily. Children 2 to 8 years: 1g (12.5ml) daily in divided dose. Infant and children up to 2 years: 500mg (6.25ml) daily in divided dose. 60ml when reconstituted. Keep container well closed. Store at room temperature not exceeding 30°C until reconstitution. After reconstitution, store below 25°C and use within 35 days, refrigeration is not required. For oral administration. Please refer to enclosed package insert for more information. KEEP OUT OF REACH OF CHILDREN JAUHI DARI KANAK-KANAK Controlled Medicine SHAKE WELL BEFORE USE GONCANG SEBELUM DIGUNAKAN MAL09082800AZ CAM R,0022IP-14 Manufactured By & Product Registration Holder :  Kotra Pharma (M) Sdn. Bhd. No. 1, 2 & 3, Jalan TTC 12, (06002-V) Cheng Industrial Estate, 75250 Melaka, Malaysia.	 ERYTHROMYCIN ES-400 SUSPENSION Each 5ml contains: Erythromycin Ethylsuccinate (micronised) equivalent to Erythromycin 400mg 60ml
		REGN - 210819	

			SUSPENSION ERYTHROMYCIN ES-400 
HOW TO PREPARE YOUR MEDICINE 1  SHAKE BOTTLE WELL TO LOOSEN GRANULES 2  PREPARE CAP AND DOSE-MASTER™ ADAPTER, THEN ADD SOME WATER TO MIX WITH THE GRANULES. AFTER THAT SHAKE AND TOP UP TO 100ML MARK 3  PLACE THE DOSE-MASTER™ ADAPTER AND CAP 4  SHAKE THE BOTTLE WELL TO MIX THE MEDICINE PROPERLY 5  PLACE THE DOSE-MASTER™ 6  WITHDRAW THE MEDICINE WITH DOSE-MASTER™ FOR ACCURATE DOSE TO COMPLETE THE COURSE OF MEDICINE  9 556586 503066	 ERYTHROMYCIN ES-400 SUSPENSION Each 5ml contains: Erythromycin Ethylsuccinate (micronised) equivalent to Erythromycin 400mg 100ml	DOSAGE: Adults: 400mg (5ml) 6 hourly or 800mg (10ml) 12 hourly up to 4g daily. Children 2 to 8 years: 1g (12.5ml) daily in divided dose. Infant and children up to 2 years: 500mg (6.25ml) daily in divided dose. 100ml when reconstituted. Keep container well closed. Store at room temperature not exceeding 30°C until reconstitution. After reconstitution, store below 25°C and use within 35 days, refrigeration is not required. For oral administration. Please refer to enclosed package insert for more information. KEEP OUT OF REACH OF CHILDREN JAUHI DARI KANAK-KANAK Controlled Medicine SHAKE WELL BEFORE USE GONCANG SEBELUM DIGUNAKAN MAL09082800AZ CAM R,0022IP-14 Manufactured By & Product Registration Holder :  Kotra Pharma (M) Sdn. Bhd. No. 1, 2 & 3, Jalan TTC 12, (09002-V) Cheng Industrial Estate, 75250 Melaka, Malaysia.	 ERYTHROMYCIN ES-400 SUSPENSION Each 5ml contains: Erythromycin Ethylsuccinate (micronised) equivalent to Erythromycin 400mg 100ml
		REGN - 210819	