AXCEL PAPAIN TABLET

Papain Pure (equivalent to proteolytic enzymes from Carica papaya 10,000 units)

What is in this leaflet

1. What Axcel Papain Tablet is used for.
2. Before you use Axcel Papain Tablet.
3. How to take Axcel Papain Tablet.
4. While you are using it.
5. Side effects.
6. Storage and disposal of Axcel Papain Tablet.
7. Product description.
8. Manufacturer and Product Registration Holder.
9. Date of revision.

What Axcel Papain Tablet is used for

Axcel Papain Tablet is use in conjunction with other physical or chemotherapeutic measures for treatment of oedema and inflammation.

Before you use Axcel Papain Tablet

-When you must not use it

Do not take Axcel Papain Tablet if:

- You have blood clotting disease.
- You have any generalised or systemic infections.

-Before you start to use it

Tell your doctor if:

- You have renal disease.
- You are suffering hepatic disease.
- You are pregnant.

-Taking other medicines

You should tell your doctor or pharmacist if you are taking the following medicine as it may interact with your Axcel Papain Tablet:

- Any anticoagulant drugs.

It may still be all right for you to take Axcel Papain Tablet and your doctor will be able to decide what is suitable for you.

How to use Axcel Papain Tablet

-How much to use

For oral administration only.

For adults: 2 tablets 4 times daily one day before surgery followed by 4 times daily for about 5 days.

-When to use it

Axcel Papain Tablet needs to be taken when you have oedema and inflammation. Usually, it is taken with the combination of other physical or chemotherapeutic measure.

-How long to use it

Take this medicine as stated in the leaflet without exceeding the dosage.

Follow any recommendations and advices from your doctor or pharmacist.

-If you forget to use it

Do not double up the dosage as Axcel Papain Tablet may give you unacceptable side effects.

-If you use too much (overdose)

It is advisable for you to contact and inform your doctor if you did take Axcel Papain Tablet too much.

While you are using it

-Things you must do

- Discuss with your doctor or pharmacist the dos' and don'ts' while taking this medicine.
- Tell your doctor if unexpectedly you need to take any other medicines.
- Take the Axcel Papain Tablet according to the recommended dosage.
- Tell your doctor or pharmacist if you experience any side effects that are not listed in this leaflet.

-Things you must not do

- Do not take your medicine more than the suggested frequency in a day.
- Do not over dosage the medicine during any single consumption.

-Things to be careful of

- Avoid prolonged use with high dosage.
- Use Axcel Papain Tablet with caution during pregnancy.
- Caution is needed while taking Axcel Papain Tablet for patients with renal or hepatic disease.

Side effects

Like all medicines, Axcel Papain Tablet can cause side effects although not everybody gets them. You are advised to consult doctor/ pharmacist for further information. The following side effects have been reported:

- Urticaria.
- Pruritus.
- Dizziness.
- Nausea.
- Vomiting.
- Diarrhoea.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835490, or visiting the website npra.moh.gov.my (Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

Storage and disposal of Axcel Papain Tablet

-Storage

Keep Axcel Papain Tablet out of reach of children.

Store below 25ºC. Keep container well closed and protect from light.

AXCEL PAPAIN TABLET

Papain Pure (equivalent to proteolytic enzymes from Carica papaya 10,000 units)

-Disposal

Do not use Axcel Papain Tablet after the expiry date which is stated on the carton.

Product description

-What it looks like

Axcel Papain Tablet is a light blue, biconvex, round tablets, 8mm in diameter with a single score- line on one side and marked KOTRA logo on the other side.

-Ingredients

- Active Ingredient (s):
  Papain

- Inactive Ingredients:
  Magnesium Stearate, Lactose, Sodium Starch Glycollate, PVP K90, Brilliant Blue Dye, Talcum Powder, Purified Water and Microcrystalline Cellulose.

-MAL number

MAL

Manufacturer and product registration holder

Kotra Pharma (M) Sdn Bhd
No. 1, 2 & 3, Jalan TTC 12,
Cheng Industrial Estate,
75250 Melaka, Malaysia.

Date of revision

27/03/19

Serial number

NPRA(R2)19/57