

AXCEL PHENTERMINE-30MG TABLETS

Phentermine (30mg)

What is in this leaflet

1. What Axcel Phentermine-30mg Tablets is used for.
2. How Axcel Phentermine-30mg Tablets works.
3. Before you use Axcel Phentermine-30mg Tablets.
4. How to take Axcel Phentermine-30mg Tablets.
5. While you are using it.
6. Side effects.
7. Storage and disposal of Axcel Phentermine-30mg Tablets.
8. Product description.
9. Manufacturer and Product Registration Holder.
10. Date of Revision.

What Axcel Phentermine-30mg Tablets is used for

Axcel Phentermine-30mg Tablets is indicated as adjunctive therapy to diet, in patients with obesity and a body mass index (BMI) of ≥ 30 , who have not responded to an appropriate weight-reducing regime alone.

How Axcel Phentermine-30mg Tablets Works

Phentermine Hydrochloride is used to suppress appetite in order to reduce weight in patients with obesity.

Before you use Axcel Phentermine-30mg Tablets

- When you must not use it

Do not take Axcel Phentermine-30mg Tablets if:

- You have hypersensitivity to sympathomimetic amines.
- You have current or past medical history of cardiovascular disease.
- You have hyperthyroidism
- You are currently or have recently taken MAO Inhibitors (MAOI) within 14 days
- You have glaucoma (eye disease which results in damage to the optic nerve and vision loss).
- You are in agitated states
- You are below 16 years old or elderly.
- You have past history of drug abuse.
- You are pregnant or nursing mother.

- Before you start to use it

Tell your doctor if:

- You are using any other drugs.
- You are pregnant or breastfeeding.
- You have hypertension or diabetes mellitus.
- You have kidney disease.

- Taking other medicines

You should tell your doctor if you are taking any other medicines, including those that you get without a prescription from pharmacy, supermarket or health food shop. Axcel Phentermine-30mg Tablets interact with these medicines when taken together:

- Monoamine Oxidase Inhibitors
- Alcohol
- Insulin and Oral Hypoglycemic Medications
- Adrenergic Neuron Blocking Drugs

Your doctor or pharmacist may have more information on medicines to be careful with or avoid while taking this medication.

How to use Axcel Phentermine-30mg Tablets

- How much to use

For oral administration only.

Dosage should be individualized to obtain an adequate response with the lowest effective dose.

The usual adult dose is one tablet daily, administered before breakfast or 1-2 hours after breakfast.

- When to use it

Your doctor will decide when it is appropriate for you to take Axcel Phentermine-30mg Tablets according to the type of condition you have.

- How long to use it

- Take this medicine as stated in the leaflet without exceeding the dosage.
- Follow any recommendations and advices from your doctor or pharmacist.

- If you forget to use it

- Do not double up the dosage as Axcel Phentermine-30mg Tablets may give you unacceptable side effects.

- If you use too much (overdose)

You must inform your doctor immediately if you have taken too much Axcel Phentermine-30mg Tablets.

While you are using it

- Things you must do

- Discuss with your doctor or pharmacist the dos' and don'ts' while taking this medicine.
- Tell your doctor if unexpectedly you need to take any other medicines.
- Take the Axcel Phentermine-30mg Tablets according to the recommended dosage.
- Tell your doctor or pharmacist if you experience any side effects that are not listed in this leaflet.

- Things you must not do

- Do not take your medicine more than the suggested frequency in a day.
- Do not over dosage the medicine during any single consumption.

- Things to be careful of

- Avoid prolonged use with high dosage.
- Caution during operating machinery or driving after taking the medicine.
- Avoid concomitant use of alcohol.

Side effects

Like all medicines, Axcel Phentermine-30mg Tablets can cause side effects although not everybody gets them. You are advised to consult doctor/ pharmacist for further information. The following side effects have been reported:

- High blood pressure
- Leaking heart valve (a condition where heart can't pump enough blood to meet body's needs)
- Palpitation (fast and irregular heartbeat)

AXCEL PHENTERMINE-30MG TABLETS

Phentermine (30mg)

- Tachycardia (abnormally rapid heart rate)
- Ischemic events (restriction in blood supply to body tissues)
- Overstimulation
- Restlessness (feeling uneasy)
- Dizziness
- Insomnia (sleep disorder)
- Euphoria (feeling of intense excitement)
- Dysphoria (discomfort or distress)
- Tremor (unintentional, rhythmic muscle movement)
- Headache
- Psychosis (mental disorder)
- Dryness of the mouth
- Bad breath
- Diarrhea
- Constipation
- Other gastrointestinal disturbances
- Urticaria (hives/ rash)
- Erectile dysfunction
- Changes in sexual desire

- Ingredients

Active Ingredient (s):

Phentermine Hydrochloride

Inactive Ingredients:

Microcrystalline Cellulose, Lactose Monohydrate, Corn Starch, FD&C Blue, Colloidal Silicon Dioxide and Magnesium Stearate.

- MAL Number

MAL16105033AZ

Manufacturer and Product Registration Holder

KOTRA PHARMA (M) SDN BHD
No. 1, 2 & 3, Jalan TTC 12,
Cheng Industrial Estate,
75250 Melaka, Malaysia.

Date of Revision

27/07/2023

Serial Number

NPRA(R2/1)230619/1086

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website nptra.gov.my [Consumers -> Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Axcel Phentermine-30mg Tablets

- Storage

Keep Axcel Phentermine-30mg Tablets out of reach of children. Store below 30°C. Protect from light.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Axcel Phentermine-30mg Tablets is a light blue colored, biconvex, oval shape, uncoated tablet embossed with kotra logo on one side and score line on other side.