

TRANSAMIN[®] CAPSULE

Tranexamic acid 250mg

What is in this leaflet

1. What Transamin Capsule is used for
2. How Transamin Capsule works
3. Before you use Transamin[®] Capsule
4. How to use Transamin Capsule
5. While you are using it
6. Side effects
7. Storage and Disposal of Transamin Capsule
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Transamin Capsule is used for

Transamin is an anti-fibrinolytic agent used for treating certain abnormal bleeding conditions.

How Transamin Capsule works

Transamin Capsule contains an active ingredient called tranexamic acid. It is used to stop or reduce unwanted bleeding. When you bleed your body forms clots to stop the bleeding. In some people the break down of clotting mechanism causing too much bleeding. Transamin stops the clots from dissolving and so reduces unwanted bleeding.

Before you take Transamin Capsule

When you must not take it

If you are receiving thrombin, an agent used in helping blood clot.

Pregnancy and lactation

If you are pregnant, trying to become pregnant or breast feeding ask your doctor or pharmacist for advice before taking Transamin Capsule

Before you start to take it

It is important to let your doctor know, if :

- You are allergic to the product before
- You have serious problems with your kidneys (kidney failure)
- You have or at risk with blood clot in your blood vessels (such as stroke and heart attack due to thrombosis)
- You have recently undergone a surgical operation
- You are at risk of excessive bleeding as a result of a bleeding disorder called consumption coagulopathy

Taking other medicines

Transamin should be administered with care when co-administered with the following medications: haemocoagulase (a medication used to treat bleeding.), batroxobin(used to prevent bleeding) and coagulation factor agents (clotting factor agents).

Inform your doctor or pharmacist that you are taking Transamin Capsule before your are given any medicine. Ask your doctor or pharmacist before you consume any health supplement, as they will have a more complete and detail list of medicines that may interact with Transamin Capsule if taken together

How to take Transamin Capsule

How much to take

3 to 8 capsules to be divided into 3 or 4 times per day. The dosage should be adjusted according to indication and patient's age.

When to take it

Each capsule should be swallowed whole after meal.

How long to take it

As prescribed by your doctor.

If you forget to take it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

If you take too much(overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

Things you must do

You should take Transamin Capsule as per prescribed by your doctor.

Things you must not do

Do not start a new medicine without first consulting a doctor.

Things to be careful of

Driving and using machines

If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

Side Effects

Transamin Capsule is generally well tolerated. The main adverse events were anorexia, nausea, vomiting, heartburn, itching, and skin rash. Incidence of convulsion was reported in dialysis patients.

TRANSAMIN[®] CAPSULE

Tranexamic acid 250mg

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835490, or visiting the website npra.gov.my [Consumers ♦ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Product Registration Holder

First Pharmaceutical Sdn Bhd
No. 20, Jalan SS 19/5
47500 Subang Jaya
Selangor, Malaysia

Date of Revision

11/02/2019

Serial Number

NPRA (R1/1) 28012019/033

Storage and Disposal of Transamin Capsule

Storage

Store below 30°C in well-closed containers.

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

What it looks like

Orange and White opaque hard capsule.

Ingredients

Active ingredient: Tranexamic acid
Inactive ingredients: Corn starch, Magnesium stearate

MAL No. 19890202ACZ

Manufacturer

CMIC CMO Co., Ltd., Shizuoka
Plant
1-588, Kanaya-Azuma, Shimada,
Shizuoka, Japan

For DAIICHI SANKYO CO., LTD.
3-5-1, Nihonbashi Honcho, Chuo-ku,
Tokyo, Japan