

Acyclovir 800 STADA

Acyclovir 800 mg

What is in this leaflet

1. What Acyclovir 800 STADA is used for
2. How Acyclovir 800 STADA works
3. Before you take Acyclovir 800 STADA
4. How to take Acyclovir 800 STADA
5. While you are using it
6. Possible side effects
7. Storage and Disposal of Acyclovir 800 STADA
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Acyclovir 800 STADA is used for

Acyclovir 800 STADA is a medicine for the treatment of viral infections (antiviral agent). Acyclovir is used for Shingles (herpes zoster).

How Acyclovir 800 STADA works

Acyclovir is an antiviral medicine, which acts on infected cells by stopping the virus growing.

Before you use Acyclovir 800 STADA

When you must not take it

- If you are allergic (hypersensitive) to medications containing acyclovir or valacyclovir, or any of the other ingredients of Acyclovir 800 STADA.

Before you start to take it

- Please discuss with your doctor or pharmacist before taking Acyclovir 800 Stada.

Taking other medicine

- Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. Probenecid (uricosuric drug for gout) reduces acyclovir excretion via the kidneys, which may increase acyclovir persistence in

the body.

How to take Acyclovir 800 STADA

How much to take

- The instructions given below apply unless your doctor has prescribed Acyclovir 800 Stada otherwise for you.
- Adults:
Shingles (herpes zoster): Take 1 tablet (800mg) of acyclovir 5 times daily every 4 hours during the day. Patients with kidney function impairment, which is particularly prevalent in elderly patients, may need a lower acyclovir dose than indicated above. If you should have kidney problems, your doctor will adjust your dosage to reflect your kidney function.

When to take it

- Swallow the tablets whole with a sufficient amount of liquid (eg. A glass of water) after meals. An adequate fluid intake must be ensured during Acyclovir 800 Stada therapy, especially in patients with impaired kidney function. For best response, Acyclovir 800 Stada therapy should be started as early as possible, ie. As soon as you notice the first signs on your skin.

How long to take it

- Your doctor will determine for how long you should continue Acyclovir 800 Stada therapy. The duration of therapy of herpes zoster is 7 days.

If you forget to take it

- Do not take a double dose to make up for a forgotten dose. Continue your treatment as prescribed. If you have forgotten several doses, or if you have not been taking enough Acyclovir 800 Stada, please consult your doctor.

If you take too much (overdose)

- Intoxication is unlikely after an Acyclovir 800 Stada overdose. A

single dose of 5g of acyclovir produced no signs or symptoms of intoxication. However, no data are available for higher single (over) doses. If you suspect an overdose, should experience pronounced side effects or are in doubt, be sure to discuss with your doctor any concerns you may have. Acyclovir is effectively eliminated by haemodialysis.

While you are using Acyclovir 800 STADA

Things you must do

- Always take Acyclovir 800 Stada exactly as your doctor or pharmacist has told you. You should check with your doctor or pharmacist if you are not sure.

Things you must not do

Please do not:

- 1) Take any other medicines on your own.
- 2) Take probenecid concurrently.

Things to be careful of

- Acyclovir has been detected in breast milk following administration of acyclovir products. Breast-feeding is therefore discouraged while taking Acyclovir 800 Stada.

Side effects

Like all medicines, Acyclovir tablets can cause side effects, although not everybody gets them.

Common side effects are:

- Dizziness
- Skin rashes
- Nausea
- Diarrhea
- Abdominal pain

Should you experience any adverse effects not described in this Insert, please be sure to inform your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug

Acyclovir 800 STADA

Acyclovir 800 mg

Reaction Monitoring by visiting the website npa.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Date of revision

17/09/2020

Serial Number

NPRA(R2/1)201125/565

Storage and Disposal of Acyclovir 800 STADAStorage

- Keep out of the reach and sight of children.
- Do not store above 30°C.

Disposal

- Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product DescriptionWhat it looks like

- Acyclovir 800 STADA is oblong, white with breaking notch on both sides, imprint with VS 3.
- It is available in pack of 35's.

Ingredients

- Active ingredient:
The active substance is: acyclovir.
- Inactive ingredient:
The other ingredients are Colloidal anhydrous silica, Copolyvidone, Microcrystalline Cellulose, Magnesium Stearate, Sodium Starch Glycollate
- MAL number(s):
MAL19984317AZ

Manufacturer

STADA Arzneimittel AG
STADASTRASSE 2-18
D-61118 BAD VILBEL
GERMANY

Product Registration Holder

DKSH Malaysia Sdn Bhd,
B-11-01, The Ascent, Paradigm
No.1, Jalan SS7/26A, Kelana Jaya
47301 Petaling Jaya
Selangor Darul Ehsan