

PACK INSERT FOR MALAYSIA

CYMEVEVE®

500 mg powder for concentrate for solution for infusion

Ganciclovir

1. DESCRIPTION

1.1 Therapeutic/Pharmacologic Class of Drug

Antiviral
ATC code: J05AB06

1.2 Type of Dosage Form

Powder for concentrate for solution for infusion

1.3 Route of Administration

Intravenous (IV) infusion

1.4 Sterile/Radioactive Statement

Sterile product

1.5 Qualitative and Quantitative Composition

Active ingredient: ganciclovir in the form of the sodium salt. Vials containing dry substance equivalent to 500 mg ganciclovir and approximately 43 mg (2 mEq) sodium. The vial content is a white to off-white solid and the constituted solution in water is colourless to light yellow.

2. CLINICAL PARTICULARS

2.1 Therapeutic Indication (s)

Cytomegalovirus retinitis in immunocompromised individuals including patients with acquired immunodeficiency syndrome (AIDS). Prevention of CMV disease in transplant patients at risk for CMV disease. Prevention and treatment of life- or sight-threatening cytomegalovirus (CMV) disease in immunocompromised individuals.

2.2 Dosage and Administration

General

Caution: do not administer by rapid or bolus IV injection! The toxicity of Cymevene may be increased as a result of excessive plasma levels.

Caution: i.m. or s.c. injection may result in severe tissue irritation due to the high pH (-11) of ganciclovir solutions. The recommended dosage, frequency, or infusion rates should not be exceeded. See Section 4.2 (Special Instructions for Use, Handling and Disposal) for details of preparation and handling of Cymevene solution.

Standard dosage for treatment of CMV retinitis

Induction treatment: 5 mg/kg given as an IV infusion over one hour, every 12 hours for 14-21 days in patients with normal renal function. **Maintenance treatment:** 5 mg/kg given as an IV infusion over one hour, once daily on 7 days per week or 6 mg/kg once daily on 5 days per week.

Standard dosage for prevention in transplant recipients

Induction treatment: 5 mg/kg given as an IV infusion over one hour, every 12 hours for 7-14 days in patients with normal renal function. **Maintenance treatment:** 5 mg/kg given as an IV infusion over one hour, once daily on 7 days per week or 6 mg/kg once daily on 5 days per week.

2.2.1 Special dosage instructions

Pediatric patients

Safety and efficacy of ganciclovir in pediatrics have not been established, including use for the treatment of congenital or neonatal CMV infections. The use of Cymevene in children warrants extreme caution due to the potential for long-term carcinogenicity and reproductive toxicity. The benefits of treatment should outweigh the risks (see section 3.2.5 *Pharmacokinetics in Special Populations, Pediatric Population*).

Geriatric patients

No studies have been conducted in adults older than 65 years of age. Since renal clearance decreases with age, Cymevene should be administered to geriatric patients with special consideration of their renal status (see Table 1 and section 3.2.5 *Pharmacokinetics in Special Populations, Geriatric population*).

Patients with renal impairment

For patients with renal impairment, the dose of Cymevene should be modified as shown in the table below.

Table 1 Cymevene dosing for patients with renal impairment receiving mg/kg dosing:

CrCl	Induction dose	Maintenance dose
≥70 ml/min	5.0 mg/kg q12h	5.0 mg/kg/day
50-69 ml/min	2.5 mg/kg q12h	2.5 mg/kg/day
25-49 ml/min	2.5 mg/kg/day	1.25 mg/kg/day
10-24 ml/min	1.25 mg/kg/day	0.625 mg/kg/day
<10 ml/min	1.25 mg/kg 3x/wk after hemodialysis	0.625 mg/kg 3x/wk after hemodialysis

Estimated creatinine clearance can be related to serum creatinine by the following formulae:

For males: $(140 - \text{age}[\text{years}]) \times (\text{bodyweight} [\text{kg}]) / (72) \times (0.011 \times \text{serum creatinine} [\mu\text{mol/l}])$

For females: $0.85 \times \text{male value}$

As dosage modifications are recommended in patients with renal impairment, serum creatinine or estimated creatinine-clearance levels should be monitored carefully.

Hepatic impairment

The safety and efficacy of Cymevene have not been studied in patients with hepatic impairment (see section 3.2.5 *Pharmacokinetics in Special Populations, Hepatic impairment*).

2.3 Contraindications

Cymevene is contraindicated in patients with hypersensitivity to ganciclovir, valganciclovir or to any of the excipients. Breastfeeding (see section 2.5.3 *Lactation*).

2.4 Warnings and Precautions

2.4.1 General

Cross hypersensitivity

Due to the similarity of the chemical structure of ganciclovir and that of aciclovir and penciclovir, a cross-hypersensitivity reaction between these drugs is possible. Caution should therefore be used when prescribing Cymevene to patients with known hypersensitivity to aciclovir or penciclovir (or to their prodrugs, valaciclovir or famciclovir respectively).

Mutagenicity, teratogenicity, carcinogenicity, fertility and contraception

In animal studies ganciclovir was found to be mutagenic, teratogenic, carcinogenic and to impair fertility. Cymevene should therefore be considered a potential teratogen and carcinogen in humans with the potential to cause birth defects and cancers. Prior to initiation of ganciclovir treatment, patients should be advised

of the potential risks to the fetus and to use contraceptive measures. Based on clinical and nonclinical studies, Cymevene may cause temporary or permanent inhibition of spermatogenesis in males (see section 2.5.1, *Females and Males of Reproductive Potential, 2.5.2 Pregnancy, 2.5.3 Lactation, 2.6, Undesirable Effects, 3.3 Preclinical Safety and 4.2, Special Instructions for Use, Handling and Disposal*).

Myelosuppression

Cymevene should be used with caution in patients with pre-existing hematological cytopenia or a history of drug-related hematological cytopenia and in patients receiving radiotherapy. Severe leukopenia, neutropenia, anemia, thrombocytopenia, pancytopenia, bone marrow failure and aplastic anemia have been observed in patients treated with Cymevene. Therapy should not be initiated if the absolute neutrophil count is less than 500 cells/ μL or the platelet count is less than 25,000 cells/ μL or the hemoglobin is less than 8 g/dL (see section 2.6, *Undesirable Effects*).

It is recommended that complete blood counts and platelet counts be monitored in all patients during therapy, particularly in patients with renal impairment (see section 2.2.1 *Special Dosage Instructions*).

In patients with severe leukopenia, neutropenia, anemia and/or thrombocytopenia, that treatment with hematopoietic growth factors and/or the interruption of therapy is recommended (see section 2.6 *Undesirable Effects*).

Use with other medicines

Seizures have been reported in patients taking imipenem-cilastatin and ganciclovir. Cymevene should not be used concomitantly with imipenem-cilastatin unless the potential benefits outweigh the potential risks (see section 2.8 *Interactions with other Medicinal Products and other Forms of Interactions*).

Candida infections including oral candidiasis
Zidovudine and Cymevene each have the potential to cause neutropenia and anemia. Some patients may not tolerate concomitant therapy at full dosage (see section 2.8 *Interactions with other Medicinal Products and other Forms of Interactions*).

Didanosine plasma concentrations may increase during concomitant use with Cymevene; therefore, patients should be closely monitored for didanosine toxicity (see section 2.8 *Interactions with other Medicinal Products and other Forms of Interactions*).

Concomitant use of other drugs that are known to be myelosuppressive or associated with renal impairment with Cymevene may result in added toxicity (see section 2.8 *Interactions with other Medicinal Products and other Forms of Interactions*).

2.4.2 Drug Abuse and Dependence

No information is available for drug abuse and dependence with Cymevene.

2.4.3 Ability to Drive and Use Machines

No studies on the effect on the ability to drive and use machines have been performed. Based on the adverse reaction profile, ganciclovir may have a minor influence on the ability to drive and use machines. Adverse reactions, for example seizures, dizziness and confusion may occur in patients receiving Cymevene (see section 2.6 *Undesirable Effects*). If they occur, such effects may affect tasks requiring alertness including the patient's ability to drive and operate machinery.

2.5 Use in Special Populations

2.5.1 Females and Males of Reproductive Potential

Fertility

In animal studies ganciclovir was found to impair fertility (see section 3.3.3 *Impairment of Fertility*). In a clinical study renal transplant patients receiving Valcyte (which is a pro-drug of Cymevene) for CMV prophylaxis for up to 200 days were compared to an untreated control group. Spermatogenesis was inhibited during treatment with Valcyte. At follow-up, approximately six months after treatment discontinuation, the mean sperm density in treated patients was comparable to that observed in the untreated control group. In Valcyte treated patients, all patients with normal sperm density ($n=7$) and 8/13 patients with low sperm density at baseline, recovered to normal counts after treatment cessation. In the control group, all patients with normal sperm density ($n=6$) and 2/4 patients with low sperm density at baseline, had normal density at the end of follow-up.

Contraception

Women of reproductive potential should be advised to use effective contraception during and for at least 30 days after treatment. Sexually active men are recommended to use condoms during and for at least 90 days after cessation of treatment with Cymevene, unless it is certain that the female partner is not at risk of becoming pregnant (see section 2.4.1 *Warnings and Precautions, General, Mutagenicity, teratogenicity, carcinogenicity, fertility and contraception and 3.3.4 Reproductive Toxicity*).

2.5.2 Pregnancy

In animal studies ganciclovir was associated with reproductive toxicity and teratogenicity (see section 3.3.3 *Impairment of Fertility and 3.3.4 Reproductive Toxicity*). The safety of Cymevene in pregnant women has not been established. However, ganciclovir readily diffuses across the human placenta. The use of Cymevene should be avoided in pregnant women unless the benefit to the mother outweighs the potential risk to the fetus. The safe use of Cymevene during labor and delivery has not been established.

2.5.3 Lactation

Peri- and postnatal development has not been studied with ganciclovir but the possibility of ganciclovir being excreted in breast milk and causing serious adverse reactions in the nursing infant cannot be discounted. Human data are not available, but animal data indicates that ganciclovir is excreted in the milk of lactating rats. Therefore, a decision should be made to discontinue the drug or discontinue nursing taking into consideration the potential benefit of Cymevene to the nursing mother.

2.5.4 Pediatric Use

See sections 2.2.1 *Special Dosage Instructions and 3.2.5 Pharmacokinetics in Special Populations*.

2.5.5 Geriatric Use

See section 2.2.1 *Special Dosage Instructions and 3.2.5 Pharmacokinetics in Special Populations*.

2.5.6 Renal Impairment

In patients with renal impairment, dosage adjustments based on creatinine clearance are required (see section 2.2.1 *Special Dosage Instructions, Patients with renal impairment and 3.2.5 Pharmacokinetics in Special Populations*).

2.5.7 Hepatic Impairment

The safety and efficacy of Cymevene have not been studied in patients with hepatic impairment (see section 2.2.1 *Special Dosage Instructions and 3.2.5 Pharmacokinetics in Special Populations*).

2.6 Undesirable Effects

2.6.1 Clinical Trials

Valganciclovir is a pro-drug of ganciclovir, and adverse reactions associated with valganciclovir can be expected to occur with ganciclovir. Therefore, adverse drug reactions reported with IV or oral ganciclovir (no longer available) or with valganciclovir are included in the table of adverse reactions (see Table 3). In patients treated with ganciclovir/valganciclovir the most serious and frequent adverse drug reactions are hematological reactions

and include neutropenia, anemia and thrombocytopenia. The frequencies presented in the table of adverse reactions are derived from a pooled population of HIV-infected patients ($n=1704$) receiving maintenance therapy with ganciclovir (GAN1697, GAN1653, GAN2304, GAN1774, GAN2226, AVI034, GAN041) or valganciclovir (WV15376, WV15705). Exception is made for agranulocytosis, granulocytopenia and anaphylactic reaction; the frequencies of which are derived from post-marketing experience.

Frequencies are presented as percentages and as CIOMS frequency categories defined as very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1,000$) and very rare ($< 1/10,000$).

The overall safety profile of ganciclovir/valganciclovir is consistent in HIV and transplant populations except that retinal detachment has only been reported in HIV patients with CMV retinitis. However, there are some differences in the frequency of certain reactions.

Intravenous ganciclovir is associated with a lower risk of diarrhea compared to oral valganciclovir. Pyrexia, candida infections, depression, severe neutropenia (ANC $< 500/\mu\text{L}$) and skin reactions are reported more frequently in patients with HIV. Renal and hepatic dysfunction is reported more frequently in organ transplant recipients.

Table 2 Frequency of Ganciclovir/Valganciclovir ADRs Reported in HIV Patients Receiving Maintenance Therapy ($n=1704$):

ADR (MedDRA) System Organ Class	Percentage	Frequency Category
Infections and infestations:		
Candida infections including oral candidiasis	22.42%	Very common
Upper respiratory tract infection	16.26%	
Sepsis	6.92%	Common
Influenza	3.23%	
Urinary tract infection	2.35%	
Cellulitis	1.47%	
Blood and lymphatic system disorders:		
Neutropenia	26.12%	Very common
Anemia	19.89%	Common
Thrombocytopenia	7.34%	Common
Leukopenia	3.93%	
Pancytopenia	1.06%	
Bone marrow failure	0.29%	Uncommon
Aplastic anemia	0.06%	Rare
Agranulocytosis*	0.02%	
Granulocytopenia*	0.02%	
Immune system disorders:		
Hypersensitivity	1.12%	Common
Anaphylactic reaction*	0.02%	Rare
Metabolism and nutrition disorders:		
Decreased appetite	12.09%	Very common
Weight decreased	6.46%	Common
Psychiatric disorders:		
Depression	6.69%	Common
Confusional state	2.99%	
Anxiety	2.64%	
Agitation	0.59%	Uncommon
Psychotic disorder	0.23%	
Thinking abnormal	0.18%	
Hallucinations	0.18%	
Nervous system disorders:		
Headache	17.37%	Very common
Insomnia	7.22%	Common
Neuropathy peripheral	6.16%	
Dizziness	5.52%	
Paraesthesia	3.58%	
Hypoaesthesia	2.58%	
Seizures	2.29%	
Dysgeusia (taste disturbance)	1.35%	
Tremor	0.88%	Uncommon
Eye disorders:		
Visual impairment	7.10%	Common
Retinal detachment**	5.93%	
Vitreous floaters	3.99%	
Eye pain	2.99%	
Conjunctivitis	1.58%	
Macular edema	1.06%	
Ear and labyrinth disorders:		
Ear pain	1.17%	Common
Deafness	0.65%	Uncommon
Cardiac disorders:		
Arrhythmia	0.47%	Uncommon
Vascular disorders:		
Hypotension	2.05%	Common
Respiratory, thoracic and mediastinal disorders:		
Cough	18.31%	Very common
Dyspnoea	11.80%	Common
Gastrointestinal disorders:		
Diarrhea	34.27%	Very common
Nausea	26.35%	Common
Vomiting	14.85%	
Abdominal pain	10.97%	
Dyspepsia	4.81%	Common
Flatulence	4.58%	
Abdominal pain upper	4.58%	
Constipation	3.70%	
Mouth ulceration	3.17%	
Dysphagia	2.93%	
Abdominal distention	2.41%	
Pancreatitis	1.64%	

ADR (MedDRA) System Organ Class	Percentage	Frequency Category
Hepatobiliary disorders:		
Blood alkaline phosphatase increased	3.58%	Common
Hepatic function abnormal	3.23%	
Aspartate aminotransferase increased	1.88%	
Alanine aminotransferase increased	1.23%	
Skin and subcutaneous tissue disorders:		
Dermatitis	11.80%	Very common
Night sweats	7.92%	Common
Pruritus	4.58%	
Rash	2.52%	
Alopecia	1.29%	
Dry skin	0.94%	Uncommon
Urticaria	0.70%	
Musculoskeletal and connective tissue disorders:		
Back pain	4.46%	Common
Myalgia	3.52%	
Arthralgia	3.35%	
Muscle spasms	2.99%	
Renal and urinary disorders:		
Renal impairment	2.52%	Common
Creatinine clearance renal decreased	2.35%	
Blood creatinine increased	1.88%	
Renal failure	0.76%	Uncommon
Hematuria	0.70%	
Reproductive system and breast disorders:		
Infertility male	0.23%	Uncommon
General disorders and administration site conditions:		
Pyrexia	33.51%	Very common
Fatigue	18.96%	Common
Injection site reaction	6.98%	Common
Pain	5.81%	
Chills	5.40%	
Malaise	2.11%	
Asthenia	2.00%	
Chest pain	0.88%	Uncommon

* The frequencies of these adverse reactions are derived from post-marketing experience.
** Retinal detachment has only been reported in studies in HIV patients treated with Cymevene for CMV retinitis.

Description of selected adverse reactions

Neutropenia

The risk of neutropenia is not predictable on the basis of the number of neutrophils before treatment. Neutropenia usually occurs during the first or second week of induction therapy. The cell count usually normalizes within 2 to 5 days after discontinuation of the drug or dose reduction (see section 2.4 *Warnings and Precautions*).

Thrombocytopenia

Patients with low baseline platelet counts ($< 100,000/\mu\text{L}$) have an increased risk of developing thrombocytopenia. Patients with iatrogenic immunosuppression due to treatment with immunosuppressive drugs are at greater risk of thrombocytopenia than patients with HIV (see section 2.4 *Warnings and Precautions*). Severe thrombocytopenia may be associated with potentially life-threatening bleeding.

Laboratory Abnormalities

Laboratory abnormalities in HIV infected patients

Laboratory abnormalities reported from three clinical trials in HIV infected patients receiving intravenous ganciclovir as maintenance treatment for CMV retinitis are listed below in Table 3. One hundred seventy-nine patients were eligible for the laboratory abnormality analysis.

Table 3 Laboratory abnormalities:

Laboratory abnormalities	N=179
Neutropenia (ANC /mm3)	
<500	25.1%
500-750	14.3%
750-1000	26.3%
Anemia (hemoglobin g/dl)	
<6.5	4.6%
6.5-8.0	16.0%
8.0-9.5	25.7%
Thrombocytopenia (platelets/mm3)	
<25,000	2.9%
25,000-50,000	5.1%
50,000-100,000	22.9%
Serum creatinine (mg/dl)	
>2.5	1.7%
1.5-2.5	13.9%

2.6.2 Post-marketing Experience

Safety reports from the post-marketing setting are consistent with safety data from clinical trials with ganciclovir and valganciclovir (see section 2.6.1 *Undesirable Effects* - Table 2).

2.7 Overdose

Overdose experience with intravenous ganciclovir

Reports of overdoses with intravenous ganciclovir, some with fatal outcomes, have been received from clinical trials and during post-marketing experience. In some of these cases no adverse events were reported. The majority of patients experienced one or more of the following adverse events:

- **Hematological toxicity:** myelosuppression including pancytopenia, bone marrow failure, leukopenia, neutropenia, granulocytopenia
 - **Hepatotoxicity:** hepatitis, liver function disorder
 - **Renal toxicity:** worsening of hematuria in a patient with pre-existing renal impairment, acute kidney injury, elevated creatinine
 - **Gastrointestinal toxicity:** abdominal pain, diarrhea, vomiting
 - **Neurotoxicity:** generalised tremor, seizure
- Hemodialysis and hydration may be of benefit in reducing blood plasma levels in patients who receive an overdose of ganciclovir (see 3.2.5 *Pharmacokinetics in Special Populations*).

2.8 Interactions with other Medicinal Products and other Forms of Interactions

Imipenem-cilastatin

Convulsions have been reported in patients who received ganciclovir and imipenem-cilastatin and a pharmacodynamic interaction between these two drugs cannot be discounted. These drugs should not be used concomitantly unless the potential benefits outweigh the risks (see section 2.4.1 *Warnings and Precautions, General, Use with other medicines*).

Potential drug interactions

Toxicity may be enhanced when ganciclovir is co-administered with other drugs known to be myelosuppressive or associated with renal impairment. This includes nucleoside analogues (e.g. zidovudine, didanosine, stavudine), immunosuppressants (e.g. ciclosporin, tacrolimus, mycophenolate mofetil), antineoplastic agents (e.g. doxorubicin, vincristine, vinblastine, hydroxyurea), and anti-infectives (e.g. trimethoprim/sulphonamides, dapsone, amphotericin B, flucytosine, pentamidine). Therefore, these drugs should only be considered for concomitant use with ganciclovir if the potential benefits outweigh the potential risks (see section 2.4.1 *Warnings and Precautions, General, Use with other medicines*).

Zidovudine

Both zidovudine and Cymevene have the potential to cause neutropenia and anemia, a pharmacodynamic interaction may occur during concomitant administration of these drugs some patients may not tolerate concomitant therapy at full dosage (see section 2.4.1 *Warnings and Precautions, General, Use with other medicines*).

Didanosine


Didanosine plasma concentrations were found to be consistently raised when given with IV ganciclovir. At intravenous doses of 5 and 10 mg/kg/day, an increase in the AUC of didanosine ranging from 38 to 67% has been observed confirming a pharmacokinetic interaction during the concomitant administration of these drugs. There was no significant effect on ganciclovir concentrations. Patients should be closely monitored for didanosine toxicity (e.g. pancreatitis) (see section 2.4.1 *Warnings and Precautions, General, Use with other medicines*).

Probencid

Probencid given with oral ganciclovir resulted in statistically decreased renal clearance of ganciclovir (20%) leading to statistically significant increased exposure (40%). These changes were consistent with a mechanism of interaction involving competition for renal tubular excretion. Therefore, patients taking probencid and Cymevene should be closely monitored for ganciclovir toxicity.

3. PHARMACOLOGICAL PROPERTIES AND EFFECTS

3.1 Pharmacodynamic Properties



(N=15) to 4.52 ±2.79 mL/min/kg (N=6) and renal clearance ranged from 2.57 ±0.69 mL/min/kg (N=15) to 3.48 ±0.68 mL/min/kg (N=20), corresponding to 90-101% of administered ganciclovir. Half-lives in patients without renal impairment ranged from 2.73 ±1.29 (N=6) to 3.98 ±1.78 hours (N=8).

3.2.5 Pharmacokinetics in Special Populations

Pediatric Population

Ganciclovir pharmacokinetics were studied in 27 neonates aged 2 to 49 days at intravenous doses of 4 mg/kg (N=14) and 6 mg/kg (N=13). Mean C_{max} was 5.5 ±6 µg/mL and 7.0 ±1.6 µg/mL, the lower and higher dose levels respectively. Mean values for V_{ss} (0.7 L/kg) and systemic clearance (3.15 ±0.47 mL/min/kg at 4 mg/kg and 3.55 ±0.35 mL/min/kg at 6 mg/kg) were comparable to those observed in adults with normal renal function. Ganciclovir pharmacokinetics were also studied in 10 children with normal renal function, aged 9 months to 12 years. The pharmacokinetic characteristics of ganciclovir were the same after single and multiple (q12h) IV doses (5 mg/kg). Exposure as measured by mean AUC₀₋₂₄ on days 1 and 14 were 19.4 ±7.1 and 24.1 ±14.6 µg·h/mL, respectively and the corresponding C_{max} values were 7.59 ±3.21 (day 1) and 8.31 ±4.9 µg/mL (day 14). These range of exposures are comparable to those observed in adults. The steady state volume of distribution after a single dose on day 1 and at the end of the repeat dose period (day 14) was 0.68 ±0.20 L/kg. Systemic clearance for the same study days was 4.66 ±1.72 (day 1) and 4.86 ±2.96 mL/min/kg (day 14). The respective mean values for renal clearance (0 – 12 h) were 3.49 ±2.40 on day 1 and 3.49 ±1.19 mL/min/kg on day 14. The corresponding mean values for the half-life were 2.49 ±0.57 (day 1) and 2.22 ±0.76 h (day 14). The pharmacokinetics of ganciclovir from this study were consistent with those in neonates and adults.

Geriatric population

No ganciclovir pharmacokinetic studies have been conducted in adults older than 65 years of age. However, because ganciclovir is mainly renally excreted and since renal clearance decreases with age a decrease in ganciclovir total body clearance and prolongation of ganciclovir elimination half-life can be anticipated in the elderly (see section 2.2.1 *Special Dosage Instructions, Geriatric patients*).

Renal impairment

The total body clearance of ganciclovir is linearly correlated with creatinine clearance. In patients with mild, moderate, and severe renal impairment, mean systemic clearances of 2.1, 1.0 and 0.3 mL/min/kg were observed. Patients with renal impairment show an increased elimination half-life. In patients with severe renal impairment elimination half-life was increased by 10-fold (see section 2.2.1 *Special Dosage Instructions, Renal impairment*).

Patients undergoing hemodialysis

Plasma concentrations of ganciclovir are reduced by about 50% during a 4 hour hemodialysis session (see section 2.7 *Overdose*). During intermittent hemodialysis, estimates for the clearance of ganciclovir ranged from 42 to 92 mL/min, resulting in intra-dialytic half-lives of 3.3 to 4.5 hours. Estimates of ganciclovir clearance for continuous dialysis were lower (4.0 to 29.6 mL/min) but resulted in greater removal of ganciclovir over a dose interval. For intermittent hemodialysis, the fraction of ganciclovir removed in a single dialysis session varied from 50-63%.

Hepatic impairment

No pharmacokinetic study has been conducted and no population PK data were collected in patients with hepatic impairment undergoing ganciclovir therapy. Hepatic impairment is not anticipated to affect the pharmacokinetics of ganciclovir since ganciclovir is excreted renally (see section 3.2.5 *Pharmacokinetics properties, Elimination*).

3.3 Preclinical Safety

3.3.1 Carcinogenicity

Ganciclovir was mutagenic in mouse lymphoma cells and clastogenic in mammalian cells. Such results are consistent with the positive mouse carcinogenicity study with ganciclovir. Ganciclovir is a potential carcinogen.

3.3.2 Genotoxicity

Ganciclovir was mutagenic in mouse lymphoma cells and clastogenic in mammalian cells.

3.3.3 Impairment of Fertility

Ganciclovir causes impaired fertility and teratogenicity in animals (see section 2.4 *Warnings and Precautions*).

Based upon animal studies where spermia was induced at ganciclovir systemic exposures below therapeutic levels, it is considered likely that ganciclovir could cause temporary or permanent inhibition of human spermatogenesis (see section 2.5.1 *Females and Males of Reproductive Potential, Fertility*).

3.3.4 Reproductive toxicity

Ganciclovir causes teratogenicity in animals.

4. PHARMACEUTICAL PARTICULARS

4.1 Storage

Storage

Do not store above 30°C.

Shelf-life

3 years

This medicine should not be used after the expiry date (EXP) shown on the pack.

Shelf-life and storage of the reconstituted solution

For immediate use after reconstitution

Shelf-life and storage of the infusion solution

Chemical and physical in-use stability of the infusion solution has been demonstrated for 24 hours at 2–8°C. Do not freeze (see section 4.2 *Special Instructions for Use, Handling and Disposal*).

4.2 Special Instructions for Use, Handling and Disposal

Caution should be exercised in the handling of Cymevene.

Since Cymevene is considered a potential teratogen and carcinogen in humans, caution should be observed in its handling (see section 2.4 *Warnings and Precautions*). Avoid inhalation or direct contact of the powder contained in the vial or direct contact of the reconstituted solution with the skin or mucous membranes. Cymevene solutions are alkaline (pH ~11). If such contact occurs, wash thoroughly with soap and water, rinse eyes thoroughly with plain water.

Wearing disposable gloves is recommended during reconstitution and when wiping the outer surface of the vials and the table after reconstitution.

Incompatibilities

Cymevene should not be mixed with other IV products.

Preparation of Cymevene reconstituted solution

1. Lyophilized Cymevene should be reconstituted by injecting 10 mL of sterile water for injection into the vial. Do not use bacteriostatic water for injection containing parabens (para-hydroxybenzoates), since these are incompatible with Cymevene sterile powder and may cause precipitation.
2. The vial should be gently swirled in order to ensure complete wetting of the product. Continue swirling until a clear reconstituted solution is obtained. Reconstituted solution should be inspected for particulate matter prior to proceeding with admixture preparation.

3. From a microbiological point of view, the reconstituted solution should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user.

Preparation of Cymevene infusion solution

Based on patient weight the appropriate calculated dose volume should be removed from the Cymevene vial (concentration 50 mg/mL) and added to an acceptable infusion fluid. Normal saline, dextrose 5% in water, Ringer's or lactated Ringer's solution are determined chemically or physically compatible with Cymevene. Infusion concentrations greater than 10 mg/mL are not recommended. From a microbiological perspective, because Cymevene is reconstituted with non-bacteriostatic sterile water, the infusion solution should be used as soon as possible. If not used immediately in-use storage times and conditions prior to use are the responsibility of the user and should not be longer than 24 hours at 2°C to 8°C.

Disposal of unused/expired medicines

The release of pharmaceuticals in the environment should be minimized. Medicines should not be disposed of via wastewater and disposal through household waste should be avoided. The following points should be strictly adhered to regarding the use and disposal of syringes and other medicinal sharps:

- Needles and syringes should never be reused.
- Place all used needles and syringes into a sharp's container (puncture-proof disposable container).
- Dispose of the full container and of the administration system according to local requirements.

5. PACKAGING AVAILABLE

Vials 500 mg..... 1

Medicine: keep out of reach of children

MYCymevene20220908

Revision date: September 2022

Manufacturer: Valdepharm, Parc Industriel d'Incarville, Parc de la Fringale, CS 10606, 27106 Val de Reuil Cedex, France.

Made for CHEPLAPHARM Arzneimittel GmbH, Germany.