

1. NAME OF THE MEDICINAL PRODUCT

Burinex®
Bumetanide

Carefully read this insert before administering this product. It contains information about your treatment. If you have any doubt or you are not sure about something, please ask your physician or Pharmacist chemist.

Keep this insert as you might need to read it again.

Verify this product fully corresponds to the one prescribed by your physician.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Bumetanide 0.50 mg/ml
Pack size: 5x4ml ampoules

For the full list of excipients, see section 6.1.

Not all pack sizes, strengths and formulations may be marketed

Sale under prescription

3. PHARMACEUTICAL FORM

Solution for injection
Clear colourless solution.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Burinex® is indicated whenever diuretic therapy is required in the treatment of oedema *e.g.* associated with congestive heart failure, cirrhosis of the liver, renal diseases including the nephrotic syndrome. Acute pulmonary oedema, drug induced fluid retention, and drug poisoning that can be treated by forced diuresis. Hypertension.

4.2 Posology and method of administration

Usually 1–2 mg intravenously or intramuscularly.

Pulmonary oedema:

Initially 2 mg by intravenous injection.

This can be repeated, if necessary, after 20 minutes. In those conditions in which an infusion appropriate, 2–5 mg may be given in 500 ml infusion fluid over 30–60 minutes.

Renal failure:

2–10 mg in 500 ml infusion fluid given over a period of 30–60 minutes. Repeated, if required, at intervals of 6–8 hours.

Drug poisoning with salicylates or barbiturates:

Initially, 2 mg intravenously, followed by 1 mg every 4 hours. Totally, 7 mg in the course of 24 hours. The usual procedure for forced alkaline diuresis should be followed. Burinex® injection may be added to the commonly used infusion fluids based on glucose, sodiumchloride, sodium bicarbonate, or potassium chloride.

4.3 Contraindications

- Hypersensitivity to formaldehyde (see Section 4.4).
- Hypersensitivity to active substance or to any of the excipients
- Severe electrolyte depletion
- Persisting anuria
- Hepatic encephalopathy including coma

4.4 Special warnings and precautions for use

Formaldehyde is a degradation product which occurs in the product in trace amounts during storage. Due to the anaphylactic potential of formaldehyde, caution should be taken.

Toxic epidermal necrolysis (TEN) and Stevens-Johnson syndrome (SJS), which can be life-threatening or fatal, have been reported in relation to non-antibiotic sulphonamide containing products, including bumetanide. Patients should be advised of the signs and symptoms of SJS and TEN. If SJS or TEN is suspected, bumetanide should be withdrawn immediately. If the patient has developed SJS or TEN, treatment with bumetanide must not be restarted in this patient at any time.

Caution is advised if bumetanide is to be administered to patients with severe hepatic impairment. Caution should be exercised when bumetanide is used in patients with hypotension. Electrolyte and fluid imbalance may occur (see section 4.8) and replacement therapy should be instituted where indicated. Serum potassium concentrations should be monitored regularly.

Administration of proton pump inhibitors has been associated with development of hypomagnesaemia. Hypomagnesaemia may be exacerbated with co-administration of Burinex® and particular attention to magnesium levels should be given when this combination is used.

As with other diuretics, bumetanide may cause an increase in blood uric acid.

Bumetanide should be used with caution in patients with potential obstruction of the urinary tract.

Caution is advised if bumetanide is to be administered to patients with severe or progressive renal impairment or with elevated urea/Blood Urea Nitrogen (BUN) or creatinine. Periodic monitoring of urine and blood glucose should be made in diabetics and patients suspected of latent diabetes.

If known hypersensitivity to sulphonamides there may be a potential risk of hypersensitivity to bumetanide.

Bumetanide found in urine by doping test is cause for disqualification of athletes.

4.5 Interaction with other medicinal products and other forms of interaction

Digitalis glycosides

Hypokalaemia increases the sensitivity to digitalis glycosides which might result in digitalis toxicity (nausea, vomiting, and arrhythmias). Potassium level and signs for digitalis toxicity should be monitored. Potassium supplementation and lower digitalis glycoside dose should be considered.

Non-depolarising neuromuscular blocking agents

Hypokalaemia increases the sensitivity to non-depolarising neuromuscular blocking agents.

Lithium

Bumetanide reduces lithium clearance resulting in high serum levels of lithium, therefore concomitant therapy requires close monitoring of serum lithium levels. Lower lithium doses may be required.

Antiarrhythmics

Concomitant use of bumetanide and class III antiarrhythmic drugs may result in increased risk of electrolyte imbalance and subsequent cardiotoxicity (QT prolongation, torsades de pointes, cardiac arrest). Patients' electrolyte levels should be monitored as should symptoms of arrhythmias.

NSAIDs

Non-steroidal anti-inflammatory drugs (NSAID) inhibit the effect of bumetanide. The effects of concurrent use should be monitored (e.g. blood pressure, signs of renal failure). Diuretics may enhance the nephrotoxicity of NSAIDs.

Antihypertensive agents and medicinal products inducing postural hypotension

Bumetanide may potentiate the effect of antihypertensive agents including diuretics and drugs inducing postural hypotension (e.g. tricyclic antidepressants). First-dose hypotension may occur.

Potassium depleting agents

The potassium depleting effect of bumetanide may be increased by other potassium depleting agents.

Aminoglycosides

The ototoxic effects of aminoglycosides may be increased by concomitant administration of potent diuretics such as bumetanide.

Probenecid

Probenecid inhibits the renal tubular secretion of bumetanide leading to a diminished natriuresis.

4.6 Fertility, pregnancy and lactation

Pregnancy

Bumetanide may cause harmful pharmacological effects during pregnancy, to the foetus or to the newborn child. Burinex® should not be used during pregnancy unless the clinical condition of the

woman requires treatment with bumetanide. It may be used only in case of heart failure when the potential benefit justifies the potential risk to the foetus.

Breastfeeding

Bumetanide should not be used during breastfeeding.

Fertility

There are no clinical studies with bumetanide regarding fertility.

4.7 Effects on ability to drive and use machines

Bumetanide has no or negligible direct influence on the ability to drive and use machines. However, the patient should be informed that dizziness may occur during treatment and take this into account while driving or using machines.

4.8 Undesirable effects

The estimation of the frequency of undesirable effects is based on a pooled analysis of data from clinical studies and spontaneous reporting.

Based on pooled data from clinical studies including more than 1000 patients who received bumetanide, approximately 12 % of patients can be expected to experience an undesirable effect.

The most frequently reported adverse reactions during treatment are headache and electrolyte imbalance (including hypokalaemia, hyponatraemia, hypochloraemia and hyperkalaemia) occurring in approximately 4% of the patients, followed by dizziness (including orthostatic hypotension and vertigo) and fatigue occurring in approximately 3% of patients.

Electrolyte disturbances can occur especially during long term treatment.

Renal failure has been reported in post-marketing safety surveillance.

Severe cutaneous adverse reactions (SCARs), including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), have been reported in association with bumetanide (see section 4.4).

Undesirable effects are listed by MedDRA system organ class (SOC) and the individual undesirable effects are listed starting with the most frequently reported. Within each frequency grouping, adverse reactions are presented in the order of decreasing seriousness.

Very common $\geq 1/10$

Common $\geq 1/100$ and $< 1/10$

Uncommon $\geq 1/1,000$ and $< 1/100$

Rare $\geq 1/10,000$ and $< 1/1,000$

Very rare $< 1/10,000$

Not known (cannot be estimated from the available data)

Blood and lymphatic system disorders	
Uncommon ($\geq 1/1,000$ and $< 1/100$)	Bone marrow failure and pancytopenia Thrombocytopenia Leukopenia including neutropenia Anaemia
Metabolism and nutrition disorders	
Common: ($\geq 1/100$ and $< 1/10$)	Electrolyte imbalance (including hypokalaemia, hyponatraemia, hypochloraemia and hyperkalaemia)

Uncommon: ($\geq 1/1,000$ and $< 1/100$)	Dehydration Glucose metabolism disorder Hyperuricaemia and gout
Nervous system disorders	
Common: ($\geq 1/100$ and $< 1/10$)	Dizziness (including orthostatic hypotension and vertigo) Fatigue (including lethargy, somnolence, asthenia and malaise) Headache
Uncommon: ($\geq 1/1,000$ and $< 1/100$)	Syncope
Ear and labyrinth disorders	
Uncommon: ($\geq 1/1,000$ and $< 1/100$)	Hearing disturbances
Cardiac disorders	
Uncommon ($\geq 1/1,000$ and $< 1/100$)	Chest pain and discomfort
Vascular disorders	
Uncommon: ($\geq 1/1,000$ and $< 1/100$)	Hypotension
Respiratory, thoracic and mediastinal disorders	
Uncommon: ($\geq 1/1,000$ and $< 1/100$)	Dyspnoea Cough
Gastrointestinal disorders	
Common: ($\geq 1/100$ and $< 1/10$)	Abdominal pain and discomfort Nausea
Uncommon: ($\geq 1/1,000$ and $< 1/100$)	Vomiting Diarrhoea Constipation Dry mouth and thirst
Skin and subcutaneous tissue disorders	
Uncommon: ($\geq 1/1,000$ and $< 1/100$)	Rash* Dermatitis and eczema Urticaria Pruritus Photosensitivity *Various types of rash reactions such as erythematous, maculo-papular and pustular have been reported
Not known (cannot be estimated from the available data)	Stevens-Johnson syndrome (SJS), Toxic epidermal necrolysis (TEN)
Musculoskeletal and connective tissue disorders	
Common: ($\geq 1/100$ and $< 1/10$)	Muscle spasms Pain and myalgia
Renal and urinary disorders	
Common: ($\geq 1/100$ and $< 1/10$)	Micturition disorder
Uncommon: ($\geq 1/1,000$ and $< 1/100$)	Renal impairment (including renal failure)
General disorders and administration site conditions	
Uncommon: ($\geq 1/1,000$ and $< 1/100$)	Oedema peripheral

Paediatric population

The safety profile of Burinex® has not been established in the paediatric population.

4.9 Overdose

In high doses and during long-term treatment loop diuretics may cause electrolyte imbalance, dehydration and polyuria.

Symptoms of electrolyte imbalance include dry mouth, thirst, weakness, lethargy, drowsiness, confusion, gastrointestinal disturbances, restlessness, muscle pain and cramps and seizures.

Treatment is by adjustment of the fluid and electrolyte imbalance.

5. PHARMACOLOGICAL PROPERTIES

5.0 Therapeutic classification

C 03 CA 02 - Sulfonamides, plain

5.1 Pharmacodynamic properties

Bumetanide is a potent high ceiling loop diuretic.

Bumetanide exerts an inhibiting effect on the reabsorption mechanism of salts in the thick ascending limb of Henle and in the renal proximal tubules.

Bumetanide hereby exerts a diuretic and natriuretic action.

5.2 Pharmacokinetic properties

After intravenous administration, diuresis occurs within a few minutes and usually ceases after approx. 2 hours. Bumetanide is eliminated with a half-life between 1 to 2 hours. In patients with hepatic or renal diseases the elimination half-life is prolonged. Bumetanide is strongly bound to plasma proteins and renal excretion accounts for about half of the total clearance. The hepatic metabolism and biliary excretion accounts for the other half. No active metabolites have been found. Burinex has a steep dose response curve.

5.3 Preclinical safety data

Bumetanide has shown no mutagenic, teratogenic or carcinogenic effects in the preclinical studies although data from investigative preclinical studies *in vitro* and *in vivo* suggest a possible effect on pre- and postnatal kidney, lung and neurogenic development. Non-clinical data reveal no special hazard for humans at the recommended therapeutic dose based on conventional studies of acute, subacute and repeated dose toxicity.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

- Disodium phosphate dihydrate
- Sodium dihydrogen phosphate dihydrate
- Xylitol
- Sterile water

6.2 Shelf life

3 years.

6.3 Special precautions for storage

Store below 25°C.

Keep the ampoule in the outer carton to protect from light.

Keep medicines out of reach of children.

7. MANUFACTURER

CENEXI, 52, rue Marcel et Jacques Gaucher
94120 Fontenay-sous Bois, France

For

Karo Pharma AB
Box 16184
103 24 Stockholm
Sweden

This leaflet was last revised in Apr 2025