

SYMBICORT TURBUHALER®

80/4.5 MCG/DOSE, 160/4.5 MCG/DOSE

Budesonide/formoterol fumarate dihydrate (80/4.5 mcg/dose)

Budesonide/formoterol fumarate dihydrate (160/4.5 mcg/dose)

What is in this leaflet

1. What Symbicort Turbuhaler is used for
2. How Symbicort Turbuhaler works
3. Before you use Symbicort Turbuhaler
4. How to use Symbicort Turbuhaler
5. While you are using it
6. Side effects
7. Storage & Disposal of Symbicort Turbuhaler
8. Product Description
9. Manufacturer
10. Product Registration Holder
11. Date of revision
12. Serial number
13. Preparing your new Symbicort Turbuhaler

What Symbicort Turbuhaler is used for

Symbicort Turbuhaler is an inhaler that is used to treat asthma in adults, adolescents aged 12-17 years and children aged 6 years and above. It is also used to treat the symptoms of Chronic Obstructive Pulmonary Disease (COPD) in adults aged 18 years and older.

Asthma

Symbicort Turbuhaler can be prescribed for asthma in two different ways.

- Some people can take Symbicort Turbuhaler when they need it – they use Symbicort Turbuhaler as an anti-inflammatory “reliever” to treat their symptoms when their asthma gets worse and to help prevent symptoms from happening (eg before exercise or exposure to other triggers such as allergens).
- Some people need to take Symbicort Turbuhaler everyday – they use their Symbicort Turbuhaler as a

daily maintenance preventer to help maintain control of their asthma symptoms and help prevent asthma attacks.

Chronic obstructive pulmonary disease (COPD)

Symbicort Turbuhaler can also be used to treat the symptoms of COPD in adults. COPD is a long-term disease of the airways in the lungs, which is often caused by cigarette smoking.

How Symbicort Turbuhaler works

Symbicort Turbuhaler is an inhaler. It contains two different medicines: budesonide and formoterol fumarate dihydrate.

- Budesonide belongs to a group of medicines called ‘corticosteroids’. It works by reducing and preventing swelling and inflammation in your lungs.
- Formoterol fumarate dihydrate belongs to a group of medicines called ‘long-acting β_2 adrenoceptor agonists’ or ‘bronchodilators’. It works by relaxing the muscles in your airways. This helps you to breathe more easily.

Before you use Symbicort Turbuhaler

- When you must not use it

Do not use Symbicort Turbuhaler if:

- You are allergic (hypersensitive) to budesonide, formoterol, or the other ingredient of this medicine (listed in section ‘Ingredients’), which is lactose (which contains small amounts of milk protein).

- Before you start to use it

Before you use Symbicort Turbuhaler, tell your doctor or pharmacist if:

- You are diabetic.
- You have a lung infection.
- You have high blood pressure or you have ever had a heart problem (including an uneven heart beat, a very fast pulse, narrowing of the arteries or heart failure).
- You have problems with your thyroid or adrenal glands.
- You have low levels of potassium in your blood.
- You have severe liver problems.

Contact your doctor if you experience blurred vision or other visual disturbances.

- Taking other medicines

Please tell your doctor or pharmacist if you are using, have recently used or might use any other medicines.

In particular, tell your doctor or pharmacist if you are using any of the following medicines:

- Beta-blocker medicines (such as atenolol or propranolol for high blood pressure), including eyedrops (such as timolol for glaucoma)
- Medicines for a fast or uneven heart beat (such as quinidine)
- Medicines like digoxin, often used to treat heart failure
- Diuretics, also known as ‘water tablets’ (such as furosemide). These are used to treat high blood pressure.
- Steroid medicines that you take by mouth (such as prednisolone)
- Xanthine medicines (such as theophylline or

- aminophylline). These are often used to treat asthma
- Other bronchodilators (such as salbutamol)
 - Tricyclic anti-depressants (such as amitriptyline) and the anti-depressant nefazodone
 - Phenothiazine medicines (such as chlorpromazine and prochlorperazine)
 - Medicines called 'HIV-protease inhibitors' (such as ritonavir) to treat HIV infection.
 - Medicines to treat infections (such as ketoconazole, itraconazole, voriconazole, posaconazole, clarithromycin and telithromycin)
 - Medicines for Parkinson's disease (such as leva-dopa)
 - Medicines for thyroid problems (such as levo-thyroxine)

If any of the above applies to you, or if you are not sure, talk to your doctor or pharmacist before using Symbicort Turbuhaler.

Also tell your doctor or pharmacist if you are going to have a general anaesthetic for an operation or for dental work.

How to use Symbicort Turbuhaler

- How much to use & when to use it

- Always use Symbicort Turbuhaler exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.
- It is important to use Symbicort Turbuhaler every day, even if you have no asthma or COPD symptoms at the time.
- If you are using Symbicort Turbuhaler for asthma, your doctor will want to regularly check your symptoms.

If you have been taking steroid tablets for your asthma or COPD, your doctor may reduce the number of tablets that you take, once you start to use Symbicort Turbuhaler. If you have been taking oral steroid tablets for a long time, your doctor may want you to have blood tests from time to time. When reducing oral steroid tablets, you may feel generally unwell even though your chest symptoms may be improving. You might experience symptoms such as a stuffy or runny nose, weakness or joint or muscle pain and rash (eczema). If any of these symptoms bother you, or if symptoms such as headache, tiredness, nausea (feeling sick) or vomiting (being sick) occur, please contact your doctor immediately. You may need to take other medication if you develop allergic or arthritic symptoms. You should speak to your doctor if you are concerned as to whether you should continue to use Symbicort Turbuhaler.

Your doctor may consider adding steroid tablets to your usual treatment during periods of stress (for example, when you have a chest infection or before an operation).

Important information about your asthma or COPD symptoms

If you feel you are getting breathless or wheezy while using Symbicort Turbuhaler, you should continue to use Symbicort Turbuhaler but go to see your doctor as soon as possible, as you may need additional treatment.

Contact your doctor immediately if:

- Your breathing is getting worse or you often wake up at night with asthma.
- Your chest starts to feel tight in the morning or your chest tightness lasts longer than usual.

These signs could mean that your asthma or COPD is not being properly controlled and you may need different or additional treatment immediately.

Asthma

Your doctor may have prescribed Symbicort Turbuhaler for you to use as:

- a) An anti-inflammatory reliever medicine only or,
- b) Both an anti-inflammatory reliever and a daily maintenance preventer medicine or,
- c) As a daily maintenance preventer only, where another medicine is used as a reliever.

Ask your doctor if you have any questions about how you should be using your Symbicort Turbuhaler.

If your asthma has been under control for some time, your doctor may tell you to take less inhalations of Symbicort Turbuhaler, prescribe you a lower strength of Symbicort Turbuhaler or recommended that you use Symbicort Turbuhaler in a different way.

If you are using more inhalations of your reliever medicine or you are wheezing or breathless more than usual tell your doctor as your asthma may be getting worse.

a) Anti-inflammatory reliever only

For Symbicort Turbuhaler 160/4.5 mcg/dose

Patients aged 12 years and over, Symbicort Turbuhaler 160/4.5 mcg/dose can be used to treat asthma symptoms when they happen and to help stop asthma symptoms from happening (e.g. just before exercise or before you get exposed to other triggers).

If you get asthma symptoms, take 1 inhalation and wait a few minutes. If you do not feel better, take another inhalation.

Your doctor will tell you how many inhalations to take before exercising or exposure to other triggers to help stop symptoms from happening.

Do not use more than 6 inhalations on a single occasion or more than 12 inhalations in any day. If your symptoms continue to worsen over 3 days, despite using additional inhalations, tell your doctor.

Have your Symbicort Turbuhaler reliever with you at all times.

Symbicort Turbuhaler 80/4.5 mcg/dose should NOT be used as Symbicort Anti-inflammatory Reliever Therapy.

Symbicort Anti-inflammatory Reliever Therapy is NOT recommended for children below 12 years of age.

b) Anti-inflammatory reliever plus maintenance therapy

Only use Symbicort Turbuhaler in this way if your doctor has told you to and if you are aged 12 or above.

Use Symbicort Turbuhaler 160/4.5 mcg/dose or 80/4.5 mcg/dose as a 'reliever inhaler' to treat asthma symptoms when they happen.

- If you get asthma symptoms, take 1 inhalation and wait a few minutes.
- If you do not feel better, take another inhalation.
- Your doctor will tell you how many inhalations of Symbicort Turbuhaler to take before exercising or exposure to other triggers to help stop symptoms from happening.
- Do not take more than 6 inhalations at a single time.

Symbicort Turbuhaler 160/4.5 mcg/dose can also be used to help stop asthma symptoms

from happening (e.g. just before exercise or before you get exposed to other triggers).

In addition, Symbicort Turbuhaler 160/4.5 mcg/dose or 80/4.5 mcg/dose should be taken every day. You can take:

- 1 inhalation in the morning **and** 1 inhalation in the evening
- or
- 2 inhalations in the morning
- or
- 2 inhalations in the evening.

For Symbicort Turbuhaler 160/4.5 mcg/dose, your doctor may increase this to 2 inhalations twice a day.

Always keep your Symbicort Turbuhaler Inhaler with you, so you can use it when you need it.

A total daily dose of more than 8 inhalations is not normally needed. However, your doctor may allow you to take up to 12 inhalations a day for a limited period.

If you regularly need to use 8 or more inhalations a day, make an appointment to see your doctor or nurse. They may need to change your treatment.

Do not use more than 12 inhalations in total in 24 hours.

c) Daily fixed dose maintenance therapy

Use your Symbicort Turbuhaler every day. This helps to prevent asthma symptoms from happening.

For Symbicort Turbuhaler 160/4.5 mcg/dose

Adults (18 years and above)

- The usual dose is 1 or 2 inhalations, twice a day.
- Your doctor may increase this to 4 inhalations, twice a day.

Adolescents (12 to 17 years)

- The usual dose is 1 or 2 inhalations, twice a day.

If your symptoms are well controlled, your doctor may ask you to take your medicine once a day.

For Symbicort Turbuhaler 80/4.5 mcg/dose

Adults (18 years and above)

- The usual dose is 1 or 2 inhalations, twice a day.
- Your doctor may increase this to 4 inhalations, twice a day.

Adolescents (12 to 17 years)

- The usual dose is 1 or 2 inhalations, twice a day

Children (6 years and older)

- The usual dose is 2 inhalations, twice a day.

If your symptoms are well controlled, your doctor may ask you to take your medicine once a day.

Symbicort Turbuhaler is not recommended to be used in children who are younger than 6 years.

Your doctor (or asthma nurse) will help you to manage your asthma. They will adjust the dose of this medicine to the lowest dose that controls your asthma. However, do not adjust the dose without talking to your doctor (or asthma nurse) first.

Use your separate 'reliever inhaler' to treat asthma symptoms when they happen.

Always keep your 'reliever inhaler' with you to use when you need it. Do not use Symbicort Turbuhaler to treat asthma symptoms - use your reliever inhaler.

Chronic Obstructive Pulmonary Disease (COPD)

For Symbicort Turbuhaler 160/4.5 mcg/dose

- Only to be used by adults (aged 18 years and above).
- The usual dose is 2 inhalations twice a day.

Your doctor may also prescribe other bronchodilator drugs, for example anticholinergics (such as tiotropium or ipratropium bromide) for your COPD disease.

Symbicort Turbuhaler 80/4.5 mcg/dose is NOT indicated for symptomatic treatment of COPD.

- How long to use it

- Always use Symbicort Turbuhaler exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.
- It is important to use Symbicort Turbuhaler every day, even if you have no asthma or COPD symptoms at the time.
- If you are using Symbicort Turbuhaler for asthma, your doctor will want to regularly check your symptoms.

- If you forget to use it

- If you forget to take a dose, take it as soon as you remember. However, if it is nearly time for your next dose, skip the missed dose.
- Do **not** take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

- If you use too much (overdose)

It is important that you take your dose as stated on the pharmacist's label or as advised by your doctor. You should not exceed your prescribed dose without seeking medical advice.

The most common symptoms that may occur after if you use more Symbicort Turbuhaler than you should are trembling, headache or a rapid heart beat.

While you are using it

- Things you must do

Use Symbicort Turbuhaler as directed by your doctor.

- Things you must not do

Do not stop using Symbicort Turbuhaler unless advised by your doctor.

- Things to be careful of

Pregnancy and breast-feeding

- If you are pregnant, or planning to get pregnant, talk to your doctor before using Symbicort Turbuhaler - do not use Symbicort Turbuhaler unless your doctor tells you to.
- If you get pregnant while using Symbicort Turbuhaler, do not stop using Symbicort Turbuhaler but talk to your doctor immediately.
- If you are breast-feeding, talk to your doctor before using Symbicort Turbuhaler.

Driving and using machines

Symbicort Turbuhaler has no or negligible effect on your ability to drive or to use tools or machines.

Symbicort Turbuhaler contains lactose

Symbicort Turbuhaler contains lactose, which is a type of sugar. If you have been told by your doctor that you have an intolerance to some sugars, talk to your doctor before using this medicine. The amount of lactose in this medicine does not normally cause problems in people who are lactose intolerant.

The excipient lactose contains small amounts of milk proteins, which may cause allergic reaction.

Side effects

Like all medicines, Symbicort Turbuhaler can cause side effects, although not everybody gets them.

If either of the following happens to you, stop using Symbicort Turbuhaler and talk to your doctor immediately:

- Swelling of your face, particularly around your mouth (tongue and/or throat and/or difficulty to swallow) or hives together with difficulties to breath (angioedema) and or sudden feeling of faintness. This may mean that you are having an allergic reaction. This happens rarely, affecting less than 1 in 1,000 people.
- Sudden acute wheezing or shortness of breath immediately after using your inhaler. **If either of these symptoms occur, stop using your Symbicort Turbuhaler Inhaler straightaway and use your 'reliever' inhaler. Contact your doctor immediately as you may need to have your treatment changed.** This happens very rarely, affecting less than 1 in 10,000 people.

Other possible side effects:

Common (may affect up to 1 in 10 people)

- Palpitations (awareness of your heart beating), trembling or shaking. If these effects occur, they are usually mild and usually disappear as you continue to use Symbicort Turbuhaler
- Thrush (a fungal infection) in the mouth. This is less likely if you rinse your mouth out with water after using your Symbicort Turbuhaler.
- Mild sore throat, coughing and a hoarse voice
- Headache

- Pneumonia (infection of the lung) in COPD patients

Tell your doctor if you have any of the following while taking Symbicort Turbuhaler, they could be symptoms of a lung infection:

- Fever or chills
- Increased mucus production, change in mucus colour
- Increased cough or increased breathing difficulties

Uncommon (may affect up to 1 in 100 people)

- Feeling restless, nervous or agitated
- Disturbed sleep
- Feeling dizzy
- Nausea (feeling sick).
- Fast heart beat
- Bruising of the skin
- Muscle cramps
- Blurred vision

Rare (may affect up to 1 in 1,000 people)

- Rash, itching
- Bronchospasm (tightening of the muscles in the airways which causes wheezing). If the wheezing comes on suddenly after using Symbicort Turbuhaler stop using Symbicort Turbuhaler and talk to your doctor immediately
- Low levels of potassium in your blood
- Uneven heart beat

Very rare (may affect up to 1 in 10,000 people)

- Depression
- Changes in behaviour, especially in children
- Chest pain or tightness in the chest (angina pectoris)
- An increase in the amount of sugar (glucose) in your blood
- Taste changes, such as an unpleasant taste in the mouth
- Changes in your blood pressure

Inhaled corticosteroids can affect the normal production of steroid hormones in your body, particularly if you use high doses for a long time. The effects include:

- Changes in bone mineral density (thinning of the bones)
- Cataract (clouding of the lens in the eye)
- Glaucoma (increased pressure in the eye)
- A slowing of the rate of growth of children and adolescents
- An effect on the adrenal gland (a small gland next to the kidney)

These effects are much less likely to happen with inhaled corticosteroids than with corticosteroid tablets.

If any of the side effects get serious or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my (Consumer → Reporting Side Effects to Medicines (ConSERF or Vaccines (AEFI))).

Storage & Disposal of Symbicort Turbuhaler

- Storage

- Keep this medicine out of the reach and sight of children.
- Do not use the inhaler after the expiry date which is stated on the carton or on the label of your inhaler after EXP. The expiry date refers to the last day of that month.
- Do not store above 30°C.
- Keep the container/cap tightly closed, in order to protect from moisture.

- Disposal

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to dispose of medicines you no longer use. These measurements will help protect the environment.

Product Description

- What it looks like

Symbicort Turbuhaler is an inhaler containing your medicine. The inhalation powder is white in colour. Each inhaler contains either 30, 60 or 120 doses and has a white body with a red turning grip. The turning grip contains a Braille code with the number 6 for identification, to differ from other AstraZeneca inhaled products.

Symbicort Turbuhaler is available in packs of 1 inhaler containing 30, 60 or 120 doses.

- Ingredients

- **Active Ingredients**
The active substances are budesonide and formoterol fumarate dihydrate.

Each inhaled dose of Symbicort Turbuhaler 80/4.5 mcg/dose contains 80 micrograms of budesonide and 4.5 micrograms of formoterol fumarate dihydrate.

Each inhaled dose of Symbicort Turbuhaler 160/4.5 mcg/dose contains 160 micrograms of budesonide and 4.5 micrograms of formoterol fumarate dihydrate.

- **Inactive Ingredients**
The other ingredient is lactose monohydrate (which contains milk proteins).

- MAL number:

Symbicort Turbuhaler
160/4.5 mcg/dose
MAL20020709AZ

Symbicort Turbuhaler 80/4.5 mcg/dose
MAL20020710AZ

Manufacturer

AstraZeneca AB
Forskargatan 18
SE-151 36 Södertälje
Sweden.

Product Registration Holder

AstraZeneca Sdn. Bhd.
Level 11 & 12, The
Bousteador, No. 10, Jalan PJU
7/6, Mutiara Damansara,
47800 Petaling Jaya, Selangor,
Malaysia.

Date of revision

21/11/2024
VV-RIM-01448652
(based on PI VV-RIM-
01325129)

Serial number

NPRA (R1/4) 20112024/212

Preparing your new Symbicort Turbuhaler

Before using your **new** Symbicort Turbuhaler Inhaler **for the first time**, you need to prepare it for use as follows:

- Unscrew the cover and lift it off. You may hear a rattling sound.
- Hold your Symbicort Turbuhaler Inhaler upright with the red grip at the bottom.
- Turn the red grip as far as it will go in one direction. Then turn it as far as it will go in the other direction (it does not matter which way you turn it first). You should hear a click sound.
- Do this again, turning the red grip in both directions.
- Your Symbicort Turbuhaler Inhaler is now ready for use.

How to take an inhalation

Every time you need to take an inhalation, follow the instructions below.

1. Unscrew the cover and lift it off. You may hear a rattling sound.
2. **Hold your Symbicort Turbuhaler Inhaler upright** with the red grip at the bottom.



3. Do not hold the mouthpiece when you load your Symbicort Turbuhaler Inhaler. To load your Symbicort Turbuhaler Inhaler with a dose, turn the red grip as far as it will go in one direction.



Then turn it as far as it will go in the other direction (it does not matter which way you turn it first). You should hear a click sound. Your Symbicort Turbuhaler Inhaler is now loaded and ready to use. Only load your Symbicort Turbuhaler Inhaler when you need to use it.

4. Hold your Symbicort Turbuhaler Inhaler away from your mouth. Breathe out gently (as far as is comfortable). Do not breathe out through your Symbicort Turbuhaler Inhaler.
5. Place the mouthpiece gently between your teeth. Close your lips. Breathe in as deeply and as hard as you can through your mouth. Do not chew or bite on the mouthpiece.



6. **Remove your Symbicort Turbuhaler Inhaler from your mouth. Then breathe out gently.** The amount of medicine that is inhaled is very small. This means you may not be able to taste it after inhalation. If you have

followed the instructions, you can still be confident that you have inhaled the dose and the medicine is now in your lungs.

7. If you are to take a second inhalation, repeat steps 2 to 6.
8. Replace the cover tightly after use.



9. Rinse your mouth with water after your daily morning and/or evening doses, and spit it out.

Do not try to remove or twist the mouthpiece. It is fixed to your Symbicort Turbuhaler Inhaler and must not be taken off. Do not use your Symbicort Turbuhaler Inhaler if it has been damaged or if the mouthpiece has come apart from your Symbicort Turbuhaler Inhaler.

As with all inhalers, caregivers should ensure that children prescribed Symbicort Turbuhaler use correct inhalation technique, as described above.

Cleaning your Symbicort Turbuhaler Inhaler

Wipe the outside of the mouthpiece once a week with a dry tissue. Do not use water or liquids.

When to start using a new inhaler

- The dose indicator tells you how many doses (inhalations) are left in your Symbicort Turbuhaler Inhaler, starting with either 30, 60 or 120 doses when it is full.



- The dose indicator is marked in intervals of 10 doses. Therefore it does not show every dose.

- When you first see a red mark at the edge of the indicator window, there are approximately 20 doses left. For the last 10 doses, the background of the dose indicator is red. When the '0' on the red background has reached the middle of the window, you must start using your new Symbicort Turbuhaler Inhaler.

Note:

- The grip will still twist and 'click' even when your Symbicort Turbuhaler Inhaler is empty.
- The sound that you hear as you shake your Symbicort Turbuhaler Inhaler is produced by a drying agent and not the medicine. Therefore the sound does not tell you how much medicine is left in your Symbicort Turbuhaler Inhaler.
- If you load your Symbicort Turbuhaler Inhaler more than once by mistake before taking your dose, you will still only receive one dose. However, the dose indicator will register all the loaded doses.